center for PHILOSOPHY OF SCIENCE

home

- ::: [about](#)
- ::: [news](#)
- ::: [links](#)
- ::: [giving](#)
- ::: [contact](#)

events

- ::: [calendar](#)
- ::: [lunchtime](#)
- ::: [annual lecture series](#)
- ::: [conferences](#)

people

- ::: [visiting fellows](#)
- ::: [resident fellows](#)
- ::: [associates](#)

joining

- ::: [visiting fellowships](#)
- ::: [resident fellowships](#)
- ::: [associateships](#)

being here

- ::: [visiting](#)
- ::: [the last donut](#)
- ::: [photo album](#)

[::: center home](#) >> [events](#) >> [conferences](#) >> [other](#) >> [2007-08](#) >> [&HPS](#)

The Discovery of Argon: A Case for Error-Statistical Reasoning

Aris Spanos

Virginia Tech,

Lord Rayleigh and Sir William Ramsay discovered the inert gas 'argon' in atmospheric air in 1895, after a series of experiments involving the production of nitrogen using two different procedures, broadly defined as 'atmospheric' and 'chemical'. The discovery process was based on a carefully designed sequence of experiments combined with an informal (by today's standards) analysis of the resulting data. This combination guided them to further, and further experiments until they were able to establish the existence of argon in the atmospheric air, by eliminating other possible explanations. The basic objective of the paper is twofold. First, to capture the sound scientific intuition underlying their reasoning and their pragmatic attempts to address Duhemian problems in their efforts to explain the source of the initial discrepancy. Second, to formalize their 'reasoning from error' which led them to infer the existence of argon in the atmospheric air and eventually to the discovery of the other noble gases. It is argued that the error statistical account proposed by Mayo (1996) provides an appropriate framework for accomplishing both objectives.

*I would like to thank Alan Chalmers for invaluable advice on the history of this episode.

I am

grateful to Deborah Mayo for numerous suggestions that helped to improve the paper.

1

1 Introduction

Lord Rayleigh (1842-1919) was awarded the 1904 Nobel Prize in physics and his collaborator, Sir William Ramsay (1852-1916), was awarded the 1904 Nobel Prize in chemistry, for their role in the discovery of argon, an inert gas in the atmosphere. The discovery of argon was of paramount importance for Chemistry because it was instrumental in the discovery of helium, neon, krypton, xenon and radon by Ramsay in the late 1890s and early 1900s, discoveries which helped to revise and complete the periodic table; see Freund (1968). At the same time Mendeleev's (1834-1907) antagonistic reaction to the discovery of argon might have been the primary reason for not being awarded the Nobel prize in Chemistry in 1906; see Gordin (2004). The discovery of the argon resulted from the careful unraveling of an empirical 'discrepancy', detected initially by Rayleigh (professor of Cambridge University's Cavendish Laboratory), in the density of nitrogen gas produced by two different procedures, broadly described as 'atmospheric' and 'chemical'. The conclusion, based on joint work between Rayleigh and Ramsay (professor of chemistry at University College, London), that the atmospheric air contains an inert gas they called argon, hitherto unknown, was reached after a long 'discovery' process based on trial and error. The idea of this paper is to revisit the Rayleigh-Ramsay historical episode with the view to 'capture' and systematize their learning from error process. Their 'trial and error' process was based on a carefully designed sequence of experiments, combined with an informal (by today's standards) analysis of the resulting data, which guided them to further and further experiments until they reached the conclusion that atmospheric nitrogen contained argon absorbed from atmospheric air. The perspective used for the proposed formalization is that of the error-statistical account (see Mayo, 1996). The error-probing reasoning can be used to formalize the scientists' resourceful ways in dealing with Duhemian problems, as they arose in their actual research situations. In particular the emphasis is place on a variety of techniques for avoiding and uncovering error (ibid, pp. 4-7) captures essential

strategies that Rayleigh and Ramsay used to both convince themselves, and the numerous skeptics, that their results were not experimental artifacts but that they had, indeed, discovered a new chemical element. The reasoning is that of arguing from error (ibid, p. 7), and it is precisely this reasoning, Mayo argues, that is formally captured in some of the standard statistical techniques, e.g., significance tests, analysis of variance, etc. Those who question whether statistical reasoning could possibly lie at the heart of scientific learning in general on the grounds that these techniques are developments of the 20th century overlook the extent to which they are actually systematizing and sharpening tools that scientists (and humans, in general) have used for many centuries to learn by induction in the face of errors and uncertainty. Indeed, the discovery of argon provides an excellent example of arguing from error which began as 'an empirical anomaly' whose thorough investigation eventually led to a substantial growth in experimental knowledge.

2

In section 2 the paper summarizes the story of the discovery of argon, paying particular attention to the systematic efforts first by Rayleigh and then jointly with Ramsay to establish the robustness of the empirical discrepancy between atmospheric and chemical nitrogen, and then explain its source, by eliminating other possible explanations.

Section 3 of the paper attempts a formalization of the statistical analysis performed by Rayleigh on his experimental data in order to make a case that the kind of inductive reasoning employed can be captured using the error-statistical approach. An important dimension of the error-statistical approach is concerned with the validity of the premises of inductive inference. It is shown that some of the (informal) empirical tests used by Rayleigh can be viewed as informal misspecification tests in order to establish the validity of the premises for his inferences. Section 4 repeats the statistical analysis of section 3 to more precise data generated by the collaborative efforts of Rayleigh and Ramsay. It is shown that the new data give rise to more incisive inferences.

2 The story of the discovery of Argon

The story of the discovery of argon is interesting from the philosophy of science perspective for several reasons. First, it constitutes a case where the new knowledge was not the result of prediction from a well-established theory, but a systematic search to explain an empirical discrepancy detected in an experiment. Indeed, one of the primary reasons for the initial reluctance to accept it was the fact that it could not be accommodated into the periodic table as understood at the time. Second, the fact that the new knowledge could not be accommodated in the theoretical framework at the time led to a major overhaul of that framework, including a complete revamping of the periodic table changing its basis from atomic weights to atomic numbers. Third, the discovery of argon provides an example where the careful analysis of the data from experiments led the scientists in question to a systematic sequence of other experiments that eventually led them to the unequivocal conclusion of the existence of argon, by trial and error; a clear example of learning from error. Fourth, the statistical analysis performed by these scientists in the 1890s is technically primitive by today's standards, but highly sophisticated from the inductive reasoning perspective. As demonstrated below, Rayleigh was able to draw reliable inductive inferences because his data analysis involved establishing a primitive form of statistical adequacy for his premises of inference; see Spanos (2006) for further discussion.

2.1 The initial detection of an empirical discrepancy

Lord Rayleigh's investigations that led to the discovery of argon began in the early 1890s with an attempt to measure the (molar) mass of nitrogen, after his successful determination of the mass of oxygen and hydrogen. In the 1890s the conventional wisdom was that atmospheric air was "a mixture of oxygen, nitrogen, and small

3

quantities of carbon dioxide and water vapour, together with a trace of ammonia." (Ramsay, 1915, p. 148). Rayleigh produced nitrogen gas from atmospheric air by removing

the other known constituents; calling it "atmospheric nitrogen". The original procedure he used was to bubble air through liquid ammonia and then pass it through

a hot tube. The water produced could be removed by drying agents and the nitrogen product joined nitrogen from the atmospheric sample. He was ready to publish his results, but decided to pursue some additional experiments in order to eliminate any potential errors due to the particular process used to produce the nitrogen:

"But then I reflected that it is always advisable to employ more than one method, than the method I had used — Mr. Vernon Harcourt's method — was not that which had been used by any of those who had preceded me in weighing nitrogen. The usual method consists in absorbing the oxygen of air by means of red-hot copper; and I thought that I ought at least to give that method a trial, fully expecting to obtain forthwith a value in harmony with that already afforded by the ammonia method." (Rayleigh, 1895, p. 702)

To his surprise, the nitrogen produced by passing air over hot copper, thus removing the oxygen as copper oxide, was lighter; he called this "chemical nitrogen". That is, when he compared the mass of nitrogen from these two different methods he found a small but 'persistent' (and repeatable) discrepancy that could not be explained away as experimental error:

"... the gas obtained by the copper method, ..., proved to be one-thousandth part heavier than that obtained by the ammonia method; and on repetition, that difference was only brought out more clearly." (1895, p. 702)

2.2 Magnifying the detected empirical discrepancy

Rayleigh's next step exemplifies the powerful intuition of an exceptional scientist:

"The next step in the enquiry was, if possible, to exaggerate the discrepancy. One's instinct at first is to try to get rid of a discrepancy, but I believe that experience shows such an endeavour to be a mistake. What one ought to do is to magnify a small discrepancy with a view to finding out the explanation; and, as it appeared in the present case that the root of the discrepancy lay in the fact that part of the nitrogen prepared by the ammonia method was nitrogen out of ammonia, although the greater part remained of common origin in both cases, the application of the principle suggested pure oxygen for atmospheric air in the ammonia method, so that the whole, instead of only part, of the nitrogen collected should be derived from the ammonia itself. The discrepancy was at once magnified some five times" (Rayleigh, 1895) [emphasis added]. That is, he devised a way to magnify the observed discrepancy by making specific changes to the experimental set-up in order to eliminate any errors due to using atmospheric air instead of pure oxygen.

4

2.3 Establishing the robustness of the empirical discrepancy

Rayleigh used a variety of different methods to produce both atmospheric and chemical nitrogen in order to eliminate any errors due to the peculiarities of the method of production. The sound experimental intuition exemplified by Rayleigh in utilizing such a variety of methods can be best understood as a way to ensure the robustness of his results to certain background assumptions (Mayo, 1996, p. 6). The idea was that if the different ways to produce, say atmospheric nitrogen, gave rise to the same measurement of density, one may argue that it is impossible that they all conspired to do so, this similarity is strong evidence of a reliable or robust measurement. In other words, if results were due to experimental artifacts, they are given an excellent chance to have been detected by at least one of the distinct ways of producing nitrogen. In this case, he wanted to ensure that the results of the two broad methods of producing nitrogen were not 'sensitive' to the particular process, instruments or chemicals used. For the atmospherically produced nitrogen Rayleigh used four (4) different methods to eliminate the oxygen using ammonia: hot iron, ferrous hydrate, hot iron, electrified and copper. For the chemically derived nitrogen he also used four different methods: nitric oxide (NO) by hot iron, nitrous oxide (N₂O) by hot iron, from N₂O by hot iron electrified, and ammonium nitrite (NH₄NO₂). Each of the four methods is used to obtain an average, and it is expected there will be some variability even if they are all measuring the same value. The question is whether they are close enough (within the experimental error bounds) to be regarded as having arisen from the same process. Rayleigh compared the averages of each particular method within the atmospheric nitrogen category: by hot iron (1893):

$$x_1 = 2.31017 + 2.30986 + 2.31010 + 2.31001$$

$$4 = 2.31003,$$

by copper (1892):

$$x_2 = 2.31026.$$

by hot iron, electrified (1894):

$$x_3 = 2.31163 + 2.30956$$

$$2 = 2.31059,$$

by ferrous hydrate (1894):

$$x_4 = 2.31024 + 2.31010 + 2.31028$$

$$3 = 2.3102,$$

with the overall average $x = 2.310221$, and concluded that the deviations:

$$(x - x_1) = .000191, (x - x_2) = .000021, (x - x_3) = -.000369, (x - x_4) = -.000039 \quad (1)$$

were 'small enough' to infer that all four production processes can be realistically viewed as giving rise to the same (homogeneous) product. That is, the differences in (1) are not 'large enough' to suggest that the nitrogen produced by each method was qualitatively different.

The same conclusion was reached by Rayleigh when comparing the means from the four different way of producing chemical nitrogen (NO — nitric oxide, N₂O — nitrous oxide, NH₄ NO₂ — ammonium nitrite):

from NO by hot iron (1893):

$$y_1 = 2.30143 + 2.29890 + 2.29816 + 2.30182$$

$$4 = 2.30008,$$

from N₂O by hot iron (1893):

$$y_2 = 2.29869 + 2.29940$$

$$2 = 2.29904,$$

from NH₄ NO₂ (1894):

$$y_3 = 2.29849 + 2.29889$$

$$2 = 2.29869$$

from N₂O by hot iron, electrified (1894):

$$y_4 = 2.30074 + 2.30054$$

$$2 = 2.30064$$

The overall average was $y = 2.299706$, and the small differences:

$$(y - y_1) = -.000374, (y - y_2) = .000666, (y - y_3) = .0010169, (y - y_4) = -.000934, \quad (2)$$

indicated that the nitrogen produced by each method was not qualitatively different.

The statistical analysis performed on this data by Lord Rayleigh was rudimentary (by today's standards), based primarily on 'averaging' and comparing the differences of these averages. Averaging was a well-known method to enhance the reliability of estimating the magnitude of a 'true' quantity when the observed data differ from it by non-systematic random effects, going back to Gauss in 1809. It was known at the time that the sample mean ($\bar{Y} = \frac{1}{n} \sum_{k=1}^n Y_k$) provides a more precise estimate of the 'true' mean (μ) than any one individual observation (Y_k) because the variance of the sample mean is n times smaller since $\text{Var}(\bar{Y}) = \frac{1}{n} \text{Var}(Y_k)$; the larger the number of observations n the higher the precision. This, and much else in his effort to probe the different ways his inferences could be in error, will be discussed more extensively in section 3.

nPn

$k=1$ Y_k) provides a more precise estimate of

the 'true' mean (μ) than any one individual observation (Y_k) because the variance of the sample mean is n times smaller since $\text{Var}(\bar{Y}) = \frac{1}{n} \text{Var}(Y_k)$

n ; the larger the number of

observations n the higher the precision. This, and much else in his effort to probe the different ways his inferences could be in error, will be discussed more extensively in section 3.

Having established the homogeneity of the two different broad ways of producing nitrogen using the atmospheric process as well as the chemical process, Rayleigh proceeded to compare the overall average weight of the nitrogen produced by the two processes x and y , respectively. The difference between the two overall means:

$$(x - y) = 0.010515, \quad (3)$$

was more than 67 times the average difference within the atmospheric nitrogen in (1)

and more than 15 times that within the chemical nitrogen in (2). His reasoning has a strong family resemblance to the Analysis of Variance (ANOVA) comparison based on within and between groups variation. On this basis Rayleigh concluded that:

"The difference is about 11 milligrams, or about one-half per cent.;

and it was sufficient to prove conclusively that the two kinds of nitrogen — the chemically derived nitrogen and the atmospheric nitrogen — differed in weight, therefore, of course, in quality, for some reason hitherto unknown." (Rayleigh, 1895, p. 702) [emphasis added].

The 'atmospheric nitrogen' had a density of 1.2572 grams per liter, whereas the 'chemical nitrogen' had a density of 1.2511 grams per liter. How reliable was this result?

6

"I need not spend time in explaining the various precautions that were necessary in order to establish surely that conclusion. One had to be on one's guard against impurities, especially against the presence of hydrogen, which might seriously lighten any gas in which it was contained." (Rayleigh, 1895, p. 702) [emphasis added].

2.4 'Reasoning from error' to explain the discrepancy

Rayleigh (1895) describes numerous precautions and experimental controls in establishing

the robustness of these results, including storing the nitrogen for eight months and re-weighting it in order to ensure that the discrepancy persisted. He concluded that the difference $(x - y) = 0.010515$ is not an artifact because it was well outside the magnitude of his experimental error. Again, Rayleigh's analysis uses reasoning which is analogous to a modern t-test assessing the difference between two means; see section 3. He published his results in a 1894 paper entitled "On an Anomaly Encountered in Determination of the Density of Nitrogen Gas". He was so concerned with this empirical discrepancy that he asked for help from other scientists, especially chemists:

"In order, if possible, to get further light upon a discrepancy which puzzled me very much, and which, at the time, I regarded only with disgust and impatience, I published a letter in Nature inviting criticisms from chemists who might be interested in such questions. I obtained various useful suggestions, but none going to the root of the matter." (Rayleigh, 1895, p. 702)

On the basis of his own evidence, Rayleigh reached a preliminary conclusion:

"Upon the assumption that similar gas should be obtained by both methods, we may explain the discrepancy by supposing either that:

h1 : the atmospheric nitrogen was too heavy on account of imperfect removal of oxygen, or that

h2 : the ammonia was too light on account of contamination with gases lighter than pure nitrogen," (Rayleigh, 1894, p. 340) [h1 and h2 inserted]

Rayleigh went on to investigate both of these possible explanations thoroughly by designing new experiments. Using the results of these experiments, in conjunction with established substantive knowledge in both physics and chemistry, he was able to eliminate both explanations.

He eliminated h1 both on experimental as well as theoretical grounds. New experiments introduced oxygen which was then burned, ensuring that no oxygen was left.

Using the similarity of the density of oxygen and nitrogen, he argued that, the observed discrepancy would have required that the nitrogen should have contained 1/30 of

its volume of oxygen, or 1/6

of that present in air, which was impossible on theoretical grounds! (ibid., 1894, p. 341)

Using similarly convincing arguments he eliminated the possible explanation h2:

"Of the possible impurities lighter than nitrogen, those most demanding

7

consideration are hydrogen, ammonia, and water vapour. The last may be dismissed at once, and the absence of ammonia is almost equally certain.

The question of hydrogen appears the most important. But this gas, and hydrocarbons, such as CH₄, could they be present, should be burnt by the copper oxide; and the experiments already referred to, in which hydrogen was purposefully introduced into atmospheric nitrogen, seem to prove conclusively that the burning would really take place." (ibid., p. 342)

Rayleigh went on to consider and eliminate other possible explanations that might

have given rise to h2, such as:

h3 : atmospheric nitrogen might contain three atom molecules, a heavier allotropic form of nitrogen, say, N₃.

It was known at the time that ozone (O₃) can be created by electrifying oxygen using 'the action of an electric discharge':

"Further experiments were tried upon the action of the silent electric discharge — both upon the atmospheric nitrogen and upon the chemically derived nitrogen — but neither of them seemed to be sensibly affected by such treatment" (Rayleigh, 1895, p. 703)

Reflecting on the hypotheses h1 and h2 we can see that they are not exhaustive. To render them exhaustive one needs to redefined them as follows:

h01 : the atmospheric nitrogen was too heavy on account of a heavier gas, and

h02 : the chemical nitrogen was too light on account of a lighter gas.

The obvious culprits for a heavier and lighter gas were oxygen and hydrogen, respectively. Both were eliminated, however, by designing experiments where the atmospheric and chemical nitrogen were purposely mixed with additional oxygen and hydrogen, respectively, and removing these contaminations using the various methods originally used to produce the nitrogen. Rayleigh proved that these methods were very effective in removing these two contaminants. His arguments constitute excellent examples of dealing with Duhem's problem by eliminating potential explanations.

At the end of a long trial and error process, Rayleigh's primary conclusion was:

"... altogether, the balance of evidence seemed to incline against the hypothesis of abnormal lightness in the chemically derived nitrogen being due to dissociation, and to suggest strongly, as almost the only possible alternative, that there must be in atmospheric nitrogen some constituent heavier than true nitrogen." (Rayleigh, 1895, p. 703) [emphasis added]

That is, Rayleigh revised his original hypothesis h1 to h01. His intuition that the atmospheric nitrogen may contain a heavier gas led him to seek advice from a famous chemist, Sir William Ramsay, and that initiated an intensive collaborative research effort that probed exhaustively the hypotheses h01 and h02. They first repeated

Rayleigh's experiments with additional controls in order to eliminate any form of impurity as the source of the discrepancy. (Rayleigh and Ramsay, 1895, pp. 191-2).

8

2.5 The discovery of Argon

On account of their training, Rayleigh focused primarily on the physical aspects and Ramsay on the chemical dimensions of this empirical discrepancy puzzle. As an experienced chemist Ramsay recalled the older experiments by Cavendish in 1785 who demonstrated that after both the oxygen and nitrogen were extracted from atmospheric

air there was still a small residue: "... That residue amounted to about 1/120 part of the nitrogen taken" (Rayleigh, 1895, p. 704).

After employing a thorough 'reasoning from error', Rayleigh and Ramsay eliminated h02, and began a joint intense effort to probe h01 by isolating and identifying the residual gas noted earlier by Cavendish:

"The simplest explanation in many respects was to admit the existence of a second ingredient in air from which oxygen, moisture, and carbonic anhydride had already been removed. The proportional amount required was not great. If the density of the supposed gas were double that of nitrogen, one-half per cent. only by volume would be needed; or, if the density were but half as much again as that of nitrogen, then one per cent. would still suffice. But in accepting this explanation, even provisionally, we had to face the improbability that a gas surrounding us on all sides, and present in enormous quantities, could have remained so long unsuspected." (Rayleigh and Ramsay, 1895, p. 192)

Ramsay devised an experiment with much smaller experimental error than that of Cavendish to remove all the nitrogen from his sample of 'atmospheric nitrogen' by passing it repeatedly over heated magnesium, with which nitrogen reacts to form magnesium nitride - a solid. As a result of this experiment he was left with 1% residual of the original volume which would not react. The same experiment applied

to 'chemical nitrogen' left no residual! In their own words:

"... the conclusion seems inevitable that "atmospheric nitrogen" is a mixture and not a simple body." (ibid., p. 208)

When they measured the density of the residual in the 'atmospheric nitrogen' they discovered that it was denser than nitrogen. In order to eliminate even the remote possibility that the unidentified gas was not the result of the process used to isolate it, Ramsay and Rayleigh devised yet another experiment where atmospheric nitrogen was passed through a long sequence of porous clay pipes surrounded by a vacuum. The portion of nitrogen diffused through the clay was less dense than the remainder, proving that the heavier gas must have been in the original air. The last alternative explanation they wanted to eliminate again was the possibility that the residual gas was a heavier allotropic form of nitrogen N₃. This possibility was championed by two famous chemists, James Dewar (professor of chemistry in the Royal Institution) and Dmitri Mendeleev (a professor of chemistry at St. Petersburg University), in their attempt to save the periodic table as it was shaped at the time; see Gordin (2004), p. 211.

9

The most crucial test to eliminate allotropic nitrogen as the heavier gas was the spectral analysis applied by Ramsay to the residual which established unequivocally that it was a hitherto unidentified gas. It should be noted that the spectral analysis was, at the time, the best test to identify the nature of any chemical element. Hence, using reasoning from error they were able to demonstrate experimentally that the discrepancy

is exclusively due to a hitherto unknown inert gas found in the atmosphere, which they called 'argon' (the Greek word for 'inactive') because of its chemical inertness.

It is now known that Argon has atomic number 18 (the total number of protons in its atomic nucleus), it has atomic weight 39.948, it constitutes 1.3 percent of the atmospheric air by weight and 0.94 percent by volume; see Emsley (2001). Having established the presence of argon in atmospheric nitrogen they went further and argued that:

"If the newly discovered gas were not in the atmosphere, the discrepancies in the density of "chemical" and "atmospheric" nitrogen would remain unexplained." (p. 235)

In Mayo's (1996) terminology, they reached this conclusion by ensuring that h01 passed a severe test in the sense that it "has withstood a scrutiny that it would very likely have failed, were it not correct." (ibid., p. 30).

2.6 Growth of experimental knowledge

Convincing the scientific establishment that they identified a new element that could not even fit into Mendeleev's periodic table, as it stood at the time, was not an easy task. To achieve that they embarked on a sequence of new improved experiments using additional controls in an attempt to eliminate as many impurities from the original data as possible. The new experiments of Rayleigh and Ramsay (1895) yielded more precise data which enabled them to enhance the discrepancy to:

$$(x - y) = 0.011167.$$

The monatomic nature of the new substance was established by measurements of constant-pressure and constant-volume heat capacities, but its existence remained controversial because it did not fit into Mendeleev's periodic table. Eventually they were able to provide conclusive physical evidence that the residual was indeed a new element, argon, and determine its density; 1.7824 grams per liter, compared to nitrogen's density of 1.2506 grams per liter. The additional probing enabled Rayleigh and Ramsay to ensure that the claim:

h4 : the atmospheric nitrogen was too heavy because it contained a heavier gas from atmospheric air, identified as argon, also passed a severe test; see Mayo (1996).

Rayleigh and Ramsay made a preliminary announcement of their discovery of argon at the Oxford meeting of the British Association in August of 1894. Predictably, their claim was met with skepticism bordering on hostility; it was not the first time, and it will not be the last, that someone has erroneously claimed the discovery of a new chemical element. This particular discovery was even more problematic since:

10

(i) it did not fit into the periodic table,
 (ii) it raised the possibility of a zero-valency group of elements, and
 (iii) its inertness made it impossible to determine its atomic weight by chemical means; the chemists of that period distrusted physical evidence concerning atomic weights. Dewar and Mendeleev were convinced that argon was in fact triadic nitrogen N₃; Dewar registered his dissension by not attending the formal presentation of the results at the Royal Society in January of 1895. Rayleigh must have been greatly frustrated by these incessant and unjustified (in his mind) criticisms to such an extent that he would soon return to his research in physics excusing his move by stating that in that field: "second rate men seem to know their place" (see Brock, 1992, p. 336). The hostility against their claim dissipated only when the other noble gases were discovered and Mendeleev's periodic table was reshaped by the early 1900's. Even Mendeleev, who argued vehemently against the claim, changed his mind:

"In 1903, he considered Ramsay's findings "some of the most brilliant experimental discoveries of the end of the 19th century," and admitted that his early hypothesis of triatomic nitrogen was incorrect. What changed his mind? Mendeleev cited five pieces of evidence that swayed him: the finding that argon's density was just barely greater than 19, while N₃ would have been around 21; Ramsay's discovery of helium in 1895, which also displayed chemical inertness; the later discoveries of the other inert gases neon and krypton; the uniqueness of their spectra; and Ramsay's proof of the constancy of chemical features when correlated with density." (Gordin, 2004, p. 211)

Ramsay (1899) went on to use the established densities, as well as the atomic weights of nitrogen and argon, to determine the 'theoretical' value of the discrepancy to be $\gamma^* = 0.01186$; note that estimated discrepancy $\gamma = 0.011167$ was reasonably accurate.

Ramsay (1915) provides a retrospective detailed account of the quest that led to the discovery of argon.

Rayleigh and Ramsay arrived at the further conclusion that if the periodic law and the discovery of a new inert elemental gas were both correct, then there must be a whole family of such elements, rendering Mendeleev's periodic table incomplete; see Freund (1968). Ramsay's subsequent work confirmed that conclusion by isolating helium first and then discovering neon, krypton, and xenon by the end of the century; adding a completely new column to the periodic table. Hence, the discovery of argon was more than just isolating yet another element. It was instrumental in giving rise to major revisions of the experimental knowledge of the 1870s.

Argon's discovery began the process of questioning the basic pillars of how matter was understood at the time:

- (i) integrability (each atom was integral - it had no substructure),
- (ii) immutability (each specific element had fixed mass and could not be transformed into another), and
- (iii) valency (a numerical charge that determined how a given atom

11

would combine with others).

The discovery of the other noble gases, followed immediately by the discovery of radioactivity, by Pierre and Marie Curie in 1898, and the electron (sub-elements of the atom) by J. J. Thompson in 1897, undermined completely the three basic pillars of knowledge on matter as it stood at the end of the 19th century; see Gordin (2004), p. 209. Argon called into question the valency of matter by initiating the zero-valency group. It also created a major problem with the periodic table because its atomic weight of 39.948 was bigger than that of potassium 39.098, but their atomic weights turned out to be 18 and 19, respectively. The periodic table was recast by Moseley in 1914 in terms of atomic numbers, replacing Mendeleev's periodic table in terms of atomic weights. It turned out that the atomic number, not only identifies the chemical properties of an element, but also facilitates the description of other aspects of atoms and nuclei. These early developments, which began with the discovery of argon, had an enormous impact on both chemistry and nuclear physics over the next several decades of the 20th century; see Brock (1992).

3 Data analysis and the Error-Statistical approach

3.1 A brief summary of the error-statistical approach

The term Error-Statistical approach was coined by Mayo (1996) to denote a modification/

extension of the framework for frequentist inductive inference, usually associated with Fisher, Neyman and Pearson. The modification/extensions come primarily in the form of:

(i) Emphasizing the learning from data (about the phenomenon of interest) objective of empirical modeling.

(ii) Paying due attention to the validity of the premises of induction via statistical adequacy, using thorough misspecification testing and respecification.

(iii) Emphasizing the central role of error probabilities in assessing the reliability of inference, both pre-data as well as post-data.

(iv) Supplementing the original framework with a post-data assessment of inference in the form of severity evaluations.

(v) Bridging the gap between theory and data using a sequence of interconnected models, theory (primary), structural (experimental), statistical (data) built on two different, but related, sources of information: substantive subject matter and statistical information (chance regularity patterns); see Spanos (1999).

(vi) Advancing thorough probing of the different ways an inductive inference might be in error, by localizing the error probe in the context of the different models in (v); see Mayo (1996), Spanos (2006).

The primary objective of this section is to revisit Rayleigh's data analysis and inference, using the error-statistical perspective. The discussion in section 2 above, describing the sequence of experiments by both Rayleigh and Ramsay and their 'learning from error' process, constitutes an excellent example of probing for error as de-

scribed in (vi) at the level of an experimental model. Some of the relevant errors in their probing were quantitative and statistical, but most of them were not. This is important to bring out because probing for error using severe testing reasoning does not have to be quantitative; see Mayo (1996).

Having said that, this section the paper focuses on formalizing the 'learning from error' process that is based on quantitative assessments of the experimental data using (i)-(v). The formalization is interesting because it demonstrates the capacity of the error-statistical approach to capture and systematize the 'learning from error' experience of scientists at the 'trenches'. It is important to emphasize that this does not amount to a reconstruction of the form advocated by Bayesians; see Howson and Urbach (1993).

3.2 Embedding the material experiment into a statistical model

The statistical model that one can use to embed Rayleigh's material experiment in its context is a simple modification of the simple Normal model where $\{Z_k, k \in \mathbb{N}\}$ is assumed to be a Normal, Independent and Identically Distributed (NIID) process with mean μ and variance σ^2 :

$$M_0 : Z_k \sim \text{NIID}(\mu, \sigma^2), k \in \mathbb{N}.$$

The modification consists in allowing for the possibility that the data constitute two heterogeneous sub-sets; have different means. That is,

$$z := (z_1, z_2, \dots, z_n) = (x_1, x_2, \dots, x_{n_1}, y_1, y_2, \dots, y_{n_2}),$$

where $n = n_1 + n_2$, constitute a realization of the same NI process but with two different means (see Cox and Hinkley, 1974), i.e. the two means simple Normal model:

$$M_1 : X_k \sim \text{NIID}(\mu_1, \sigma^2), Y_k \sim \text{NIID}(\mu_2, \sigma^2), k \in \mathbb{N}. \quad (4)$$

Viewed in the context of M_1 , Rayleigh's substantive (primary) hypothesis can be recast in the form of the difference between the two means:

$$H_0 : \mu := (\mu_1 - \mu_2) = 0, \text{ vs. } H_1 : \mu := (\mu_1 - \mu_2) > 0. \quad (5)$$

The alternative hypothesis is one-sided because there was substantive information ensuring that $\mu_1 > \mu_2$; rendering the subset $\mu_1 < \mu_2$ of the parameter space irrelevant.

Using Neyman-Pearson (N-P) hypothesis testing we can test (5) using (Cox and Hinkley, 1974):

$$\tau(Z) =$$

$$(X - Y)$$

$$\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}$$

$$n_1$$

$$+$$

$$\frac{s_2^2}{n_2}$$

$$n_2$$

$$H_0: v \sim \text{St}(v), C_1(\alpha) := \{z : \tau(z) > c_\alpha\}, (6)$$

$$X = 1$$

$$n_1 P_{n_1}$$

$$j=1 X_j, Y = 1$$

$$n_2 P_{n_2}$$

$$j=1 Y_j, s_{21}$$

$$= \frac{s_1^2}{n_1}$$

$$j=1 (X_j - \bar{X})^2$$

$$(n_1 - 1), s_{22}$$

$$= \frac{s_2^2}{n_2}$$

$$j=1 (Y_j - \bar{Y})^2$$

$$(n_2 - 1),$$

and ' $H_0: v \sim \text{St}(v)$ ' reads 'distributed under the null as Student's t with $v = (n_1 + n_2 - 2)$ degrees of freedom'. For $\alpha = .05$, $c_\alpha = 1.734$ ($\alpha = .025$, $c_\alpha = 2.101$).

13

3.3 Establishing a statistical discrepancy

Table 1 - Rayleigh data (1894)

Obs. Atmospheric (x) in g Chemical (y) in g

1. 2.31017 2.30143

2. 2.30986 2.29890

3. 2.31010 2.29816

4. 2.31001 2.30182

5. 2.31024 2.29869

6. 2.31010 2.29940

7. 2.31028 2.29849

8. 2.31163 2.29889

9. 2.30956 2.30074

10. 2.31026 2.30054

Rayleigh's data z_0 in table 1 yield the following estimates:

$$\bar{x} = 2.310221, \bar{y} = 2.299706, s_{21}$$

$$= .000000292699, s_{22}$$

$$= .000001724, n_1 = n_2 = 10,$$

$$(7)$$

which give rise to an observed test statistic:

$$\tau(z_0) = (\bar{x} - \bar{y}) \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

$$n_1$$

$$+$$

$$\frac{s_2^2}{n_2}$$

$$n_2$$

$$= \frac{(2.310221 - 2.299706) \sqrt{.000000292699}}{\sqrt{.000001724}}$$

$$10 = 23.415[.0000000], (8)$$

$$10$$

$$= 23.415[.0000000], (8)$$

where the number in square brackets denotes the p -value. The tiny p -value leads to a rejection of the null of no discrepancy, and in conjunction with a small sample size, indicates that z_0 provides strong evidence against the null; see Mayo and Spanos (2006).

Rayleigh's intuition that the detected discrepancy was not an artifact, is confirmed by this formal test, but the issue is whether one can appraise the size of the discrepancy $\mu = (\mu_1 - \mu_2) > 0$ warranted by his data z_0 . That is, go beyond the established statistical significance to the substantive significance. The traditional N-P hypothesis testing does not allow one to establish the presence of a substantive discrepancy because

it's too coarse in the sense that it concludes with the 'accept/reject H_0 decision.

One way to establish the presence or absence of a substantive discrepancy one can use the severity evaluation proposed by Mayo (1996).

3.4 Establishing a substantive discrepancy

In the context of Mayo's (1996) error-statistical account, a test T provides evidence for a hypothesis H only to the extent that the data x_0 , not only accord with hypothesis H , but, in addition, such a good result would have been highly unlikely if H were false. This requires the evaluation of the test's probativeness with respect to the ways a hypothesis H might be false. If a test T has very high probativeness to detect all

departures from H , when the test is applied, and no departure from H is detected, then we are justified in concluding that there is no evidence that any of the probed departures are present. Moreover, the high probativeness of the test allows us to go a step further and conclude that the absence of departures provides evidence that H is true — the higher the probativeness to detect departures from H the stronger the evidence for H . That is, we have evidence for H only to the extent that the hypothesis withstood a severe probe of the ways it could be false and survived; see Mayo and Spanos (2006).

In the case of Neyman-Pearson testing where H_0 is rejected, the severe testing reasoning can be used to establish the maximum size of the discrepancy from H_0 warranted by data z_0 . In this case one can argue that, by rejecting the null, one 'passes' the claim $\mu = (\mu_2 - \mu_1) > \gamma$, for some $\gamma \geq 0$. The idea underlying the severity assessment is to evaluate the probativeness of test T_α in detecting departures from the claim $\mu > \gamma$, i.e. departures in the direction $\mu \leq \gamma$, under the scenario that the $\mu > \gamma$ is false. That is, the relevant severity assessment comes in the form of evaluating 'the probability of observing sample realizations z less in accord with $\mu > \gamma$ than z_0 , under the scenario that $\mu > \gamma$ is false, or equivalently, $\mu \leq \gamma$ is true'. More formally (see Mayo and Spanos, 2006):

$\text{Sev}(\tau(z_0), \mu > \gamma) = P(\tau(Z) \leq \tau(z_0); \mu > \gamma \text{ is false}) = P(\tau(Z) \leq \tau(z_0); \mu \leq \gamma \text{ is true})$,
Evaluation of severity for different values of γ , in the case of test T_α , takes the form:

$\text{Sev}(23.415, \mu > \gamma) = P(\tau(Z) \leq 23.415 - \gamma)$

n_1

$+ s_2^2$

$n_2 - 1$,

giving rise to the numerical results shown below:

Relevant Inference $(\mu_2 - \mu_1) > \gamma$,

γ .008 .009 .01 .0105 .011 .012

$\text{Sev}(23.415, \mu > \gamma)$ 1.000 0.998 0.867 0.513 0.147 0.002

(9)

On the basis of the above severity evaluations, one can deduce, with high severity, that the maximum discrepancy licensed by data z_0 is in the range [.009, .01).

3.5 Statistical adequacy: securing the reliability of inference

It is often insufficiently appreciated that the above statistical analysis that gave rise to the inference of the presence of both a statistical and a substantive discrepancy, depends crucially on the probabilistic assumptions constituting the underlying statistical model (the premises of inference); see Mayo and Spanos (2004). When any of

15

the assumptions of the statistical model M_1 are invalid, any inferences based on it are likely to be unreliable because the nominal error probabilities tend to differ from the actual error probabilities (Mayo, 1996). An important facet of the error-statistical account is the statistical adequacy of the premises (see (ii) in section 3.1).

The model assumptions underlying M_1 are:

[i] Normality: $X_k \sim N(\cdot)$, $Y_k \sim N(\cdot)$, $k \in N = \{1, 2, \dots\}$,

[ii] Independence over k ,

[iii] Different means, $E(X_k) = \mu_1$, $E(Y_k) = \mu_2$, but constant over k ,

[iv] Same variance, $\text{Var}(X_k) = \text{Var}(Y_k) = \sigma^2$, constant over k .

How does one assess the appropriateness of these assumptions? In practice the assessment takes the form of an informal checking using data plots, as well as a formal testing using misspecification tests.

3.5.1 Misspecification Testing (assumptions [i]-[ii])

The k -plots of data $x := (x_1, x_2, \dots, x_{n_1})$ and $y := (y_1, y_2, \dots, y_{n_2})$, shown in fig. 1,

suggest no obvious departures from assumptions [i]-[iii] ($\mu_1 > \mu_2$). The Normality assumption [i] seems reasonable; the absence of any cycles in the k-plots indicates no departures from assumption [ii]; the arithmetic average seems constant over the observation period, suggesting no departures from assumption [iii]; see Spanos (1999), ch. 5.

In d e x

Data

1 2 3 4 5 6 7 8 9 1 0

2 .3 1 2

2 .3 1 0

2 .3 0 8

2 .3 0 6

2 .3 0 4

2 .3 0 2

2 .3 0 0

2 .2 9 8

V a r i a b l e

x

y

T i m e S e r i e s P l o t o f x , y

Fig. 1: k-plots of (x_k, y_k) , $k = 1, \dots, n$.

(10)

A closer comparison of the two plots in fig. 1, however, indicates that there might be a problem with assumption [iv] because the y_k data (dotted line) exhibit more variation.

The small sample size is an issue for formal misspecification testing, but it's not a major handicap.

- Testing Normality using the Shapiro-Wilks test yielded: $S-W(z) = .879[.121]$.

- Testing Independence using a runs test yielded: $R(z) = -0.000078[.782]$; neither test indicates any departures - see Spanos (1986), Mayo and Spanos (2004).

16

Assumptions [iii] and [iv] will be formally tested in next two sub-sections. Testing [iii] formally will confirm Rayleigh's sound scientific intuition when he deduced the homogeneity of the two types of nitrogen using a simple comparison based on (1)-(2).

3.5.2 Testing the constancy of the two means (assumption [iii])

As mentioned in section 3, before Rayleigh went on to compare the difference

$(x-y)=0.010515$

based on the two broad categories of nitrogen, atmospheric and chemical, he performed an informal homogeneity of the means assessment within each group, and concluded that each of the two categories separately comprised homogenous nitrogen. This was an important assessment for Rayleigh because if nitrogen within each of the two categories was not homogeneous the comparison of the overall means of the two groups would have been meaningless.

In modern statistical terminology Rayleigh performed a misspecification test (see Spanos, 1986, and Mayo and Spanos, 2004) to assess the homogeneity of the means within each group. That is, he tested:

[iii] $E(X_k) = \mu_1$, $E(X_k) = \mu_2$, constant over $k \in N$.

The homogeneity within each of the two groups is assumed by the statistical model, and thus ensuring its validity is part of securing the reliability of inference; testing without the boundaries of the statistical model - see Spanos (1999). One way one can formally test assumption [iii] is to use an ANOVA test (see Cox and Hinkley, 1974) as a misspecification test.

The ANOVA test for m sub-groups. with n_i observations in the i -th sub-group, takes the general form:

$F(Y) = \frac{1}{m-1} \sum_{i=1}^m n_i (y_i - \bar{y})^2$

$= \frac{1}{m-1} \sum_{i=1}^m n_i (y_i - \bar{y})^2$

$= \frac{1}{m-1} \sum_{i=1}^m n_i (y_i - \bar{y})^2$

$= \frac{1}{m-1} \sum_{i=1}^m n_i (y_i - \bar{y})^2$

$= 1$

$(n_i - 1)s_i^2$
 $n - m$
 $H_0 \vee F((m-1), (n-m)), C1 := \{y : F(y) > \alpha\}, (11)$
 where $n = P_m$
 $i=1$ n_i is the total number of observations;
 $y_i = 1$
 $n_i P_{ni}$
 $j=1$ y_{ij} is the sample mean of the i -th sub-group;
 y is the overall mean based on all observations;
 s_i^2
 $= 1$
 $(n_i - 1)P_{ni}$
 $j=1(y_{ij} - y_i)^2$ is the sample variance of the i -th sub-group;
 'H0 v F(k, l)' reads 'under H0 is F distributed with (k, l) degrees of freedom, and C1 denotes the rejection region for an α -significance level test. The F-test statistic is the ratio of the variation explained by the mean heterogeneity P_m
 $i=1 n_i(y_i - y)^2$ and
 the residual variation P_m
 $i=1(n_i - 1)s_i^2$
 .

The ANOVA information for the chemical nitrogen data in table 1 is given in table 2.

Table 2 - ANOVA for chemically produced nitrogen gas

Source of variation DF Sum of squares

Explained: P_m

$i=1 n_i(y_i - y)^2$ 3 .0000004198

Residual: P_m

$i=1(n_i - 1)s_i^2$

6 .0000022143

Total: P_m

$i=1 P_{ni}$

$j=1(y_{ij} - y)^2$ 9 .0000026341

17

where 'DF' stand for 'Degrees of Freedom'. The observed value of the test statistic:

$F(y_0) = (.0000004198/3)$

$(.0000022143/6) = 0.37917[.772]$,

in conjunction with a high p-value (in square brackets) indicate that there is no strong evidence against the homogeneity assumption. No evidence against H_0 , combined with the information that the nitrogen gas produced by these methods is qualitatively similar, can be interpreted as indicating that all the chemically based methods do produce homogeneous nitrogen gas.

Similarly, for the atmospherically produced nitrogen gas data (table 1), the ANOVA information is given in table 3.

Table 3 - ANOVA for atmospherically produced nitrogen gas

Source of variation DF Sum of squares

Explained: P_m

$i=1 n_i(x_i - \bar{x})^2$ 3 .00000042781

Residual: P_m

$i=1(n_i - 1)s_i^2$

6 .0000022143

Total: P_m

$i=1 P_{ni}$

$j=1(x_{ij} - \bar{x})^2$ 9 .0000026421

The value of the test statistic:

$F(x_0) = (.00000042781/3)$

$(.0000022143/6) = 0.3864[.767]$,

and the associated p-value indicate no evidence for departures from the homogeneity assumption.

Hence, the misspecification tests for the homogeneity of the means within each sub-group (assumption (iii)) revealed no departures for the data in table 1, confirming

Rayleigh's original inference based on an informal ANOVA test. It is interesting to note that the numerator of the F-test statistic (11) provides a more sophisticated way to evaluate the individual differences $(y - y_i)$, $i = 1, 2, \dots, m$, than the comparison in (2) used by Rayleigh.

3.5.3 Testing the equality of the two variances (assumption [iv])

Another model assumption which needs to be tested is [iv], the homogeneity of the variances between the two categories, which can be assessed by testing the hypotheses:

$$H_0 : \sigma^2_1$$

$$= \sigma^2_2$$

$$, \text{ vs. } H_1 : \sigma^2_2$$

$$> \sigma^2_1$$

$$, (12)$$

in the context of the 'encompassing' statistical model:

$$M_2 : X_k \sim \text{NIID}(\mu_1, \sigma^2_1)$$

$$), Y_k \sim \text{NIID}(\mu_2, \sigma^2_2)$$

$$), k \in N. (13)$$

Note that under H_0 , model M_2 reduces to the original model M_1 . A well-known test for (12) is:

$$v(Z) = \frac{s^2_2}{s^2_1}$$

$$\frac{n_2 - 1}{n_1 - 1}$$

$$) / (\frac{s^2_1}{s^2_2})$$

$$H_0 \sim F(n_2 - 1, n_1 - 1), C_1 := \{z : v(z) > c_\alpha\}, (14)$$

$$18$$

where $F(n_2 - 1, n_1 - 1)$ denotes the F-distribution with $(n_2 - 1, n_1 - 1)$ degrees of freedom; see Cox and Hinkley (1974). For $F(9, 8)$, $\alpha = .05$, $c_\alpha = 3.179$, and thus for the data in table 1:

$$v(z_0) = \frac{s^2_2}{s^2_1}$$

$$= \frac{.000001724}{.000000292699}$$

$$= 5.890[.007],$$

$$.000000292699 = 5.890[.007],$$

with p-value .007, in conjunction with a small sample size, provide strong evidence against the null.

In view of this rejection and the reasonableness of the other probabilistic assumptions [i]-[iii] underlying M_1 , it might seem judicious to respecify the original model

by replacing it with M_2 . However, it can be shown that using a test of the primary

hypothesis (5) which allows for the heterogeneity of the two variances, such as the Welch (1938) test, does not affect the inferences in sections 3.3-3.4.

4 'Sharpening' the Experimental Results

In a deliberate effort to improve the quality of the original data (table 1), Rayleigh

and Ramsay (1895) implemented additional controls and took more precautions to eliminate as many of the experimental errors as possible. Despite the fact that table

4 contains fewer observations, the 'cleaning up' of the data renders them more

'trustworthy' in measuring the discrepancy in 'weight' between the chemical and

atmospheric

nitrogen gas.

Table 4 - Rayleigh-Ramsay data (1895)

Atmospheric (x) Chemical (y)

1. 2.3103 2.3001

2. 2.3100 2.2990

3. 2.3102 2.2987

4. 2.2985

5. 2.2987

The Rayleigh-Ramsay data (table 4) yield the following estimates:

$$\bar{x} = 2.310166667, \bar{y} = 2.299, s^2_1$$

$$= .00000002334, s^2_2$$

$$= 0.00000041, n_1 = 3, n_2 = 5,$$

where the test statistic is:

$$T(z_0) = \frac{(\bar{x} - \bar{y})}{\sqrt{\frac{s^2_1}{n_1} + \frac{s^2_2}{n_2}}}$$

$$n_1$$

$$\begin{aligned}
 &+ \\
 &s_{22} \\
 &n_2 \\
 &= (2.310166667 - 2.299) \sqrt{.00000002334} \\
 &3 + 0.00000041 \\
 &5
 \end{aligned}$$

$$= 37.2678[.000000],$$

which leads to a strong rejection of the null of no discrepancy at $\alpha = .05$, $ca = 1.943$ ($\alpha = .025$, $ca = 2.447$). In view of the fact that the sample size is small, the p-value in square brackets suggests that data z_0 provide strong evidence against the null; see Mayo and Spanos (2006).

19

Evaluating the severity of the claim $\mu := (\mu_1 - \mu_2) > \gamma$, for different values of γ yields:

Relevant Inference $(\mu_2 - \mu_1) > \gamma$,

γ .008 .009 .01 .0105 .0108 .011 .012

Sev(37.2678, $\mu > \gamma$) 1.000 1.000 0.996 0.966 0.867 0.701 0.016

(15)

On the basis of the above severity evaluations based on test $\{\tau(Z), C_1\}$, with only 8 observations, one can deduce that the maximum size of the discrepancy warranted by the Rayleigh-Ramsay data z_0 is in the range $[\.009, .011]$.

As in the case of the data in table 1, the reliability of the inference, as well as the severity evaluations (15), depend crucially on the appropriateness of the assumed statistical model M_1 . It turns out that assumption (iv) is less problematic in this case and the use of the Welch (1939) test does not change the above inferences.

On the basis of the above severity evaluations one can deduce that the maximum discrepancy warranted by data z_0 is within the range $[\.01, .011]$. Taking into account the fact that no experimental data, even today, can be totally free of all possible impurities, as well as the small sample size, the severity evaluation of the maximum size of the discrepancy warranted by the data in table 4, established the severity evaluations

in (15), turned out to provide a very good approximation to the substantive value of $\gamma^* = .01186$ given by Ramsay (1899).

In concluding this section, it is interesting to note that the use of Confidence Intervals (CI), as a method to capture the substantive discrepancy γ^* , can be shown to be less effective in this particular case; the general reasons for this are discussed in Mayo and Spanos (2006).

5 Conclusion

The error-statistical perspective has been used to capture, as well as systematize, the 'learning from error' process that led Rayleigh and Ramsay to the discovery of argon. Their probing for error at all stages of the 'trial and error' process, and their careful but rudimentary statistical analysis of the data from their experiments, played an important role in guiding their learning process. It is shown that the error-statistical account, based on severe testing reasoning (Mayo, 1996), can be used to describe, codify and provide a naturalistic account of their quest. In particular, this approach was shown to accomplish three related objectives:

(a) it provides a coherent rationale for the probing for error process that systematically eliminates potential explanations in order to address Duhem's problem 'locally', at the level of a particular experiment,

20

(b) it captures the intuition underlying their informal probing for error procedures, including their 'diagnostic checks' to secure the validity of the (implicit) statistical model,

(c) it formalizes and systematizes their intuitive inductive reasoning that gave rise to a reliable measure of the magnitude of the discrepancy warranted by their data, using post-data severe testing evaluations.

Of particular interest is their use of primitive (by today's standards) statistical analysis in guiding them toward additional experiments which eventually led to the establishment of a substantive discrepancy between the density of atmospheric and chemical nitrogen. That in turn led them to the discovery of argon in the atmospheric

air, which precipitated a major revision of the periodic table. The importance of the discovery of argon was recognized early in the 20th century by awarding two Nobel prizes to Rayleigh and Ramsay in 1904, in physics and chemistry, respectively, and not awarding one in chemistry to Mendeleev in 1906.

University of Pittsburgh

Revised 3/10/08 - Copyright 2006