Evolution of a Strategy for the Enantioselective Synthesis of (–)-Cajanusine

Renyu Guo, Brittany P. Witherspoon, and M. Kevin Brown*

Cite This: *J. Am. Chem. Soc.* 2020, 142, 5002–5006

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: The first enantioselective synthesis of (–)-cajanusine is presented. Key features of the route include a rapid synthesis of the [4.2.0]bicyclooctane core by an enantioselective isomerization/stereoselective [2+2]-cycloaddition strategy as well as prominent use of catalytic methods for bond construction. The evolution of the approach is also presented that highlights unexpected roadblocks and how novel solutions were developed.

The stilbenoid oligomers constitute a broad family of natural products with high structural variance, many of which are derived from the common stilbenoid resveratrol.¹ Most stilbenoid oligomers arise from radical-mediated cyclizations that lead to the formation of products with ring sizes five and greater. However, cyclobutane stilbenoid dimers that likely arise from a formal [2+2]-cycloaddition of the alkene units are known.¹ In 2014, a new cyclobutane-containing stilbenoid dimer, cajanusine (1), was reported.² Here, the cyclobutane is likely generated from a formal [2+2]-cycloaddition with the oxygenated aryl group and the stilbenoid alkene unit, thus giving rise to a novel scaffold within this family of natural products (Scheme 1).²

Scheme 1. Cajanusine and Biosynthetic Precursors

Due to our laboratory's ongoing interest and work in developing enantioselective reactions to access cyclobutanes, we were drawn to targeting (–)-cajanusine (1) for synthesis.³ The pentasubstituted cyclobutane ring, as part of a highly functionalized [4.2.0]bicyclooctane core, was viewed as a challenge to access with traditional methods and strategies. Our group recently disclosed an enantioselective isomerization/stereoselective [2+2]-cycloaddition strategy for the synthesis of [4.2.0]bicyclooctanes.^{4–6} Given this premise, we focused on an approach that would implement this strategy toward (–)-cajanusine (1). As shown in Scheme 2, we expected that stereoselective [2+2]-cycloaddition of 2 would generate 3. Since the stereochemistry at C7 is opposite that of the natural product, an epimerization would be necessary. Release of steric pressure of the *syn*-1,2-diarylcyclobutane to

Scheme 2. Initial Strategy for the Synthesis of Cajanusine

the *anti*-1,2-diarylcyclobutane would serve as a driving force for the reaction. Elaboration of 4 to (–)-cajanusine (1) was envisioned to occur through stereoselective conjugate addition, followed by oxidation and prenylation of the C3 and C4 positions, respectively.

Our initial efforts commenced with synthesis of β,γ -unsaturated alkynyl ketone 8, an intermediate necessary for the enantioselective isomerization/stereoselective [2+2]-cycloaddition process (Scheme 3). Cross-coupling of dihydropyran 5 with PhMgBr under Ni catalysis as described by Wenkert provided the desired alcohol,⁷ which was oxidized to the aldehyde under Cu-catalyzed aerobic conditions reported by Stahl⁸ and then converted to epoxide 6. Addition of the

Received: January 11, 2020

Published: March 9, 2020

Scheme 3. First-Generation Approach

lithium acetylide of **7** and subsequent oxidation provided the β,γ -unsaturated alkynyl ketone **8**.

Enantioselective isomerization proceeded smoothly to generate the allenic ketone (not isolated), which, upon treatment with $\text{Bi}(\text{OTf})_3$, likely promoted the [2+2]-cycloaddition via pre-transition state **10**.⁴ Examination of the unpurified reaction mixture by ^1H NMR revealed that, in addition to the expected product **3**, cyclobutene **11** was formed as a significant product.⁴ Cyclobutene **11** likely arises from **3** under the reaction conditions via an acid-promoted alkene isomerization driven by relief of steric pressure of the *syn*-1,2-diarylcyclobutane motif. Furthermore, any attempts at purification by chromatography resulted in complete isomerization to cyclobutene **11**, which could be isolated in 35% yield and 98:2 er.

Given that epimerization at C7 is necessary for the synthesis of cajanusine, the isomerization reaction was deemed fortuitous. With access to cyclobutene **11**, isomerization of the alkene into conjugation with the ketone with concomitant formation of the desired stereoisomer **12** (aryl groups now anti) could be achieved upon treatment with DBU (Scheme 3). However, it appears that $K_{\text{eq}} \approx 1$, because the second isomerization event could not be driven toward completion under thermodynamic conditions. Preliminary ground-state calculations on the relative energies of **3**, **11**, and **12** gave them to be 1.98, 0.00, and -0.04 kcal/mol, respectively, which is consistent with experimental results. Despite the incomplete conversion to **12**, it was proposed that, after conjugate addition, cyclobutene **11** could be recycled if it was re-isolated from the reaction mixture. However, despite extensive experimentation, conjugate addition of a styrenyl-derived nucleophile to **12** could not be achieved. This is likely due to the steric demands in the generation of a quaternary carbon adjacent to the aryl group. This obstacle, coupled with the incomplete isomerization to **12**, which limited material throughput, forced us to consider other synthetic strategies toward (–)-cajanusine.

A second-generation approach was devised based on the premise that much of the difficulties experienced were due to an early-stage incorporation of the aryl group at C7 that forced a challenging conjugate addition. Therefore, strategies that installed the aryl group subsequent to the conjugate addition

were investigated (Scheme 4). Since the enantioselective isomerization/stereoselective [2+2]-cycloaddition process

Scheme 4. Second-Generation Approach

gives rise to the *syn*-diastereomer **14**, the aryl group at C7 needed to be incorporated from a functional handle that allowed for stereoselective C–C bond formation. With these criteria in mind, we considered a cross-coupling approach involving formation of a radical at C7 (**16**) that would allow for capture to occur on the convex face of the [4.2.0]-bicyclooctane core. Given the advances in Ni-catalyzed cross-coupling that involve the intermediacy of alkyl radicals, this strategy was deemed promising.⁹

With the consideration that an alkyl halide or redox-active ester would be necessary for the installation of the aryl group via Ni-catalyzed cross-coupling, a versatile protected alcohol was used as a handle. The requisite β,γ -alkynyl ketone **17** was

Scheme 5. Synthesis of (–)-Cajanusine

prepared in a sequence analogous to that shown in [Scheme 3](#) in five steps ([Scheme 5](#)).¹⁰ The enantioselective isomerization / stereoselective [2+2]-cycloaddition sequence proceeded to form **18** in good yield, diastereoselectivity, and enantioselectivity.⁴ Conjugate addition of *E*-styrenylboronic acid could be achieved through the use of Rh catalysis which, after desilylation, provided **19** in 73% yield.¹¹ Other approaches involving cuprates proved less fruitful and resulted in the formation of several unidentified impurities.

We elected to explore incorporation of the aryl group by cross-coupling of a redox-active ester (**20**, **27**), which could be prepared from **19** by oxidation to the acid and coupling with the necessary reagent. As has been noted, cross-coupling reactions of redox-active esters are substrate dependent, and it is therefore difficult to predict the success of any given set of conditions.^{12a} Therefore, an initial evaluation of 18 cross-coupling reactions¹² and 8 reductive cross-coupling reactions¹³ of various redox-active esters (**20**, **27**, and **28**) was conducted (Scheme 6).¹⁰ In general, the reductive cross-coupling reactions returned mostly starting material, whereas the standard cross-coupling reactions consumed the starting material. Of the 26 reactions evaluated, only a Ni-catalyzed cross-coupling¹⁴ of arylzinc **21** and an Fe-catalyzed cross-coupling¹⁵ of diarylzinc reagent **29**, both with redox-active ester **20** (TCNHPI = tetrachloro-*N*-hydroxyphthalimide), delivered the product in promising yields (20% and 11%, respectively). The Ni-catalyzed cross-coupling was selected for further optimization because of the higher initial yield (Scheme

6, inset table, entry 1). A screen of ligands revealed that BPhen was superior to dtbpy (Scheme 6, inset table, entry 5).¹⁶ While further improvement in yield was not observed with other temperatures or increased quantities of arylzinc reagent **21** (Scheme 6, inset table entries 6–8), higher catalyst loadings did result in increased yield (Scheme 6, inset table entry 9).

At this stage, completion of the synthesis required incorporation of the substituents at C4 and C3. This sequence was initiated through selective enolization of the sterically less demanding α -position of the ketone with *N*-MePMP (*N*-methyl pentamethylpiperidine) and Me_3SiOTf (Scheme 5).¹⁷ It is important to note that deprotonation with LDA, LHMDs, LiTMP, or $\text{LiN}(t\text{-Bu})\text{CPh}_3$ ¹⁸ resulted in the formation of enolate isomers. Conversion of the enol silane to the α,β -unsaturated carbonyl was achieved by treatment with IBX-MPO.¹⁹ Copper-catalyzed conjugate addition of Bpin and subsequent oxidation resulted in the formation of dione **23** (1:2:1 mixture of keto and enol tautomers).²⁰ Direct installation of the prenyl group proved challenging, as has been noted in the syntheses of other prenylated natural products.²¹ Therefore, an approach involving allylation followed by cross-metathesis with isobutene was explored. Surprisingly, it was difficult to incorporate a single allyl group, as bis-allylation was difficult to suppress. Thus, an efficient two-step sequence was developed in which Pd-catalyzed bis-allylation²² was followed by Ni-catalyzed mono-deallylation.²³ Finally, regioselective O-methylation (4:1 C3-OMe versus C5-OMe, > 20:1 after purification), cross-metathesis²⁴ with

Scheme 6. Optimization of the Cross-Coupling

isobutene, and deprotection allowed for synthesis of (–)-cajanusine (1). Due to the scalability and simplicity of many of the steps, >100 mg of cajanusine was easily prepared (98 mg was prepared in one reaction).

In summary, the first enantioselective synthesis of (–)-cajanusine is presented. The evolution of the strategy is highlighted to showcase the unexpected challenges that occurred and the solutions that were developed. In addition, the synthesis demonstrates the utility of various catalytic methods to function in complex settings and further underscores the utility of the enantioselective isomerization/stereoselective [2+2]-cycloaddition method developed in our laboratory.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.0c00359>.

Experimental procedures; analytical data for all new compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

M. Kevin Brown – Department of Chemistry, Indiana University, Bloomington, Indiana 47405, United States;
orcid.org/0000-0002-4993-0917; Email: brownmkb@indiana.edu

Authors

Renyu Guo – Department of Chemistry, Indiana University, Bloomington, Indiana 47405, United States
Brittany P. Witherspoon – Department of Chemistry, Indiana University, Bloomington, Indiana 47405, United States

Complete contact information is available at:
<https://pubs.acs.org/10.1021/jacs.0c00359>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Indiana University and the NIH (R01GM110131 and R35GM131755) for financial support. This project was partially funded by the Vice Provost for Research through the Research Equipment Fund and the NSF (CHE1726633). We thank Qi Cui (Peking University) for calculations on intermediates 3, 11, and 12. We thank Prof. Ying Wang (Jinan University) for providing a sample of cajanusine.

■ REFERENCES

- (1) Shen, T.; Wang, X.-N.; Lou, H.-X. Natural Stilbenes: an Overview. *Nat. Prod. Rep.* **2009**, 26, 916.
- (2) Li, X.-L.; Zhao, B.-X.; Huang, X.-J.; Zhang, D.-M.; Jiang, R.-W.; Li, Y.-J.; Jian, Y.-Q.; Wang, Y.; Li, Y.-L.; Ye, W.-C. (+)- And (–)-Cajanusine, a Pair of New Enantiomeric Stilbene Dimers with a New Skeleton From the Leaves of *Cajanus cajan*. *Org. Lett.* **2014**, 16, 224.
- (3) (a) Conner, M. L.; Xu, Y.; Brown, M. K. Catalytic Enantioselective Allenolate–Alkene [2+2] Cycloadditions. *J. Am. Chem. Soc.* **2015**, 137, 3482–3485. (b) Xu, Y.; Hong, Y. J.; Tantillo, D. J.; Brown, M. K. Intramolecular Chirality Transfer [2+2] Cycloadditions of Allenolates and Alkenes. *Org. Lett.* **2017**, 19, 3703–3706. (c) Wiest, J. M.; Conner, M. L.; Brown, M. K. Synthesis of (–)-Hebelophyllene-E: an Entry to Geminal Dimethyl-Cyclobutanes by [2+2] Cycloaddition of Alkenes and Allenolates. *Angew. Chem., Int. Ed.* **2018**, 57, 4647–4651. (d) Wiest, J. M.; Conner, M. L.; Brown, M. K. Allenolates in Enantioselective [2+2] Cycloadditions: From a Mechanistic Curiosity to a Stereospecific Transformation. *J. Am. Chem. Soc.* **2018**, 140, 15943–15949. (e) Conner, M. L.; Wiest, J. M.; Brown, M. K. Thioallenolates in Catalytic Enantioselective [2+2]-Cycloadditions with Unactivated Alkenes. *Tetrahedron* **2019**, 75, 3265–3271.
- (4) Line, N. J.; Witherspoon, B. P.; Hancock, E. N.; Brown, M. K. Synthesis of Ent-[3]-Ladderanol: Development and Application of Intramolecular Chirality Transfer [2+2] Cycloadditions of Allenic Ketones and Alkenes. *J. Am. Chem. Soc.* **2017**, 139, 14392.
- (5) For [2+2]-cycloaddition with allenic ketones, see: Zhao, J.-F.; Loh, T.-P. Acid-Catalyzed Intramolecular [2+2] Cycloaddition of Ene-Allenones: Facile Access to Bicyclo [N.2.0] Frameworks. *Angew. Chem., Int. Ed.* **2009**, 48, 7232.
- (6) For enantioselective isomerization to access allenolates, see: (a) Liu, H.; Leow, D.; Huang, K.-W.; Tan, C.-H. Enantioselective Synthesis of Chiral Allenolates by Guanidine-Catalyzed Isomerization of 3-Alkynoates. *J. Am. Chem. Soc.* **2009**, 131, 7212. (b) Inokuma, T.; Furukawa, M.; Uno, T.; Suzuki, Y.; Yoshida, K.; Yano, Y.; Matsuzaki, K.; Takemoto, Y. Bifunctional Hydrogen-Bond Donors That Bear a Quinazoline or Benzothiadiazine Skeleton for Asymmetric Organo-catalysis. *Chem. - Eur. J.* **2011**, 17, 10470.

- (7) Wenkert, E.; Michelotti, E. L.; Swindell, C. S.; Tingoli, M. Transformation of Carbon-Oxygen Into Carbon-Carbon Bonds Mediated by Low-Valent Nickel Species. *J. Org. Chem.* **1984**, *49*, 4894.
- (8) Hoover, J. M.; Stahl, S. S. Highly Practical Copper(I)/TEMPO Catalyst System for Chemoselective Aerobic Oxidation of Primary Alcohols. *J. Am. Chem. Soc.* **2011**, *133*, 16901.
- (9) Hu, X. Nickel-Catalyzed Cross Coupling of Non-Activated Alkyl Halides: a Mechanistic Perspective. *Chem. Sci.* **2011**, *2*, 1867.
- (10) See the [Supporting Information](#) for details.
- (11) Hayashi, T.; Yamasaki, K. Rhodium-Catalyzed Asymmetric 1,4-Addition and Its Related Asymmetric Reactions. *Chem. Rev.* **2003**, *103*, 2829.
- (12) (a) Sandfort, F.; O'Neill, M. J.; Cornella, J.; Wimmer, L.; Baran, P. S. Alkyl-(Hetero)Aryl Bond Formation via Decarboxylative Cross-Coupling: a Systematic Analysis. *Angew. Chem., Int. Ed.* **2017**, *56*, 3319. (b) Liu, X.-G.; Zhou, C.-J.; Lin, E.; Han, X.-L.; Zhang, S.-S.; Li, Q.; Wang, H. Decarboxylative Negishi Coupling of Redox-Active Aliphatic Esters by Cobalt Catalysis. *Angew. Chem., Int. Ed.* **2018**, *57*, 13096. (c) Chen, T.-G.; Zhang, H.; Mykhailiuk, P. K.; Merchant, R. R.; Smith, C. A.; Qin, T.; Baran, P. S. Quaternary Centers by Nickel-Catalyzed Cross-Coupling of Tertiary Carboxylic Acids and (Hetero)-Aryl Zinc Reagents. *Angew. Chem., Int. Ed.* **2019**, *58*, 2454.
- (13) (a) Huihui, K. M. M.; Caputo, J. A.; Melchor, Z.; Olivares, A. M.; Spiewak, A. M.; Johnson, K. A.; Dibenedetto, T. A.; Kim, S.; Ackerman, L. K. G.; Weix, D. J. Decarboxylative Cross-Electrophile Coupling of *N*-Hydroxyphthalimide Esters with Aryl Iodides. *J. Am. Chem. Soc.* **2016**, *138*, 5016. (b) Anka-Lufford, L. L.; Huihui, K. M. M.; Gower, N. J.; Ackerman, L. K. G.; Weix, D. J. Nickel-Catalyzed Cross-Electrophile Coupling with Organic Reductants in Non-Amide Solvents. *Chem. - Eur. J.* **2016**, *22*, 11564. (c) Suzuki, N.; Hofstra, J. L.; Poremba, K. E.; Reisman, S. E. Nickel-Catalyzed Enantioselective Cross-Coupling of *N*-Hydroxyphthalimide Esters with Vinyl Bromides. *Org. Lett.* **2017**, *19*, 2150. (d) Zhang, J.; Li, Z.; Zhuo, J.; Cui, Y.; Han, T.; Li, C. Tandem Decarboxylative Cyclization/Alkenylation Strategy for Total Syntheses of (+)-Longirabdiol, (–)-Longirabdolactone, and (–)-Effusin. *J. Am. Chem. Soc.* **2019**, *141*, 8372.
- (14) (a) Cornella, J.; Edwards, J. T.; Qin, T.; Kawamura, S.; Wang, J.; Pan, C.-M.; Gianatassio, R.; Schmidt, M.; Eastgate, M. D.; Baran, P. S. Practical Ni-Catalyzed Aryl-Alkyl Cross-Coupling of Secondary Redox-Active Esters. *J. Am. Chem. Soc.* **2016**, *138*, 2174. (b) Beck, J. C.; Lacker, C. R.; Chapman, L. M.; Reisman, S. E. A Modular Approach to Prepare Enantioenriched Cyclobutanes: Synthesis of (+)-Rumphellaone A. *Chem. Sci.* **2019**, *10*, 2315. (c) Shang, M.; Feu, K. S.; Vantourout, J. C.; Barton, L. M.; Osswald, H. L.; Kato, N.; Gagaring, K.; McNamara, C. W.; Chen, G.; Hu, L.; Ni, S.; Fernández-Canelas, P.; Chen, M.; Merchant, R. R.; Qin, T.; Schreiber, S. L.; Melillo, B.; Yu, J.-Q.; Baran, P. S. Modular, Stereocontrolled *C* β -H/*C* α -C Activation of Alkyl Carboxylic Acids. *Proc. Natl. Acad. Sci. U. S. A.* **2019**, *116*, 8721.
- (15) Toriyama, F.; Cornella, J.; Wimmer, L.; chen, T.-G.; Dixon, D. D.; Creech, G.; Baran, P. S. Redox-Active Esters in Fe-Catalyzed C–C Coupling. *J. Am. Chem. Soc.* **2016**, *138*, 11132.
- (16) See the Supporting Information of ref 14a.
- (17) Hu, P.; Chi, H. M.; DeBacker, K. C.; Gong, X.; Keim, J. H.; Hsu, I. T.; Snyder, S. A. Quaternary-Centre-Guided Synthesis of Complex Polycyclic Terpenes. *Nature* **2019**, *569*, 703.
- (18) Busch-Petersen, J.; Corey, E. J. Sterically Shielded Secondary *N*-Tritylamines and *N*-Tritylamide Bases, Readily Available and Useful Synthetic Reagents. *Tetrahedron Lett.* **2000**, *41*, 2515.
- (19) Nicolaou, K. C.; Gray, D. L. F.; Montagnon, T.; Harrison, S. T. Oxidation of Silyl Enol Ethers by Using IBX and IBX•N-Oxide Complexes: a Mild and Selective Reaction for the Synthesis of Enones. *Angew. Chem., Int. Ed.* **2002**, *41*, 996.
- (20) Mun, S.; Lee, J.-E.; Yun, J. Copper-Catalyzed β -Boration of α , β -Unsaturated Carbonyl Compounds: Rate Acceleration by Alcohol Additives. *Org. Lett.* **2006**, *8*, 4887.
- (21) For a review, see: Yang, X.-W.; Grossman, R. B.; Xu, G. Research Progress of Polycyclic Polyprenylated Acylphloroglucinols. *Chem. Rev.* **2018**, *118*, 3508.
- (22) Ruprah, P. K.; Cros, J.-P.; Pease, J. E.; Whittingham, W. G.; Williams, J. M. J. Studies Towards the Total Synthesis of Cycloaraneosene and Ophiobolin M: A General Strategy for the Construction of the 5–8 Bicyclic Ring System. *Eur. J. Org. Chem.* **2002**, *2002*, 3145.
- (23) Nečas, D.; Turský, M.; Kotora, M. Catalytic Deallylation of Allyl- and Diallylmalonates. *J. Am. Chem. Soc.* **2004**, *126*, 10222.
- (24) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. Efficient and Recyclable Monomeric and Dendritic Ru-Based Metathesis Catalysts. *J. Am. Chem. Soc.* **2000**, *122*, 8168.