

# COALGEBRAS ON MEASURABLE SPACES

Ignacio Darío Viglizzo

Submitted to the faculty of the Graduate School

in partial fulfillment of the requirements

for the degree

Doctor of Philosophy

in the Department of Mathematics,

Indiana University

August 2005

Accepted by the Graduate Faculty, Indiana University, in partial fulfillment of the requirements of the degree of Doctor of Philosophy.

Doctoral  
Committee

---

Lawrence S. Moss, Ph.D.  
(Principal Advisor)

---

Esfandiar Haghverdi, Ph.D.

---

David McCarty, Ph.D.

July 21, 2005

---

Alberto Torchinsky, Ph.D.

Copyright © 2005

Ignacio Darío Viglizzo

ALL RIGHTS RESERVED

To Ana

# Acknowledgements

There are many people without whose help I could not have completed this dissertation. These few words here reflect but a small part of the gratitude I own all of them.

My advisor Larry Moss supported me every step of the way. Since my arrival to Bloomington he shared generously his time, knowledge, insight and passion for new challenges. He encouraged me to pursue my ideas and helped me to bring them to fruition. I had the privilege of closely collaborating with him, learning first hand from his expertise.

The members of my Research Committee, Esfandiar Haghverdi, David McCarty and Alberto Torchinsky were also my professors. They always found time to extend their friendship and talk about the bigger picture of the mathematics we discussed. I greatly appreciate and will always remember these discussions.

The faculty of the Departments of Mathematics, Computer Science and Philosophy enriched and stimulated my intellectual life. I'm particularly grateful to Jon Barwise, Grahame Bennet, Michael Dunn, Douglas Hofstadter, Daniel Leivant and Slawomir Solecki.

The staff of the Department of Mathematics was exceptionally helpful all along. Misty Cummings was always a step ahead of my needs, so I never had to worry about any paperwork.

My fellow students and friends in the Departments of Mathematics, Computer Science and

Astronomy, who were always there with me to share the good times, bad times, homework and office spaces: Stefano Borgo, Sofía Brenes Barahona, Anastasiya Chagrova, Fulya Erdinc, Jim Hendrickson, Stella Kafka, Tei Laine, Francisco Lara-Dammer, Hayoung Lee, Maricarmen Martínez, Lei Qian, Thomas Reichherzer, Alek Slominski, Nik Swoboda and Alejandro Valerio. With Saleh Aliyari, Seunghwan Lee and Fernando Tohmé I also shared particularly fruitful discussions.

In Bloomington I found many friends: Sarah Ellis, Julie England, Kate Holden, Jeff and Dru Jarret, Waki Murayama, Gregory Rawlins, Steve Simms, Raja Thiagarajan and the rest of the SFDG. All of them made living here a daily joy.

My friends and family in Bahía Blanca, Argentina, never failed to encourage and support me. I hope they know how much that meant to me.

No words will ever be enough to thank my wife Ana Maguitman for all she does and is.

A teaching assistantship from the Department of Mathematics and a Fulbright fellowship provided financial support during the past six years at Indiana University. I thank these institutions for the trust they put in me.

# Abstract

Given an endofunctor  $T$  in a category  $\mathcal{C}$ , a coalgebra is a pair  $(X, c)$  consisting of an object  $X$  and a morphism  $c : X \rightarrow T(X)$ .  $X$  is called the *carrier* and the morphism  $c$  is called the *structure map* of the  $T$ -coalgebra.

The theory of coalgebras has been found to abstract common features of different areas like computer program semantics, modal logic, automata, non-wellfounded sets, etc. Most of the work on concrete examples, however, has been limited to the category **Set**. The work developed in this dissertation is concerned with the category **Meas** of measurable spaces and measurable functions.

Coalgebras of measurable spaces are of interest as a formalization of Markov Chains and can also be used to model probabilistic reasoning. We discuss some general facts related to the most interesting functor in **Meas**,  $\Delta$ , that assigns to each measurable space, the space of all probability measures on it. We show that this functor does not preserve weak pullbacks or  $\omega^{op}$ -limits, conditions assumed in many theorems about coalgebras. The main result will be two constructions of final coalgebras for many interesting functors in **Meas**. The first construction (joint work with L. Moss), is based on a modal language that lets us build formulas that describe the elements of the final coalgebra. The second method makes use of a subset of the projective limit of the final sequence for the functor in question. That is, the sequence  $1 \leftarrow T1 \leftarrow T^2 1 \leftarrow \dots$  obtained by iteratively applying

the functor to the terminal element 1 of the category. Since these methods seem to be new, we also show how to use them in the category **Set**, where they provide some insight on how the structure map of the final coalgebra works.

We show as an application how to construct universal Type Spaces, an object of interest in Game Theory and Economics. We also compare our method with previously existing constructions.



# Contents

Acknowledgements	v
Abstract	vii
<b>1 Introduction</b>	<b>1</b>
<b>2 Coalgebras and Limits</b>	<b>4</b>
2.1 Coalgebras . . . . .	4
2.2 Limits . . . . .	6
2.3 Bisimulations . . . . .	9
2.4 Final Sequences and Final Coalgebras . . . . .	11
<b>3 Measurable Spaces</b>	<b>19</b>
3.1 Measure Theory . . . . .	19
3.2 Weak Pullbacks and the Functor $\Delta$ . . . . .	23
3.3 $\omega^{op}$ -limits and the Functor $\Delta$ . . . . .	25

3.4	Kolmogorov's consistency Theorem . . . . .	30
3.5	A Lemma on generators of $\sigma$ -algebras . . . . .	33
<b>4</b>	<b>Modal Languages and Satisfied Theories</b>	<b>38</b>
4.1	Modal Languages for Measure Polynomial Functors . . . . .	38
4.2	The Space of Satisfied Theories . . . . .	43
4.3	From $S^*$ to $S(Id^*)$ . . . . .	46
4.4	Final Coalgebras . . . . .	52
<b>5</b>	<b>Final Coalgebras Found in Final Sequences</b>	<b>54</b>
5.1	The Space of Realised Points in the Final Sequence . . . . .	54
5.2	From $Z_S$ to $S(Z)$ . . . . .	57
5.3	Final Coalgebras . . . . .	62
<b>6</b>	<b>Probabilistic Kripke Polynomial Functors in Set</b>	<b>65</b>
<b>7</b>	<b>Type Spaces</b>	<b>73</b>
7.1	Motivation . . . . .	73
7.2	Measure polynomial functors in many variables . . . . .	82
7.3	Universal type spaces . . . . .	85
7.4	A brief review of the literature on type spaces . . . . .	86
	Armbruster, Böge and Eisele . . . . .	86

Mertens and Zamir . . . . .	89
Heifetz and Samet . . . . .	94
Other related work . . . . .	95
<b>8 Conclusions</b>	<b>97</b>
<b>Bibliography</b>	<b>100</b>

# Introduction

The main subject of this dissertation is the theory of coalgebras on measurable spaces. Coalgebras are simply defined as pairs  $(X, c)$  consisting of an object in a category  $\mathcal{C}$  and a morphism  $c : X \rightarrow FX$  where  $F$  is an endofunctor of the category  $\mathcal{C}$ . For some time now, the standard introduction to coalgebras have been the tutorials by Gumm [Gum99] and Rutten [Rut00]. In those works, numerous examples of applications of the abstract theory are shown, including non-wellfounded sets (see also Aczel [Acz88] and Turi and Rutten [TR98]), infinite data structures like streams and trees, different kinds of automata (including non-deterministic ones), labeled transition systems, graphs, Kripke models or frames and even some dynamical systems. All these examples are realised as coalgebras for functors in the category **Set** of sets and the usual set functions. The existence of notions like probabilistic automata and transition systems, continuous time systems and Markov chains, make it desirable to extend the theory of coalgebras to the category **Meas** of measurable spaces and measurable functions, where in particular, probability measures can be studied. Probabilities come into the picture through the analysis of the endofunctor  $\Delta$  in **Meas**. For a given measurable space  $X$ ,  $\Delta X$  is the space of all the probability measures that can be defined over  $X$ .

The first explorations of coalgebras on some measure spaces [RdV99] were led by the observation

---

that bisimulation, as defined for probabilistic systems by Larsen and Skou [LS91] was a particular case of the general coalgebraic concept due to Aczel and Mendler, [AM89] (see also [Mos99]). These studies were restricted to specific measure spaces in categories of metric and ultrametric spaces [RdV99]. Probabilistic transition systems have been studied by many computer scientists, see for example the papers by Desharnais et al., [BDEP97], [DEP02], [DGJP03], and also by van Breugel et al., [vBSW02], [vBMOW03], [vBMOW05]. For a survey on the different functors studied, see Bartels, Sokolova and de Vink, [BSdV03].

One application has been the guiding example for the research presented here. This is the definition of *type spaces* from game theory applied to economics. In particular, the problem of finding the *universal type space*, which turns out to be a final coalgebra (this is, a final object in the category of coalgebras for the functor that represent type spaces). Here the probabilities are used to model subjective beliefs of players in a game. That these beliefs can be modeled with probability measures is called the *Bayesian approach*. This branch of studies is independent from the work in computer science, but as we'll see, both of these are encompassed by the class of functors we deal with in this thesis.

Final coalgebras for a wide class of endofunctors in  $\mathbf{Meas}$  are obtained through two related methods. The first one was developed jointly with Lawrence S. Moss, and makes use of a modal language  $\mathcal{L}(T)$  defined in terms of the functor  $T$ . This language proves to be expressive enough to let us build a coalgebra whose elements are certain sets of its formulas. Modal languages like these had been studied by Jacobs [Jac01], Rößiger [Röß99, Röß00, Röß01] and Kurz [Kur01]. No logical system is proposed for the languages introduced. Instead, theories are built by collecting sets of formulas satisfied in some model.

In the second construction we use the *final sequence* for the functor. This is the sequence obtained by applying the functor to the final object in the category, and iterating the procedure  $\omega$

---

many times. We then take the projective limit and identify a subspace that will be the carrier of the final coalgebra. The final sequence has been used before to obtain final coalgebras, and in some cases its projective limit is the final coalgebra itself [SP82], but when the functors involved do not satisfy the condition of being  $\omega^{op}$ -continuous, more work needs to be done, as is the case here. For the category  $\mathbf{Set}$ , the final sequence has been studied by Worrell [Wor99] and Kurz and Pattinson [KP02]. This thesis is the first study of the final sequence in  $\mathbf{Meas}$ .

Chapter 2 introduces the general theory of coalgebras, and the construction of final coalgebras in  $\mathbf{Set}$  using the final sequence. An example in which this construction fails is included and is original to this dissertation.

In Chapter 3 we include the basic measure theory that will be needed to prove the main results. Lemma 3.6 is new, and allows us to build the languages in the Chapter 4 without using negation.

Chapter 4 presents the results that already appeared in [MV04] and [MV05].

Part of the results of Chapter 5 will appear in [Vig05].

Chapter 6 shows how to adapt the construction in Chapter 5 to the category  $\mathbf{Set}$ . These results are announced in [Vig05] but their proof appears here for the first time.

Chapter 7 expands on the treatment of type spaces sketched in [MV04], using the results from chapters 4 and 5 to construct the *universal type space*. We also offer a review of the bibliography on the topic, trying to clarify the relationship between different formalizations of type space that have been proposed.

Finally, in Chapter 8 we recapitulate the work and point out directions for further research and some open problems.

## 2

---

# Coalgebras and Limits

In this chapter we introduce coalgebras and some facts of category theory that will be helpful in organizing the material that follows. We also present a well known method for constructing final coalgebras, and an example where this method fails, motivating the new method from chapters 4, 5 and 6.

## 2.1 Coalgebras


**Definition 2.1.** Given a category  $\mathcal{C}$ , and an endofunctor  $T : \mathcal{C} \rightarrow \mathcal{C}$ , a  $T$ -coalgebra or coalgebra is a pair  $(X, c)$  where  $X$  is an object of  $\mathcal{C}$  and  $c$  is morphism from  $X$  to  $T(X)$ .

**Example 2.1.** Consider the category **Set** of sets and set functions, and let the functor  $T$  be given by  $T(X) = A \times X$  for a fixed set  $A$ . If  $f : X \rightarrow Y$ ,  $T(f) = 1_A \times f$  is the function from  $A \times X$  to  $A \times Y$  that sends an element  $\langle a, x \rangle$  to  $\langle a, f(x) \rangle$ . Then a coalgebra is a set  $X$ , together with a function  $c : X \rightarrow A \times X$  that assigns to each element in  $X$  a pair in  $A \times X$ . For example, if  $X = \{x, y\}$  and  $A = \{a, b\}$  then the map  $c(x) = \langle a, y \rangle, c(y) = \langle b, x \rangle$  makes  $X$  into a coalgebra.

**Example 2.2.** Automata can be modeled as coalgebras for the functor  $TX = X^A \times 2 \times 2$ , where

$X$  is the set of states of the automaton,  $A$  is the alphabet over which the automaton operates and  $2 = \{0, 1\}$  is used to mark which states are initial or final.

As a particular example, consider the automaton below, which recognizes the regular language  $\{a^n b | n \geq 0\}$ .



We can regard it as the coalgebra with  $X = \{x, y, z\}$ ,  $A = \{a, b\}$  and  $c : X \rightarrow X^A \times 2 \times 2$  given by:

$$c(x) = \langle (a \mapsto x, b \mapsto y), 1, 0 \rangle$$

$$c(y) = \langle (a \mapsto z, b \mapsto z), 0, 1 \rangle$$

$$c(z) = \langle (a \mapsto z, b \mapsto z), 0, 0 \rangle$$

So this indicates, for example, that  $x$  is an initial state,  $y$  is final, and  $z$  is neither initial nor final.

**Example 2.3.** Let  $\mathcal{D}$  be the functor in **Set** that assigns to each set  $X$ , the set of all functions  $\mu : X \rightarrow [0, 1]$  such that  $\mu$  has finite support and  $\sum_{x \in X} \mu(x) = 1$ . Given a function  $f : X \rightarrow Y$ , and  $\mu \in \mathcal{D}X$ ,  $(\mathcal{D}f)(\mu)(y) = \sum_{f(x)=y} \mu(x)$  if  $y \in f(X)$ , and 0 otherwise. Then a coalgebra for  $\mathcal{D}$  can be seen as a model for (discrete time, finitely branching) Markov Chains, and variations of this lead to diverse kinds of *probabilistic transition systems*. See for example, [LS91, Mos99, BSdV03].

For  $\mu \in \mathcal{D}X$ , we denote with  $\text{Supp}\mu$  the set of  $x \in X$  such that  $\mu(x) > 0$ . Notice that if  $f : X \rightarrow Y$  and  $\mu \in \mathcal{D}X$ , then

$$|\text{Supp}((\mathcal{D}f)\mu)| \leq |\text{Supp}(\mu)|. \quad (2.1)$$

**Example 2.4.** Given a set  $E$ , we can consider for each set  $X$  the set of all functions from  $E$  to



$X$ , denoted as  $X^E$ . If  $f : X \rightarrow Y$ , then we can define  $f^E : X^E \rightarrow Y^E$  by  $f^E(\alpha) = f \circ \alpha$  for every  $\alpha \in X^E$ . Thus  $\cdot^E$  is a functor in **Set**.

**Definition 2.2.** Given two coalgebras  $(X, c)$  and  $(Y, d)$ , a *T-coalgebra morphism* is a morphism  $f : X \rightarrow Y$  such that  $Tf \circ c = d \circ f$ , i.e., the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ c \downarrow & & \downarrow d \\ T(X) & \xrightarrow{T(f)} & T(Y) \end{array}$$

The notion of  $T$ -coalgebra morphism agrees with the previously existing notions of morphisms for automata and probability transition systems. That is, the theory of coalgebras captures nicely the notion of morphism in many different examples, including the automata from Example 2.2.

The collection of all coalgebras for a functor  $T$  on  $\mathcal{C}$ , together with the  $T$ -coalgebra morphisms form a category, usually denoted as  $\mathcal{C}_T$ .

## 2.2 Limits

**Definition 2.3.** (following [AHS90])

- A *diagram* in a category  $\mathcal{C}$  is a functor  $D : \mathcal{J} \rightarrow \mathcal{C}$ . The domain,  $\mathcal{J}$  is called the *scheme* of the diagram. For each object  $i$  of  $\mathcal{J}$ , we denote  $D(i)$  with  $D_i$ .
- A *source* in  $\mathcal{C}$  is a pair  $(A, (f_i)_{i \in I})$  consisting of an object  $A$  in  $\mathcal{C}$  and a family of morphisms  $f_i : A \rightarrow A_i$  indexed by some class  $I$ . In particular, we are going to consider  $I$  to be the class of objects in the category  $\mathcal{J}$ .  $A$  is called the *domain of the source* and the family  $(A_i)_{i \in I}$  is called the *codomain of the source*.

- A  $\mathcal{C}$  source  $(A \xrightarrow{f_i} D_i)$  is said to be *natural* for  $D$  provided that for each  $\mathcal{J}$ -morphism  $i \xrightarrow{d} j$  the triangle in  $\mathcal{C}$

$$\begin{array}{ccc} A & & \\ \downarrow f_i & \searrow f_j & \\ D_i & \xrightarrow{Dd} & D_j \end{array}$$

commutes. Natural sources are also called *cones*.

- A *limit* of  $D$  is a natural source  $(L \xrightarrow{l_i} D_i)_{i \in I}$  for  $D$  with the universal property that each natural source  $(A \xrightarrow{f_i} D_i)_{i \in I}$  for  $D$  uniquely factors through it; i.e., for every such source there exists a unique morphism  $f : A \rightarrow L$  with  $f_i = l_i \circ f$  for each  $i \in I$ . This limit is also called *projective limit* or *inverse limit*.
- A *weak limit* of  $D$  is a natural source  $(L \xrightarrow{l_i} D_i)_{i \in I}$  for  $D$  with the property that each natural source  $(A \xrightarrow{f_i} D_i)_{i \in I}$  for  $D$  factors through it; i.e., for every such source there exists a (not necessarily unique) morphism  $f : A \rightarrow L$  with  $f_i = l_i \circ f$  for each  $i \in I$ .
- We say that a functor  $T$  *preserves* a limit  $(L \xrightarrow{l_i} D_i)_{i \in I}$  if  $(TL \xrightarrow{Tl_i} TD_i)_{i \in I}$  is also a limit.


We will use two particular cases of limits: pullbacks and  $\omega^{op}$ -limits.

**Example 2.5.** *Pullbacks* are limits of diagrams:

$$\begin{array}{ccc} & & 1 \\ & & \downarrow f \\ 2 & \xrightarrow{g} & 3 \end{array}$$

Thus a pullback is an object  $L$  together with morphisms  $l_1 : L \rightarrow D_1$  and  $l_2 : L \rightarrow D_2$  such that  $f \circ l_1 = g \circ l_2$  (notice that  $l_3 : L \rightarrow D_3$  is usually ignored in the definition of pullback, because it's determined by  $l_3 = f \circ l_1 = g \circ l_2$ ), and for any object  $A$  and morphisms  $f_i : A \rightarrow D_i$ , such that

$f \circ f_1 = g \circ f_2$  there exists a unique morphism  $t : A \rightarrow L$  such that  $l_i \circ t = f_i$  for  $i = 1, 2$ .



In the category **Set**, pullbacks exist and are constructed as follows: Let  $L = \{\langle d_1, d_2 \rangle \in D_1 \times D_2 : (Df)(d_1) = (Dg)(d_2)\}$ . Then  $L$ , with the projections  $\pi_1 : L \rightarrow D_1$  and  $\pi_2 : L \rightarrow D_2$ , is a pullback.

**Example 2.6.**  $\omega^{op}$ -limits have as scheme the category obtained from the poset of the natural numbers with the reverse order:

$$0 \xleftarrow{f_0} 1 \xleftarrow{f_1} 2 \xleftarrow{\quad} \dots$$

We will call the diagrams of this scheme  $\omega^{op}$ -sequences, and denote them by the sequence of objects and morphisms  $(D_n, Df_n)_{n \in \omega}$  of which they are composed.

The limits of  $\omega^{op}$  sequences in **Set** can be constructed by taking the subset  $P$  of the product of all the  $D_n$ 's:

$$P = \{(d_n)_n \in \prod_{n \geq 0} D_n : (Df_n)(d_{n+1}) = d_n\} \quad (2.2)$$

Then the restrictions to  $P$  of the projections  $\pi_n : \prod_m D_m \rightarrow D_n$  make  $P$  the limit of the sequence.

## 2.3 Bisimulations

**Definition 2.4.** For an endofunctor  $T$  in  $\mathbf{Set}$ , a *bisimulation* between two  $T$ -coalgebras  $(X, c)$  and  $(Y, d)$  is a binary relation  $R \subseteq X \times Y$  such that there exists a morphism  $b : R \rightarrow T(R)$  that makes the two projections,  $\pi_X$  and  $\pi_Y$  into  $T$ -coalgebra morphisms.

$$\begin{array}{ccccc}
 X & \xleftarrow{\pi_X} & R & \xrightarrow{\pi_Y} & Y \\
 \downarrow c & & \downarrow b & & \downarrow d \\
 T(X) & \xleftarrow{T\pi_X} & T(R) & \xrightarrow{T\pi_Y} & T(Y)
 \end{array}$$

**Definition 2.5.** An endofunctor  $T$  *preserves weak pullbacks* if it transforms weak pullbacks into weak pullbacks.

**Example 2.7.** Many of the usually considered endofunctors in  $\mathbf{Set}$  preserve weak pullbacks. For example, the constant functor for any fixed set  $A$ , the exponential functor from Example 2.1, and the power set functor. If two functors  $U$  and  $V$  preserve weak pullbacks, then so does their composition  $U \circ V$  and their product  $U \times V$  (see [Gum99]).

On the other hand, not all functors in  $\mathbf{Set}$  preserve weak pullbacks. The functor  $(\cdot)_2^3$  defined by

$$X_2^3 = \{\langle x, y, z \rangle \mid |\{x, y, z\}| \leq 2\}$$

and

$$f_2^3(\langle x, y, z \rangle) = \langle f(x), f(y), f(z) \rangle$$

does not (see [AM89]).

**Definition 2.6.** A functor  $T$  is said to *weakly preserve pullbacks* if it transforms pullbacks into weak pullbacks.

In a category with pullbacks, preserving weak pullbacks and weakly preserving pullbacks are equivalent [Gum01]. The importance of this condition in the theory of coalgebra is explained by the following facts (among others).

**Theorem 2.1.** [Rut00] *Let  $T : \mathbf{Set} \rightarrow \mathbf{Set}$  be a functor that preserves weak pullbacks; and let  $f : (X, c) \rightarrow (Z, e)$  and  $g : (Y, d) \rightarrow (Z, e)$  be  $T$ -coalgebra morphisms. Then the pullback  $(P, k, l)$  of  $f$  and  $g$  in  $\mathbf{Set}$  is a bisimulation on  $X$  and  $Y$ .*

**Theorem 2.2.** [Rut00] *If the functor  $T$  preserves weak pullbacks, and  $X, Y$  and  $Z$  are  $T$ -coalgebras, then the composition  $R \circ Q$  of two bisimulations  $R \subseteq X \times Y$  and  $Q \subseteq Y \times Z$  is a bisimulation between  $X$  and  $Z$ .*

**Definition 2.7.** A *final* (or *terminal*) object in a category  $\mathcal{C}$  is an object  $1$  such that for every object  $A$  in  $\mathcal{C}$  there exists a unique morphism  $!_A : A \rightarrow 1$ .

It follows from the definition that all terminal objects in a category are isomorphic: if  $1$  and  $1'$  are terminal, then there are unique morphisms  $!_{1'} : 1' \rightarrow 1$  and  $!_1 : 1 \rightarrow 1'$ . Their composition  $!_{1'} \circ !_1$  must be  $!_1$ , which must also be the identity on  $1$ . In a similar fashion,  $!_1 \circ !_{1'}$  is the identity on  $1'$ , so  $1$  and  $1'$  are isomorphic.

**Example 2.8.** In the category  $\mathbf{Set}$ , any singleton is a final object. Since all terminal objects are isomorphic, we can select an arbitrary singleton  $1 = \{*\}$  and call it *the* terminal object of  $\mathbf{Set}$ .

Final objects in the category of coalgebras for a functor  $T$  are called final coalgebras, and they have some remarkable properties that make them interesting.

**Theorem 2.3.** (Lambek's Lemma, [Lam68]) *If  $(Z, e)$  is a final coalgebra, then  $e$  is an isomorphism between  $Z$  and  $TZ$ .*

**Theorem 2.4.** (Coinduction proof principle) [Rut00] *If there exists a bisimulation  $R \subseteq Z \times Z$  in a final coalgebra  $Z$  for a functor on  $\mathbf{Set}$ , and a pair  $\langle z, z' \rangle \in R$ , then  $z = z'$ .*

## 2.4 Final Sequences and Final Coalgebras

The main result in this section is Theorem 2.5, due to Smyth and Plotkin, [SP82], which provides a construction of final coalgebras, but we also present an original example showing that the construction cannot always be applied.

**Definition 2.8.** The *final sequence* for a functor  $T$  in a category with final object  $1$  is the one obtained by applying the functor  $T$  to  $1$ . Since  $1$  is final, there is a unique map  $!_{T1} : T1 \rightarrow 1$ , and we can apply  $T$  to this map, obtaining an infinite chain:

$$1 \xleftarrow{!} T1 \xleftarrow{T!} T^2 1 \xleftarrow{T^2!} T^3 1 \xleftarrow{\dots} \dots$$

This is a particular case of an  $\omega^{op}$ -sequence.

Given a  $T$ -coalgebra  $(X, c)$ , there is a way of regarding  $X$  as a natural source  $(X \xrightarrow{h_n^c}, T^n 1)_{n \in \omega}$  mapping  $X$  to the final sequence. Let  $h_0^c$  be the unique map  $!_X : X \rightarrow 1$ ,

$$h_0^c = !_X. \tag{2.3}$$

Given  $h_n^c : X \rightarrow T^n 1$ , let

$$h_{n+1}^c = Th_n^c \circ c. \tag{2.4}$$

**Lemma 2.1.** *Given a  $T$ -coalgebra  $(X, c)$ , for all  $n \geq 0$ ,  $h_n^c = T^n! \circ h_{n+1}^c$ .*

*Proof.* Easy to prove by induction on  $n$ : for  $n = 0$ ,  $h_0^c = !_X = !_X \circ h_1^c$  and if  $h_{n-1}^c = T^{n-1}! \circ h_n^c$ , then  $h_n^c = Th_{n-1}^c \circ c = T(T^{n-1}! \circ h_n^c) \circ c = T^n! \circ Th_n^c \circ c = T^n! \circ h_{n+1}^c$ .  $\square$

In the case of the category **Set**, the Lemma above says that for every element  $x$  of a  $T$ -coalgebra

$(X, c)$ ,  $h^c(x)$  is in the projective limit  $P$  defined as in (2.2).

$$\begin{array}{ccc}
 X & \xrightarrow{c} & TX \\
 h_n^c \downarrow & \searrow h_{n+1}^c & \downarrow Th_n^c \\
 T^n 1 & \xleftarrow{T^n!} & T^{n+1} 1
 \end{array}$$

Let  $h^c : X \rightarrow P$  be the function defined by

$$\pi_n(h^c(x)) = h_n^c(x) \quad (2.5)$$

for all  $n \geq 0$ . For a category in general,  $h^c$  is the mediating morphism given by the condition on  $P$  being a limit.

**Lemma 2.2.** *Given a  $T$ -coalgebra morphism  $f : (X, c) \rightarrow (Y, d)$ , for all  $n \geq 0$ ,  $h_n^d \circ f = h_n^c$  and therefore,  $h^d \circ f = h^c$ .*

*Proof.* We prove by induction over  $n$  that  $\pi_n h^d f = \pi_n h^c$ , this is,  $h_n^d \circ f = h_n^c$ :

For 0,  $h_0^d f = !_Y f = !_X = h_0^c$ . Assuming that  $h_n^d f = h_n^c$ , we get that  $h_{n+1}^d f = Th_n^d \circ d \circ f = Th_n^d \circ T f \circ c = T(h_n^d \circ f) \circ c = Th_n^c \circ c = h_{n+1}^c$ .  $\square$

We will ignore the superindex of  $h$  whenever doing so does not lead to any confusion.

For any  $n < m$ , let  $\tau_{mn} : T^m 1 \rightarrow T^n 1$  be defined by  $\tau_{mn} = T^n! \circ T^{n+1}! \circ \dots \circ T^{m-1}!$ . Also let  $\tau_{mm} = 1_{T^m 1}$ . We have

$$\begin{array}{ccccccc}
 & & \tau_{mn} & & & & \\
 & & \curvearrowright & & & & \\
 T^n 1 & \xleftarrow{T^n!} & T^{n+1} 1 & \xleftarrow{T^{n+1}!} & \dots & \xleftarrow{T^{m-1}!} & T^m 1
 \end{array}$$

**Theorem 2.5.** [SP82] *Let  $\mathcal{C}$  be a category with final object  $1$ , and  $T$  an endofunctor of  $\mathcal{C}$ . Suppose that  $P$  is the limit of the final sequence and that  $T$  preserves the limit. Then the final  $T$ -coalgebra exists and is  $(P, \rho)$ , where  $\rho : P \rightarrow TP$  is the mediating morphism obtained by  $TP$  being a limit.*

*Proof.* Let  $(X, c)$  be an arbitrary  $T$ -coalgebra, and assume there exists a  $T$ -coalgebra morphism  $f : (X, c) \rightarrow (P, \rho)$ . By the proof of Lemma 2.2,  $h_n^\rho \circ f = h_n^c$  for all  $n \geq 0$ . Since  $P$  is a limit, if such  $f$  exists, it must be unique. To prove its existence, it's enough to notice that  $h^c$  gives a natural source over the final sequence, so there must be a mediating morphism  $f : X \rightarrow P$ .

To prove that  $f$  is indeed a  $T$ -coalgebra morphism, note that  $h_n^\rho = \pi_n$ , and for all  $n \geq 0$  we have that  $T\pi_n \circ \rho \circ f = Th_n^\rho \circ \rho \circ f = h_{n+1}^\rho \circ f = h_{n+1}^c = Th_n^c \circ c = T(h_n^\rho \circ f) \circ c = T\pi_n \circ Tf \circ c$ , so  $\rho \circ f = Tf \circ c$ .  $\square$

The functors that preserve the  $\omega^{op}$ -limits as in the Theorem above are called  $\omega^{op}$ -continuous. Not all the functors in **Set** are  $\omega^{op}$ -continuous. Here are two important examples:

**Example 2.9.** ([Wor99]) Let  $\mathcal{P}$  be the (covariant) finite powerset functor. Let  $P$  be again the limit of the final sequence, and let  $A_n = \mathcal{P}^n 1$ . There is a unique morphism  $g : \mathcal{P}P \rightarrow P$  such that for all  $n \geq 1$ ,  $\pi_n g = \mathcal{P}\pi_{n-1}$ . If  $g$  were an isomorphism, then for some  $S \in \mathcal{P}P$ ,  $g(S) = \langle A_n \rangle$ . This sequence of elements is in  $P$  since  $A_n \in A_{n+1}$  and  $\mathcal{P}^n!(A_n) = A_{n-1}$ . But now we have that  $\pi_n g(S) \pi_n g(S) = A_{n-1}$  so  $|S| \geq |\pi_n g(S)| = |A_{n-1}|$  for all  $n \geq 1$ , which contradicts the fact that  $S$  is finite.

**Example 2.10.** Consider the functor  $\mathcal{D}$  on **Set**, the two element set  $A = \{a, b\}$  and the  $\omega^{op}$ -sequence

$$1 \xleftarrow{k_0} A \xleftarrow{k_1} A^2 \xleftarrow{k_2} A^3 \xleftarrow{\quad} \dots$$

where  $k_n : A_{n+1} \rightarrow A_n$  is the projection of the first  $n$  components. Let  $\mu_n \in \mathcal{D}A^n$  be the constant



function  $\frac{1}{2^n}$ . It's clear that for all  $n \geq 0$ ,  $(\mathcal{D}k_{n-1})\mu_n = \mu_{n-1}$ . The limit of the chain is  $A^\omega$ , and letting  $\pi_n : A^\omega \rightarrow A^n$  be the projection of the first  $n$  components, it's clear that there is no  $\mu \in \mathcal{D}A^\omega$  such that for all  $n \geq 0$ ,  $(\mathcal{D}\pi_n)\mu = \mu_n$ .

The example above shows that  $\mathcal{D}$  does not preserve  $\omega^{op}$ -sequences in general. But it could be the case that it preserves the limit of the final sequence, which is all that's required for applying Theorem 2.5. In fact, this is true, but not very interesting, since  $\mathcal{D}1$  consists of a single probability distribution (the one that assigns 1 to the single element in the set  $1 = \{*\}$ ). However, if we are interested in more complex functors, like the ones used to model labeled transition systems, we can adapt the counterexample to the final sequence of the functor, as in the following example.

**Example 2.11.** Let  $A = \{a, b\}$  and let  $S = A^\omega$ , the product of countably many copies of  $A$ . Let  $T$  be the functor  $TX = \mathcal{D}(S \times X)$ . Next we define on  $S$  the measures  $\nu_n$  by

$$\nu_n(s) = \begin{cases} \frac{1}{2^n} & \text{if } s_{n+i} = a \text{ for all } i \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

Now let  $\kappa_n$  be the product measure on  $S^n$

$$\kappa_n = \nu_n \times \nu_n \times \dots \times \nu_n. \tag{2.6}$$

Note that the support of  $\kappa_n$  has  $(2^n)^n$  elements.

The next step of this construction is to embed  $S^n$  in  $S \times T^{n-1}1$  for  $n \geq 1$ . Some notation first: given a set  $X$  and  $x \in X$ ,  $\delta_x \in \mathcal{D}X$  is the probability distribution:

$$\delta_x(y) = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{otherwise} \end{cases}$$

We can consider  $\delta$  as a natural transformation from the identity functor  $Id$  to  $\mathcal{D}$ . It is easy to see it is an injective one, that is, for every set  $X$ ,  $\delta_X : X \rightarrow \mathcal{D}X$  is injective. Furthermore,  $\delta$  is the unit of the monad for  $\mathcal{D}$ , with multiplication  $\gamma_X : \mathcal{D}\mathcal{D}X \rightarrow \mathcal{D}X$  given by

$$\gamma_X(\mu)(x) = \sum_{\nu \in \mathcal{D}X} \nu(x)\mu(\nu)$$

for all  $\mu \in \mathcal{D}\mathcal{D}X$  and  $x \in X$ . This is a particular case of what's known as the Giry monad (see [Gir82] and Lemma 3.2 of this dissertation).

Now we define the maps  $a_n : S^n \rightarrow S \times T^{n-1}1$  for  $n \geq 1$  as follows:  $a_1 : S \rightarrow S \times 1 = S \times T^01$  is

$$a_1(s) = \langle s, * \rangle \tag{2.7}$$

$$a_{n+1}(s_0, s_1, \dots, s_n) = \langle s_0, \delta_{a_n(s_1, \dots, s_n)} \rangle \tag{2.8}$$

In other words,  $a_{n+1}$  is the map:

$$S^{n+1} = S \times S^n \xrightarrow{1_S \times a_n} S \times (S \times T^{n-1}1) \xrightarrow{1_S \times \delta_{S \times T^{n-1}}} S \times \mathcal{D}(S \times T^{n-1}) = S \times T^n 1$$

The maps  $a_n$  allow us to define

$$\gamma_n = (\mathcal{D}a_n)\kappa_n \tag{2.9}$$

Since  $\mathcal{D}a_n : \mathcal{D}S^n \rightarrow \mathcal{D}(S \times T^{n-1})1 = T^n 1$ ,  $\gamma_n \in T^n 1$ . Furthermore, since  $\delta$  is injective, it's easy to check that so is  $a_n$  for all  $n$ . As a consequence,

$$|Supp(\gamma_n)| = |Supp(\kappa_n)| = (2^n)^n = 2^{(n^2)} \tag{2.10}$$

Now let  $k_n : S^{n+1} \rightarrow S^n$  be the map sending  $(s_0, s_1, \dots, s_n)$  to  $(s_0, \dots, s_{n-1})$  and consider the following diagram for  $n \geq 1$ :

$$\begin{array}{ccccc}
 \dots & \xleftarrow{k_{n-1}} & S^n & \xleftarrow{k_n} & S^{n+1} & \xleftarrow{\dots} & \dots \\
 & & \downarrow a_n & & \downarrow a_{n+1} & & \\
 \dots & \xleftarrow{\dots} & S \times T^{n-1}1 & \xleftarrow{1_S \times T^{n-1}!} & S \times T^n 1 & \xleftarrow{\dots} & \dots
 \end{array} \tag{2.11}$$

We prove by induction that the square commutes. For  $n = 1$ , we have that  $a_1(k_1(s_0, s_1)) = a_1(s_0) = \langle s_0, * \rangle$ , while  $1_S \times!(a_2(s_0, s_1)) = 1_S \times! \langle s_0, \delta_{\langle s_1, * \rangle} \rangle = \langle s_0, * \rangle$ , so

$$\begin{array}{ccc}
 S & \xleftarrow{k_1} & S^2 \\
 \downarrow a_1 & & \downarrow a_2 \\
 S \times T^0 1 & \xleftarrow{1_S \times!} & S \times T^1 1
 \end{array}$$

commutes.

For the inductive step, note that the square in (2.11) can be rewritten as follows, using the observation that  $k_n = 1_S \times k_{n-1}$ :

$$\begin{array}{ccc}
 S \times S^{n-1} & \xleftarrow{1_S \times k_{n-1}} & S \times S^n \\
 \downarrow 1_S \times a_{n-1} & & \downarrow 1_S \times a_n \\
 S \times (S \times T^{n-2} 1) & \xleftarrow{1_S \times (1_S \times T^{n-2}!)} & S \times (S \times T^{n-1} 1) \\
 \downarrow 1_S \times \delta_{S \times T^{n-2} 1} & & \downarrow 1_S \times \delta_{S \times T^{n-1} 1} \\
 S \times (S \times T^{n-1} 1) & \xleftarrow{1_S \times T^{n-1}!} & S \times T^n 1
 \end{array}$$

So the problem reduces to proving that the following diagram commutes:

$$\begin{array}{ccc}
 S^{n-1} & \xleftarrow{k_{n-1}} & S^n \\
 a_{n-1} \downarrow & & \downarrow a_n \\
 S \times T^{n-2} \mathbf{1} & \xleftarrow{1_S \times T^{n-2} \mathbf{1}} & S \times T^{n-1} \mathbf{1} \\
 \delta_{S \times T^{n-2} \mathbf{1}} \downarrow & & \downarrow \delta_{S \times T^{n-1} \mathbf{1}} \\
 T^{n-1} \mathbf{1} & \xleftarrow{T^{n-1} \mathbf{1}} & T^n \mathbf{1}
 \end{array}$$

But in this diagram the upper square commutes by inductive hypothesis, and the lower one because of the naturality of the natural transformation  $\delta$ , and using the fact that  $\mathcal{D}(1_S \times T^{n-2} \mathbf{1}) = T^{n-1} \mathbf{1}$ .

Now that we know that (2.11) commutes, we can apply the functor  $\mathcal{D}$  and get that:

$$\begin{array}{ccc}
 \dots & \xleftarrow{\mathcal{D}k_{n-1}} \mathcal{D}S^n & \xleftarrow{\mathcal{D}k_n} \mathcal{D}S^{n+1} \xleftarrow{\dots} \\
 & \downarrow \mathcal{D}a_n & \downarrow \mathcal{D}a_{n+1} \\
 \dots & \xleftarrow{T^n \mathbf{1}} T^n \mathbf{1} & \xleftarrow{T^{n+1} \mathbf{1}} T^{n+1} \mathbf{1} \xleftarrow{\dots}
 \end{array} \tag{2.12}$$

also commutes. Therefore:

$$\begin{aligned}
 T^n \mathbf{1} \gamma_{n+1} &= T^n \mathbf{1} (\mathcal{D}a_{n+1}) \kappa_{n+1} && \text{by (2.9)} \\
 &= \mathcal{D}((1_S \times T^{n-1} \mathbf{1}) a_n) \kappa_{n+1} \\
 &= \mathcal{D}(a_{n-1} k_n) \kappa_{n+1} && \text{by (2.12)} \\
 &= (\mathcal{D}a_{n-1}) (\mathcal{D}k_n) \kappa_{n+1} \\
 &= (\mathcal{D}a_{n-1}) \kappa_n \\
 &= \gamma_n && \text{by (2.9)}
 \end{aligned}$$

To summarize: we have  $\gamma_n \in T^n \mathbf{1}$  with  $T^n \mathbf{1} \gamma_{n+1} = \gamma_n$  for all  $n \geq 0$ , so the sequence  $\Gamma =$

---

$(\gamma_0, \gamma_1, \dots)$  is in the limit  $P$  of the final sequence. If we assume that  $T$  preserves this limit, then there exists  $\rho : P \rightarrow TP$  so that for all  $n \geq 0$ ,  $T\pi_n \circ \rho = \pi_{n+1}$ . Then,  $\rho(\Gamma) \in TP = \mathcal{D}(S \times P)$ , and therefore  $\rho(\Gamma)$  has finite support. On the other hand, by equations (2.1) and (2.10),  $2^{(n^2)} \leq |\text{Supp}(\rho(\Gamma))|$  for all  $n$ , contradiction.

We will present in Chapter 6 a construction that allows us to find the final coalgebra for this functor  $T$ .

# 3

---

## Measurable Spaces

### 3.1 Measure Theory

**Definition 3.1.** A *ring* (or *boolean ring*) of sets is a non empty class  $\mathcal{B}$  of sets such that if  $E \in \mathcal{B}$  and  $F \in \mathcal{B}$  then  $E \cup F \in \mathcal{B}$  and  $E \setminus F \in \mathcal{B}$ . If furthermore, the class  $\mathcal{B}$  is closed under countable unions, then it's a  $\sigma$ -*ring*.

An *algebra* (or *boolean algebra*) of subsets of a set  $M$  is a nonempty class  $\mathcal{B}$  of subsets of  $M$  such that if  $E \in \mathcal{B}$  and  $F \in \mathcal{B}$  then  $E \cup F \in \mathcal{B}$  and  $E^c \in \mathcal{B}$ , where  $E^c$  denotes the complement of  $E$  with respect to  $M$ . It follows that  $M \in \mathcal{B}$ . It is also a  $\sigma$ -*algebra* if it is closed under countable unions.

**Definition 3.2.** A *measurable space* is a pair  $(M, \Sigma)$  where  $M$  is a set and  $\Sigma$  is a  $\sigma$ -algebra of subsets of  $M$ . The subsets of  $M$  which are in  $\Sigma$  are called *measurable sets*.

A function between measurable spaces  $f : (M, \Sigma) \rightarrow (M', \Sigma')$  is said to be *measurable* if for every  $E \in \Sigma'$ ,  $f^{-1}(E) \in \Sigma$ . We will denote with  $\text{Meas}$  the category of measurable spaces and measurable functions.

**Definition 3.3.** A *measure* on a measurable space  $(M, \Sigma)$  is a map  $\mu : \Sigma \rightarrow [0, \infty]$  such that

$\mu(\emptyset) = 0$  and  $\mu(\bigcup_n E_n) = \sum_n \mu(E_n)$  for any pairwise disjoint family  $\{E_n\} \subseteq \Sigma$ . If  $\mu(M) = 1$ , then  $\mu$  is called a *probability measure*.

**Definition 3.4.** A *family of generators*  $\mathcal{F}$  of a  $\sigma$ -algebra  $\Sigma$  is a family of subsets such that the smallest  $\sigma$ -algebra including  $\mathcal{F}$  is  $\Sigma$ . This is denoted by  $\sigma(\mathcal{F}) = \Sigma$ .

We will consider the endofunctor  $\Delta$  in  $\text{Meas}$  that assigns to each measurable space  $M$ , the set  $\Delta M$  of all the probability measures over  $M$ , endowed with the  $\sigma$ -algebra  $\Sigma_\Delta$  generated by the sets of the form  $\beta^p(E)$  where  $p \in [0, 1]$ ,  $E$  is a measurable subset of  $M$  and

$$\beta^p(E) = \{\mu \in \Delta M : \mu(E) \geq p\}. \quad (3.1)$$

The same  $\sigma$ -algebra can be generated by considering only the sets  $\beta^p(E)$  for rational values of  $p$ .

If  $f : (M, \Sigma) \rightarrow (N, \Sigma')$  is measurable, we define  $\Delta f : \Delta M \rightarrow \Delta N$  as follows: for  $\mu \in \Delta(M)$  and  $E \in \Sigma'$ ,

$$(\Delta f)(\mu)(E) = \mu(f^{-1}(E)).$$

As an easy consequence of the definition, we have that

**Lemma 3.1.** *If  $f : M \rightarrow M'$  is a measurable function, then for all measurable  $E \subseteq M'$  and  $p \in [0, 1]$ ,*

$$\beta^p(f^{-1}(E)) = (\Delta f)^{-1}(\beta^p(E))$$

*Proof.* Both sets are  $\{\mu \in \Delta M \mid \mu(f^{-1}(E)) \geq p\}$ . □

From Lemma 3.1, it follows that  $\Delta f$  is measurable, and it's easy to check now that  $\Delta$  is a functor on  $\text{Meas}$ .

**Definition 3.5.** Given an element  $m$  in a measurable space  $M$ ,  $\delta_m \in \Delta M$  is the Dirac measure supported at  $m$ . It assigns 1 to every subset of  $M$  containing  $m$ , and 0 to the rest.

For each measurable space  $M$ , the map  $\delta_M : M \rightarrow \Delta M$  maps  $m \in M$  to  $\delta_M(m) = \delta_m \in \Delta M$ .

This application  $\delta_M$  is measurable: for every  $E \in \Sigma$ . If  $p \in (0, 1]$ ,

$$\begin{aligned} \delta^{-1}(\beta^p(E)) &= \{x \in X : \delta_x \in \beta^p(E)\} \\ &= \{x \in X : \delta_x(E) \geq p\} \\ &= \{x \in X : x \in E\} = E \end{aligned}$$

and if  $p = 0$  we have  $\delta^{-1}\beta^0(E) = X$ .

**Integration.** In a measurable space  $X$  it is usual to consider the real-valued functions  $\chi_E$  for measurable subsets  $E$ .  $\chi_E(x) = 1$  if  $x \in E$ , and 0 otherwise. A *simple function* is a function  $f = \sum_{i=1}^n \alpha_i \chi_{E_i}$  where  $E_1, \dots, E_n$  are disjoint measurable sets and  $\alpha_i \neq 0$  for  $i = 1, \dots, n$ . Given a measure  $\mu$  on  $X$ , if each  $E_i$  has finite measure (as is the case when  $\mu$  is a probability measure), then the simple function  $f$  is said to be *integrable* and  $\int_X f d\mu$  is defined to be  $\sum_{i=1}^n \alpha_i \mu(E_i)$ . The *distance* between two simple integrable functions  $f$  and  $g$  is  $\int_X |f - g| d\mu$ . More in general, if  $f$  is a measurable real-valued function, and there exists a sequence of simple integrable functions  $f_n$  that converge in measure to  $f$  (that is,  $\lim_n \mu(\{x \mid |f_n(x) - f(x)| > \epsilon\}) = 0$  for all  $\epsilon > 0$ ) such that the distance between  $f_n$  and  $f_m$  tends to 0 when  $m$  and  $n$  tend to infinity, then  $f$  is said to be integrable and its integral is defined to be the limit of the integrals of the simple functions  $f_n$  (see [Hal50], Chapter V for more specific details).



**The Giry Monad.**  $\delta : Id \rightarrow \Delta$  is a natural transformation. A second natural transformation  $\gamma : \Delta\Delta \rightarrow \Delta$  given by

$$\gamma_M(\mu)(E) = \int_{\nu \in \Delta(M)} \nu(E) d\mu,$$

gives us the following:

**Lemma 3.2 (Giry [Gir82]).**  $(\Delta, \delta, \gamma)$  is a monad on Meas.

**Products and coproducts.** Products and coproducts exist in Meas, and they can be constructed in a similar fashion to the corresponding constructions in Set (furthermore, Meas is a complete category, [Gir82]). The  $\sigma$ -algebra of the product is the one generated by the “rectangles” formed by taking the cartesian product of measurable sets, while the  $\sigma$ -algebra of the coproduct is formed by taking (disjoint) unions of measurable sets in each of the summands.

If  $M \times N$  is a product of measurable spaces and  $\mu$  is a probability measure on it,  $(\Delta\pi_M)\mu$  is a measure on  $M$  and it’s called the *marginal* of  $\mu$  over  $M$ . Of course,  $(\Delta\pi_N)\mu$  is the marginal of  $\mu$  over  $N$ . If on the other hand,  $\mu \in \Delta M$  and  $\nu \in \Delta N$ , it is possible to define a probability measure  $\mu \times \nu \in \Delta(M \times N)$  by taking, for measurable subsets  $E$  and  $F$  in  $M$  and  $N$ , respectively

$$\mu \times \nu(E \times F) = \mu(E)\nu(F)$$

this definition is then extended to all the measurable subsets of  $M \times N$ . It is clear that the marginals of  $\mu \times \nu$  are  $\mu$  and  $\nu$ .

If  $(M_i, \Sigma_i)_{i \in I}$  is a family of measurable spaces, then  $(M, \Sigma) = \prod_i (M_i, \Sigma_i)$  is also a measurable space, where  $\Sigma$  is taken to be the  $\sigma$ -algebra generated by the union of all the  $(\pi_i)^{-1}(\Sigma_i)$ , for all the projections  $\pi_i : M \rightarrow M_i$ .

### 3.2 Weak Pullbacks and the Functor $\Delta$

A natural question that arises in this context is whether the functor  $\Delta$  preserves weak pullbacks. Here we answer negatively the equivalent question of whether  $\Delta$  weakly preserves pullbacks.

Pullbacks in **Meas** are constructed in a similar way as in **Set**. Given measurable functions  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$ , one takes the set  $P = \{\langle x, y \rangle : f(x) = g(y)\}$  endowed with the smallest  $\sigma$ -algebra that makes the projections measurable.


Let's recall that on the real line, the *Borel sets* are defined to be those in the  $\sigma$ -algebra generated by the collection of all the open intervals  $(a, b)$  with  $a < b \in \mathbb{R}$ . The length of an interval gives us a set function  $\lambda((a, b)) = b - a$ , which can be extended to the completion of the Borel sets of  $\mathbb{R}$  (this is, all the unions of Borel sets with subsets of sets of measure zero). This extension is called the *Lebesgue measure*, and the sets in the completion are *Lebesgue measurable*. Its restriction to the interval  $[0, 1]$  yields a probability measure over that same interval.

Now let  $X = Y = Z$  be the interval  $[0, 1]$  on the reals. Let  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$  be the maps:

$$f(x) = \begin{cases} x & \text{if } x \neq 1/2 \\ 0 & \text{if } x = 1/2 \end{cases}$$

and  $g(y) = 1 - y$ . It is easy to check that  $f$  and  $g$  are measurable functions and  $\{\langle x, y \rangle \mid f(x) = g(y)\} = \emptyset$ .

Now consider  $Q = \{q\}$ , and functions  $k : Q \rightarrow \Delta X, l : Q \rightarrow \Delta Y$  given by  $k(q) = l(q)$  equal to the Lebesgue measure over  $[0, 1]$ .  $(\Delta f)k(q) = (\Delta g)l(q)$  is again the Lebesgue measure, so the diagram commutes, but there is no measurable function  $j : Q \rightarrow \emptyset$ .




This counterexample can be “fixed” by changing  $f$  to be the identity on  $[0, 1]$ . Could this be the case for all counterexamples? That is, could this be a pathological case arising from the fact that two measurable functions can be essentially the same but differ on a set of measure zero? In other words, is there a pair of functions  $\hat{f}, \hat{g}$  so that  $f = \hat{f}$  and  $g = \hat{g}$  almost everywhere, but yet  $\hat{f}$  and  $\hat{g}$  have a non-empty pullback that can be weakly preserved by  $\Delta$ ? The answer is again negative as the following construction shows.

We can assume that  $\hat{f} = 1_{[0,1]}$  and  $\hat{g} = f$  on a set  $E$  of measure zero, with  $\hat{g} = g$  everywhere else. So the pullback of  $\hat{f}$  and  $\hat{g}$  is  $P = \{\langle x, x \rangle : x \in E\}$ . For  $q \in [0, 1]$ , consider the functions  $c_q : [0, 1] \rightarrow \mathbb{R}$  defined as follows:

$$c_q(x) = \begin{cases} 1 + q & \text{if } x \in [0, 1/4) \cup (3/4, 1] \\ 1 - q & \text{if } x \in [1/4, 3/4] \end{cases}$$

The functions  $c_q$  are naturally associated with a measure in  $\Delta[0, 1]$ , by taking  $\int_F c_q(x) dx$  where  $dx$  represents the usual Lebesgue measure over the reals and  $F$  is the measurable set for which we are calculating the probability.

Now let  $Q = [0, 1]$  and  $k(q) = c_q = l(q)$ . It is clear that  $\Delta 1_{[0,1]} k(q) = c_q = \Delta \hat{g} l(q)$ . Furthermore, the functions  $k$  and  $l$  are measurable. Assume that there is a measurable function  $j : Q \rightarrow \Delta P$  such that the following diagram commutes



Then, when we calculate  $k(q)(E)$  we get:  $\int_E c_q(x)dx = 0$ , while on the other hand  $\Delta\pi_1 j(q)(E) = j(q)(\pi_1^{-1}(E)) = j(q)(P) = 1$ , contradiction.

We conclude that  $\Delta P$  is not a weak pullback for the functions  $\Delta\hat{f}$  and  $\Delta\hat{g}$ , and therefore  $\Delta$  does not weakly preserve pullbacks.

### 3.3 $\omega^{op}$ -limits and the Functor $\Delta$

To present an example where  $\Delta$  does not preserve an  $\omega^{op}$ -limit, we need first to introduce some more definitions from measure theory. In the first place,  $\omega^{op}$  limits can be constructed in **Meas** as follows. Suppose we have an  $\omega^{op}$ -sequence  $(A_n, k_n)_{n \in \omega}$ . Just as in **Set**, we take

$$P = \{(a_n) \in \prod_{n \geq 0} A_n : k_n(a_{n+1}) = a_n\}$$

Now, if  $\Sigma_n$  is the  $\sigma$ -algebra on  $A_n$ , notice that  $\pi_n^{-1}(\Sigma_n)$  is a  $\sigma$ -algebra on  $P$  and  $\pi_n^{-1}(\Sigma_n) \subseteq \pi_{n+1}^{-1}(\Sigma_{n+1})$  (this follows from  $k_n \circ \pi_{n+1} = \pi_n$ ). The  $\sigma$ -algebra we will use on  $P$  is

$$\Sigma = \sigma\left(\bigcup_{n \geq 0} \pi_n^{-1}(\Sigma_n)\right).$$

**Definition 3.6.** ([Hal50], p. 42) A class of sets  $\mathcal{H}$  is *hereditary* if  $E \in \mathcal{H}$  and  $F \subseteq E$  implies  $F \in \mathcal{H}$ .

The least hereditary family containing a ring  $\mathcal{R}$  will be denoted with  $\mathcal{H}(\mathcal{R})$ .

An *outer measure* is an extended real-valued, non negative, monotone and countably subadditive set function  $\mu^*$  defined on an hereditary  $\sigma$ -ring  $\mathcal{H}$  and such that  $\mu^*(\emptyset) = 0$ .

Extended real-valued functions can take the value  $+\infty$ , while countably subadditive means that if a sequence  $\{E_i\}$  of sets in  $\mathcal{H}$  whose union is also in  $\mathcal{H}$ , we have

$$\mu^*\left(\bigcup_{i=1}^{\infty} E_i\right) \leq \sum_{i=1}^{\infty} \mu^*(E_i)$$

**Theorem 3.1.** [Hal50] *If  $\mu$  is a measure on a ring  $\mathcal{R}$ , and if for every  $E \in \mathcal{H}(\mathcal{R})$ ,*

$$\mu^*(E) = \inf\left\{\sum_{n=1}^{\infty} \mu(E_n) : E_n \in \mathcal{R}, n = 1, 2, \dots; E \subseteq \bigcup_{n=1}^{\infty} E_n\right\}$$

*then  $\mu^*$  is an extension of  $\mu$  to an outer measure on  $\mathcal{H}(\mathcal{R})$ ; if  $\mu$  is finite, so is  $\mu^*$ .*

The outer measure  $\mu^*$  may be described as the lower bound of sums of the type  $\sum_{n=1}^{\infty} \mu(E_n)$  where  $\{E_n\}$  is a sequence of sets in  $\mathcal{R}$  whose union contains  $E$ .  $\mu^*$  is called the outer measure *induced* by  $\mu$ . In particular, if  $\mu$  is a measure defined on a  $\sigma$ -algebra  $\Sigma$ , then

$$\mu^*(E) = \inf\{\mu(F) : E \subseteq F \text{ and } F \in \Sigma\}$$

Analogously, the *inner measure*  $\mu_*$  induced by  $\mu$  can be defined, for every  $E \in \mathcal{H}(\mathcal{R})$  as

$$\mu_*(E) = \sup\{\mu(F) : F \subseteq E \text{ and } F \in \Sigma\}.$$

**Definition 3.7.** A subset  $X_0$  of a measure space  $(X, \Sigma, \mu)$  is *thick* if for every measurable set  $E$   $\mu_*(E - X_0) = 0$ . This is the same as saying that  $\mu_*(X - X_0) = 0$ . If  $\mu$  is totally finite (i.e.,  $\mu(X) < \infty$ ;

all probability measures are totally finite), then  $X_0$  is thick if and only if  $\mu^*(X_0) = \mu(X)$ .

In the case of the unit interval with the Lebesgue measure on it, thick just means a set with outer Lebesgue measure equal to 1. Note that the outer measure  $\lambda^*$  induced by  $\lambda$  is defined for every subset of the reals, while  $\lambda$  itself is not. On the Lebesgue measurable sets,  $\lambda^*$  and  $\lambda$  agree, so  $\lambda$  is a *regular* measure (as every measure over a metric space is; see Definition 3.11 and the comments before Theorem 3.4).

**Theorem 3.2.** ([Hal50], p. 75) *If  $X_0$  is a thick subset of a measure space  $(X, \Sigma, \mu)$ ,  $\Sigma_0 = \Sigma \cap X_0$  and for  $E \in \Sigma$ ,  $\mu_0(E \cap X_0) = \mu(E)$ , then  $(X_0, \Sigma_0, \mu_0)$  is a measure space.*

**Lemma 3.3.** *Let  $\lambda$  be the Lebesgue measure on the interval  $[0, 1]$  of the reals. There exist sets  $X_0 \supset X_1 \supset X_2 \supset \dots$  in  $[0, 1]$  with  $\lambda^*(X_k) = 1$  for all  $k$  and  $\bigcap_k X_k = \emptyset$ .*

The proof of this Lemma makes use of sets like the non-measurable one constructed by Vitali, and is sketched in [Dud89] Exercise 2, p. 81.

**Example 3.1.** We consider  $(X_n, \Sigma_n, \mu_n)$  to be defined as in the Theorem 3.2. Let  $i_n : X_{n+1} \hookrightarrow X_n$  be the inclusion map, and  $(Y_n, \Gamma_n) = \prod_{k=0}^n (X_k, \Sigma_k)$ .

Notice that for all  $n \geq 0$ ,  $(\Delta i_n)\mu_{n+1} = \mu_n$ . Indeed, if  $E_n \in \Sigma_n$ , then there is a Lebesgue measurable set  $E$  such that  $E_n = E \cap X_n$ . Then,

$$\begin{aligned} (\Delta i_n)(\mu_{n+1})(E_n) &= \mu_{n+1}(i_n)^{-1}(E_n) \\ &= \mu_{n+1}(i_n)^{-1}(E \cap X_n) \\ &= \mu_{n+1}(E \cap X_{n-1}) \\ &= \mu(E) \\ &= \mu_n(E_n) \end{aligned}$$

Let  $s_n : X_n \rightarrow Y_n$  be the application  $x \mapsto (x, \dots, x)$ . To see that  $s_n$  is measurable, consider the sets

$$D_n = \{y \in Y_n : y_0 = y_1 = \dots = y_n\}$$

$D_n$  is measurable in  $Y_n$ :

$$I_k = \bigcup_{j=1}^k \left( \prod_{i=0}^n \left( \left[ \frac{j-1}{k}, \frac{j}{k} \right] \cap X_i \right) \right)$$

is a series of “cubes” containing  $D_n$  and  $D_n = \bigcap_{k=1}^{\infty} I_k$ .

Now for any  $E \in \Gamma_n$ ,

$$\begin{aligned} s_n^{-1}(E) &= \{x \in X_n : (x, \dots, x) \in E\} \\ &= E \cap D_n \end{aligned}$$

and this is a measurable subset of  $Y_n$ .

Let  $k_n : Y_{n+1} \rightarrow Y_n$  be the measurable map sending  $(y_0, y_1, \dots, y_n, y_{n+1})$  to  $(y_0, y_1, \dots, y_n)$ , and let  $P$  be the projective limit of the chain formed by the spaces  $Y_n$  and the maps  $k_n, n \geq 0$ , with  $\pi_n : P \rightarrow Y_n$  as before.

It's easy to check that for all  $n \geq 0$ , the following diagram commutes:

$$\begin{array}{ccc} X_{n+1} & \xrightarrow{s_{n+1}} & Y_{n+1} \\ i_n \downarrow & & \downarrow k_n \\ X_n & \xrightarrow{s_n} & Y_n \end{array}$$

Let  $\nu_n = (\Delta s_n)\mu_n \in \Delta Y_n$ . We check that for all  $n \geq 0$ :

$$\begin{aligned} (\Delta k_n)(\nu_{n+1}) &= (\Delta k_n)((\Delta s_{n+1})\mu_{n+1}) \\ &= (\Delta s_n \circ i_n)\mu_{n+1} \\ &= (\Delta s_n)\mu_n \\ &= \nu_n \end{aligned}$$

Suppose toward a contradiction that  $\Delta$  preserved limits of  $\omega^{op}$ -chains, so that  $\Delta P$  is the limit of  $\Delta Y_0 \leftarrow \Delta Y_1 \leftarrow \dots$ . Then there would be some  $\nu \in \Delta P$  such that for all  $n$ ,  $(\Delta \pi_n)(\nu) = \nu_n$ . For each  $n$ , let

$$C_n = \pi_n^{-1}(s_n(X_n)) = \pi_n^{-1}(D_n).$$

This is the set of elements of  $X$  whose first  $n$  coordinates are equal.  $\bigcap_n C_n = \emptyset$ , since  $x \in \bigcap_n C_n$  iff for all  $n$ ,  $x_n \in D_n$  iff  $x_0 \in X_n$  for all  $n$ , but we know that  $\bigcap_n X_n = \emptyset$ .

Moreover, each set  $C_n$  is measurable, being the inverse images under  $\pi_n$  of  $D_n$ . But note that

$$\begin{aligned} \nu(C_n) &= \nu(\pi_n^{-1}(s_n(X_n))) \\ &= (\Delta \pi_n)\nu(s_n(X_n)) \\ &= \nu_n(s_n(X_n)) \\ &= (\Delta s_n)\mu_n(s_n(X_n)) \\ &= \mu_n((s_n)^{-1}s_n(X_n)) \\ &= \mu_n(X_n) \\ &= 1 \end{aligned}$$

So  $\nu(C_n) = 1$  for all  $n$ , yet  $\bigcap_n C_n = \emptyset$ . This contradiction shows that  $\nu$  cannot exist.



### 3.4 Kolmogorov's consistency Theorem

We have seen that the functor  $\Delta$  does not preserve  $\omega^{op}$ -limits in general, but under certain extra topological assumptions, however, functors closely related to  $\Delta$  do preserve the  $\omega^{op}$ -limits. This is usually known as Kolmogorov's consistency theorem.

**Definition 3.8.** Given a topological space  $(X, \mathcal{T})$ , a *Borel measure* over  $X$  is a measure over the measurable space  $(X, \sigma(\mathcal{T}))$ .

Whenever  $X = (X, \mathcal{T})$  is a topological space, the  $\sigma$ -algebra we will consider over it is  $\sigma(\mathcal{T})$ . The sets in this  $\sigma$ -algebra are called the *Borel sets* of  $X$ , and  $\Delta X$  will be the space of all Borel probability measures over  $X$ .

**Definition 3.9.** A set  $F$  is the *support* of a probability  $\mu$  on a topological space (denoted  $F = \text{Supp}(\mu)$ ) if  $F$  is the smallest closed set whose complement has measure 0.

Note that the support of a measure  $\mu$  need not exist.

Let's recall that a *Polish space* is a separable and completely metrizable topological space. That is, it admits a countable dense set and a metric compatible with the topology such that the space is complete under that metric. The usual examples of Polish spaces are the real line and the Cantor space. Any metric space that is not complete, like the rational numbers, fails to be Polish.

**Definition 3.10.** A measurable space  $(X, \Sigma)$  is a *standard Borel space* if there is a Polish topology  $\mathcal{T}$  on  $X$  such that  $\sigma(\mathcal{T}) = \Sigma$ .

If  $X$  is a standard Borel space, so is  $\Delta X$ , and the projective limit of standard Borel spaces is also a standard Borel space.

**Theorem 3.3.** (see [Par67], [Kec95]) Let  $(X_n, \Sigma_n)$  be a sequence of standard Borel measurable spaces and  $f_n : X_{n+1} \rightarrow X_n$  a surjective measurable map for  $n \geq 0$ . If  $\mu_n$  is a probability measure

on  $X_n$  such that  $\mu_{n+1}f_n^{-1} = \mu_n$  for all  $n \geq 0$ , then there is a unique probability measure  $\mu$  on  $(X, \Sigma)$  the projective limit of the  $(X_n, \Sigma_n)$  such that for all  $n \geq 0$ ,  $\mu\pi_n^{-1} = \mu_n$ .

In terms of our previous notation,  $(\mu_n)_{n \in \omega}$  is an element of the projective limit of the sequence formed by the spaces  $\Delta X_n$  and the functions  $\Delta f_n$ . Then there exists  $\mu \in \Delta P$ , where  $P$  is the projective limit of the sequence of spaces  $X_n$  and functions  $f_n$ , such that  $(\Delta\pi_n)\mu = \mu_n$ . This establishes an isomorphism between the projective limit of the spaces  $\Delta X_n$  and  $\Delta P$ . In other words, we get the following Corollary:

**Corollary 3.1.** *In the category of standard Borel spaces and measurable functions, the functor  $\Delta$  is  $\omega^{op}$ -continuous.*

There are other sets of conditions under which Kolmogorov's theorem holds. To point them out we need first to recall a few definitions.

**Definition 3.11.** A Borel measure  $\mu$  is said to be *regular* if for any  $\mu$ -measurable set  $E$ ,

$$\begin{aligned} \mu(E) &= \inf\{\mu(O) : O \text{ is open and } E \subseteq O\} \\ &= \sup\{\mu(C) : C \text{ is compact and } E \supseteq C\}. \end{aligned}$$

We will denote with  $\Delta_r(X)$  the set of all the regular Borel measures over a topological space  $X$ .

**Definition 3.12.** Given a topological space  $X$ ,  $C(X)$  is the normed space of all continuous real-valued functions on  $X$ , with the supremum norm.

$C_b(X)$  is the space of all the *bounded* continuous real-valued functions.

It is well known that  $C_b(X)$  is a *Banach space*, i.e., a normed linear space which is complete.

**Definition 3.13.** The *dual* of a Banach space  $X$  is the space  $X'$  of all the continuous linear functions from  $X$  to the reals.

For a compact space  $X$ ,  $C(X) = C_b(X)$ , since all the real-valued functions over  $X$  are bounded. If  $\mu$  is a probability measure on  $X$ , we can consider  $\mu$  as an element of  $C'(X)$ , the dual of  $C(X)$  as follows: for each  $f \in C(X)$ , let  $L_\mu(f) = \int_X f(x)d\mu$ . Then  $L_\mu : C(X) \rightarrow \mathbb{R}$ .

The *weak-\** topology on  $C'(X)$  is defined as the smallest topology such that for each  $f \in C(X)$  the map  $L \mapsto L(f)$  where  $L$  ranges on  $C'(X)$  is continuous.

When restricted to the space  $\Delta X$ , the weak-*\** topology has as subbasis the sets:

$$U_{\mu_0, \epsilon, f} = \{ \mu \in \Delta X : | \int f d\mu - \int f d\mu_0 | < \epsilon \}$$

If  $X$  is a Polish space, or a Compact Hausdorff space, this is actually the same topology as the one generated by the sets  $\beta^p(E)$  for Borel sets  $E$  and  $p \in [0, 1]$  (see [Bla51]). In general, though, the weak-*\** topology is weaker.

If a topological space  $X$  is Hausdorff, then  $\Delta_r(X)$ , endowed with the weak-*\** topology is also a Hausdorff space. And if  $X$  is compact, so is  $\Delta_r(X)$  (Alaoglu's Theorem. See, e.g., [Hei93], [Kec95]). Furthermore, if  $X$  is metric, every probability measure on  $X$  is regular, so  $\Delta_r X = \Delta X$  (Ulam's theorem, see [Dud89]).

**Theorem 3.4.** (see [Hei93], also [Mèt63], Theorem III.3.2) *Let  $X_n$  be a sequence of Hausdorff topological spaces and  $f_n : X_{n+1} \rightarrow X_n$  a surjective continuous map for  $n \geq 0$ . If  $\mu_n$  is a regular Borel probability measure on  $X_n$  such that  $\mu_{n+1}f_n^{-1} = \mu_n$  for all  $n \geq 0$ , then there is a unique regular Borel probability measure  $\mu$  on the projective limit of the  $X_n$  such that for all  $n \geq 0$ ,  $\mu\pi_n^{-1} = \mu_n$ .*

**Corollary 3.2.** *The functor  $\Delta_r$  is  $\omega^{op}$ -continuous in the category **Haus** of Hausdorff topological spaces and continuous functions between them.*

The theorems 3.3 and 3.4 have been exploited to obtain some of the results mentioned in Chapter

7, and can be easily extended to yield final coalgebras for the class of measure polynomial functors defined in Chapter 4, via Theorem 2.5.

### 3.5 A Lemma on generators of $\sigma$ -algebras

**Definition 3.14.** A  $\pi$ -system is a family of subsets of a set  $M$  which is closed under finite intersections. A  $\lambda$ -system is a family including  $M$  and closed under complements and countable intersections.

**Theorem 3.5.** (*Dynkin's  $\pi$ - $\lambda$  Theorem, [Dyn61]*) *If  $\mathcal{A}$  is a  $\pi$ -system and  $\mathcal{L}$  is a  $\lambda$ -system, then  $\mathcal{A} \subseteq \mathcal{L}$  implies  $\sigma(\mathcal{A}) \subseteq \mathcal{L}$ .*

The following Lemma about probability measures is a consequence of Dynkin's  $\pi$ - $\lambda$  Theorem and will be used later.

**Lemma 3.4.** [*Dyn61*] *Suppose that  $\mu_1, \mu_2$  are probability measures on  $\sigma(\mathcal{F})$ , where  $\mathcal{F}$  is a  $\pi$ -system. If  $\mu_1$  and  $\mu_2$  agree on  $\mathcal{F}$ , then they agree on  $\sigma(\mathcal{F})$ .*

**Lemma 3.5.** (*Heifetz and Samet, [HS98]*) *Let  $\mathcal{F}$  be a boolean algebra of sets that generates the  $\sigma$ -algebra  $\Sigma$  on a measurable space  $(M, \Sigma)$ . Then the  $\sigma$ -algebra  $\mathcal{F}_\Delta$  generated by the family of sets*

$$\{\beta^p(E) : E \in \mathcal{F} \text{ and } p \in [0, 1]\}$$

*is the same as  $\Sigma_\Delta$ , the  $\sigma$ -algebra generated by the sets  $\beta^p(E)$  with  $E \in \Sigma$ .*

We have strengthened this Lemma in the following way:

**Lemma 3.6.** *Let  $\mathcal{F}$  be a  $\pi$ -system of sets that generates the  $\sigma$ -algebra  $\Sigma$  on a measurable space  $M$ . Then the  $\sigma$ -algebra  $\mathcal{F}_\Delta$  is the same as  $\Sigma_\Delta$ .*

To prove it, we first need some facts about real numbers.

**Definition 3.15.** Given a real number  $x$ , the *floor* of  $x$  is the greatest integer that is less or equal than  $x$ . We denote it with  $\lfloor x \rfloor$ .

It follows from the definition that

$$\lfloor x \rfloor - 1 = \lfloor x - 1 \rfloor \leq x - 1 < \lfloor x \rfloor \leq x < \lfloor x \rfloor + 1$$

Given  $p \in [0, 1]$  and  $n \in \mathbb{N}$ ,  $k = \lfloor np \rfloor$  is such that

$$\frac{k}{n} \leq p < \frac{k+1}{n}$$

Furthermore, from

$$\frac{np-1}{n} < \frac{\lfloor np \rfloor}{n} \leq \frac{np}{n}$$

it follows that

$$\lim_{n \rightarrow \infty} \frac{\lfloor np \rfloor}{n} = p$$

The following easy facts will also be used in the proof of Lemma 3.6:

- $p \leq q$  iff for all  $n \geq 1$ ,  $p < q + \frac{1}{n}$ .
- If for all  $n, a_n \geq 0$  then  $\sum_{n=1}^{\infty} a_n \geq p$  iff for all  $m \geq 1$ ,  $\exists k(\sum_{n=1}^k a_n \geq p - \frac{1}{m})$ .

*Proof. (of Lemma 3.6) :* Let  $\mathcal{F}' = \bigcap_p (\beta^p)^{-1}(\mathcal{F}_\Delta)$ . That is,

$$\mathcal{F}' = \{F \in \Sigma : \forall p \in [0, 1](\beta^p(F) \in \mathcal{F}_\Delta)\}.$$

So for all  $p$ ,  $\beta^p(\mathcal{F}') \subseteq \mathcal{F}_\Delta$ . Clearly  $\mathcal{F} \subseteq \mathcal{F}'$ . We show below that  $\mathcal{F}'$  is a  $\lambda$ -system. Therefore by Dynkin's Theorem, we have  $\Sigma = \sigma(\mathcal{F}) \subseteq \mathcal{F}'$ . It follows that for all  $p$ ,  $\beta^p(\Sigma) \subset \mathcal{F}_\Delta$ . So  $\Sigma_\Delta = \sigma(\bigcup_p \beta^p(\Sigma)) \subset \mathcal{F}_\Delta$ , as desired.

- Since  $\beta^p(M) = \Delta M \in \mathcal{F}_\Delta$  for all  $p \in [0, 1]$ ,  $M$  is in  $\mathcal{F}'$ .
- If we assume that  $E \in \mathcal{F}'$ , then for all  $q \in [0, 1]$ ,  $\beta^q(E) \in \mathcal{F}_\Delta$ . In particular, for a fixed  $p \in [0, 1]$  and all  $n \geq \frac{1}{p}$ ,  $\beta^{(1-p)+\frac{1}{n}}(E) \in \mathcal{F}_\Delta$ . But then  $\beta^p(E^c) = \bigcap_n (\beta^{(1-p)+\frac{1}{n}}(E))^c \in \mathcal{F}_\Delta$  so  $E^c \in \mathcal{F}'$ .

With some more detail:

$$\begin{aligned}
 \beta^p(E^c) &= \{\mu \in \Delta X : \mu(E^c) \geq p\} \\
 &= \{\mu \in \Delta X : 1 - \mu(E) \geq p\} \\
 &= \{\mu \in \Delta X : \mu(E) \leq 1 - p\} \\
 &= \{\mu \in \Delta X : \forall n \geq \frac{1}{p} \mu(E) < (1 - p) + \frac{1}{n}\} \\
 &= \bigcap_{n \geq \frac{1}{p}} (\beta^{1-p+\frac{1}{n}}(E))^c
 \end{aligned}$$

- Now, if  $\{E_n\}$  is a sequence of disjoint sets from  $\mathcal{F}'$ , then we need to prove that  $\bigcup_n E_n \in \mathcal{F}'$ .

First we prove that if  $A, B$  are disjoint sets from  $\mathcal{F}'$ , then  $A \cup B \in \mathcal{F}'$ .

Claim: Let  $I_n = \{l, m\} : l, m \in \mathbb{N}, l, m \leq n$  and  $l + m \geq \lfloor np \rfloor - 2$ . Then

$$\beta^p(A \cup B) = \bigcap_n \bigcup_{I_n} (\beta^{\frac{l}{n}}(A) \cap \beta^{\frac{m}{n}}(B)).$$

Assume that  $\mu \in \beta^p(A \cup B)$ . Then  $\mu(A \cup B) = \mu(A) + \mu(B) \geq p$ . Let  $\mu(A) = a; \mu(B) = b$  and  $r = a + b$ . Fix a natural number  $n$ . Let  $k_a = \lfloor na \rfloor \leq n, k_b = \lfloor nb \rfloor \leq n$ . Then we have that  $k_a + k_b > na - 1 + nb - 1 = nr - 2 \geq \lfloor nr - 2 \rfloor = \lfloor nr \rfloor - 2 \geq \lfloor np \rfloor - 2$ , so  $\langle k_a, k_b \rangle \in I_n$ . Since  $k_a/n \leq a$ ,  $\mu \in \beta^{k_a/n}(A)$ , and in a similar fashion,  $\mu \in \beta^{k_b/n}(B)$ . This proves that

$\mu \in \beta^{\frac{l}{n}}(A) \cap \beta^{\frac{m}{n}}(B)$  for some  $\langle l, m \rangle \in I_n$ .

Now assume that for all  $n$ ,  $\mu \in \bigcup_{I_n} (\beta^{\frac{l}{n}}(A) \cap \beta^{\frac{m}{n}}(B))$ . Then for all  $n$  and some  $l, m$ ,  $\mu(A \cup B) = \mu(A) + \mu(B) \geq \frac{l+m}{n} \geq \frac{\lfloor np \rfloor - 2}{n}$ . Since this last one converges to  $p$  when  $n \rightarrow \infty$ , we conclude  $\mu(A \cup B) \geq p$ , i.e.,  $\mu \in \beta^p(A \cup B)$ .

Since  $A, B \in \mathcal{F}'$ , then for all  $m, l, n$ ,  $\beta^{\frac{l}{n}}(A) \in \mathcal{F}_\Delta$  and  $\beta^{\frac{m}{n}}(B) \in \mathcal{F}_\Delta$ , and so are the countable intersections and unions. Therefore then  $A \cup B \in \mathcal{F}'$ .

It follows that any finite union of sets in  $\mathcal{F}'$  is also in  $\mathcal{F}'$ , and this is what we use below.

Now consider the sequence of disjoint sets  $\{E_n\}$ . We claim that

$$\beta^p\left(\bigcup_n E_n\right) = \bigcap_m \bigcup_n \beta^{p-\frac{1}{m}}\left(\bigcup_{k=1}^n E_k\right)$$

Consider the following equivalent statements:

$$\begin{aligned} \mu &\in \beta^p\left(\bigcup_n E_n\right) \\ \mu\left(\bigcup_n E_n\right) &= \sum_n \mu(E_n) \geq p \\ \forall m \geq \frac{1}{p} \exists n_m \mu\left(\bigcup_{k=1}^{n_m} E_k\right) &\geq p - \frac{1}{m} \\ \forall m \geq \frac{1}{p} \exists n_m \mu &\in \beta^{p-\frac{1}{m}}\left(\bigcup_{k=1}^{n_m} E_k\right) \\ \forall m \geq \frac{1}{p} \mu &\in \bigcup_n \beta^{p-\frac{1}{m}}\left(\bigcup_{k=1}^n E_k\right) \\ \mu &\in \bigcap_{m \geq \frac{1}{p}} \bigcup_n \beta^{p-\frac{1}{m}}\left(\bigcup_{k=1}^n E_k\right) \end{aligned}$$

□

This Lemma will be important in the proof of Lemma 4.6 in the next chapter.



---

## Modal Languages and Satisfied Theories

### 4.1 Modal Languages for Measure Polynomial Functors

**Definition 4.1.** The class of *measure polynomial functors* is the smallest class of functors on  $\text{Meas}$  containing the identity  $Id$ , the constant functor  $M$  for each measurable space  $M$ , and closed under binary products, coproducts and applications of the functor  $\Delta$ .

The main goal of this chapter is to provide a construction of final coalgebras for all the measure polynomial functors in the category  $\text{Meas}$ . The method of work is related to the construction of canonical models in modal logic, but also presents significant differences. We begin the construction by defining the *ingredients* of such functors, which will be used to index the formulas in a modal language  $\mathcal{L}(T)$ .

**Definition 4.2.** Given a measure polynomial functor  $T$  we define the set of *ingredients of  $T$* ,  $\text{Ing}(T)$  as follows:  $\text{Ing}(Id) = \{Id\}$ ;  $\text{Ing}M = \{M, Id\}$ ;  $\text{Ing}(U \times V) = \{U \times V\} \cup \text{Ing}(U) \cup \text{Ing}(V)$ ;  $\text{Ing}(U + V) = \{U + V\} \cup \text{Ing}(U) \cup \text{Ing}(V)$  and  $\text{Ing}(\Delta S) = \{\Delta S\} \cup \text{Ing}(S)$ . All the ingredients of  $T$  are measure polynomial functors, and for every measure polynomial functor  $T$ ,  $\text{Ing}(T)$  is a finite set that includes  $T$ .

The ingredients of a measure polynomial functor are defined in terms of the syntax with which the polynomial is presented.

**Example 4.1.** Let  $T = Id + \Delta(Id \times M)$  for some fixed measurable space  $M$ . Then

$$\text{Ing}(T) = \{Id, M, Id \times M, \Delta(Id \times M), T\}$$

**Definition 4.3.** For each measure polynomial functor  $T$ , we are going to define a language  $\mathcal{L}(T)$ . Each formula in  $\mathcal{L}(T)$  is associated to one of the ingredients  $S$  of  $T$ . We denote this by  $\varphi : S$ , where  $S$  is an ingredient of  $T$ . Then the formulas are formed as follows:

- For each  $S \in \text{Ing}(T)$ ,  $\text{true}_S : S$  is a formula.
- If  $M$  is a measurable space which is an ingredient of  $T$  and  $A$  is a measurable subset of  $M$ , or  $A = \{m\}$  for some  $m \in M$ , then  $A : S$  is a formula. We call these formulas *formulas of constant sort*.
- If  $\varphi : S$  and  $\psi : S$  are formulas, then  $\varphi \wedge \psi : S$  is also a formula.
- If  $U \times V \in \text{Ing}(T)$ , and  $\varphi : U$  and  $\psi : V$  are formulas, then  $\langle \varphi, \psi \rangle_{U \times V} : U \times V$  is also a formula.
- If  $U + V \in \text{Ing}(T)$  and  $\varphi : U$  is a formula, then  $\text{inl}_{U+V} \varphi : U + V$  is also a formula.
- If  $U + V \in \text{Ing}(T)$  and  $\varphi : V$  is a formula, then  $\text{inr}_{U+V} \varphi : U + V$  is also a formula.
- If  $\Delta S \in \text{Ing}(T)$ ,  $p \in [0, 1]$  and  $\varphi : S$  is a formula such that all its subformulas of constant sort are measurable sets in the appropriate spaces, then  $\beta^p \varphi : \Delta S$  is a formula.
- If  $\varphi : T$  is a formula, then  $[\text{next}] \varphi : Id$  is a formula.

We will indicate with  $\varphi :: S$  that  $\varphi$  is a formula of sort  $S$  such that all its subformulas of constant sort are measurable sets. Note that  $\varphi :: S$  implies  $\varphi : S$ .

**Example 4.2.** Let  $T = Id + \Delta(Id \times M)$  as in the Example 4.1. Here are some examples of formulas and their sorts:

$$\begin{array}{lll}
\text{true}_{Id} : Id & A : M & \langle \text{true}_{Id}, A \rangle : Id \times M \\
\beta^p \langle \text{true}_{Id}, A \rangle : \Delta(Id \times M) & \text{inl true}_{Id} : T & \text{inr } \beta^p \langle \text{true}_{Id}, A \rangle : T \\
[\text{next}] \text{inl true}_{Id} : Id & [\text{next}] \text{inr } \beta^p \langle \text{true}_{Id}, A \rangle : Id & 
\end{array}$$

Languages like  $\mathcal{L}(T)$  defined here appeared in papers by Rößiger, [Röß99, Röß01], Kurz, [Kur01], and Jacobs [Jac01] on coalgebraic generalizations of modal logic. Those languages had semantics in coalgebras in the category  $\text{Set}$ . For our work on  $\text{Meas}$  here, the precedent is the paper by Heifetz and Samet [HS98], where they develop a language for the specific case of type spaces, which we will analyze in Chapter 7. It should also be compared to the language introduced by Larsen and Skou in [LS91] to test for bisimulation in probabilistic systems.

**Definition 4.4.** Given a measure polynomial functor  $T$  and a  $T$ -coalgebra  $(X, c)$ , the formulas of the language  $\mathcal{L}(T)$  of each sort  $S$  will be interpreted in  $S(X)$  as follows:

$$\begin{array}{ll}
\llbracket \text{true}_S \rrbracket_S^c = SX & \llbracket \text{inl } \varphi \rrbracket_{U+V}^c = \text{inl} (\llbracket \varphi \rrbracket_U^c) \\
\llbracket A \rrbracket_M^c = A & \llbracket \text{inr } \varphi \rrbracket_{U+V}^c = \text{inr} (\llbracket \varphi \rrbracket_V^c) \\
\llbracket \varphi \wedge \psi \rrbracket_S^c = \llbracket \varphi \rrbracket_S^c \cap \llbracket \psi \rrbracket_S^c & \llbracket \beta^p \varphi \rrbracket_{\Delta S}^c = \beta^p (\llbracket \varphi \rrbracket_S^c) \\
\llbracket \langle \varphi, \psi \rangle \rrbracket_{U \times V}^c = \llbracket \varphi \rrbracket_U^c \times \llbracket \psi \rrbracket_V^c & \llbracket [\text{next}] \varphi \rrbracket_{Id}^c = c^{-1} (\llbracket \varphi \rrbracket_T^c)
\end{array}$$

Here  $A$  is any measurable subset or singleton of  $M$ , a constant functor which is an ingredient of  $T$ .

We say that a point  $x \in SX$  *satisfies* the formula  $\varphi : S$  if  $x \in \llbracket \varphi \rrbracket_S^c$ . Notice that if  $\varphi :: S$ , then  $\llbracket \varphi \rrbracket_S$  is measurable.

**Example 4.3.** Let's make Example 4.2 where  $T = Id + \Delta(Id \times M)$  somewhat more concrete by

letting  $M$  be the interval  $[0, 1]$  of the reals, and  $A = (1/2, 2/3]$ . Let  $X = \{x, y, z\}$  and let

$$c(x) = \text{inl } (y)$$

$$c(y) = \text{inr } (\delta_z \times \mu)$$

$$c(z) = \text{inr } \left( \left( \frac{1}{3} \delta_x \times \lambda \right) + \left( \frac{2}{3} \delta_y \times \lambda \right) \right)$$

where  $\lambda$  is the Lebesgue measure over  $[0, 1]$ .

$$\llbracket \text{true}_{Id} \rrbracket_{Id}^c = X$$

$$\llbracket (1/2, 2/3] \rrbracket_{[0,1]}^c = (1/2, 2/3]$$

$$\llbracket \langle \text{true}_{Id}, (1/2, 2/3] \rangle \rrbracket_{Id \times [0,1]}^c = X \times (1/2, 2/3]$$

$$\llbracket \beta^{1/6} \langle \text{true}_{Id}, (1/2, 2/3] \rangle \rrbracket_{\Delta(Id \times [0,1])}^c = \{ \mu \in \Delta(X \times [0, 1]) : \mu(X \times (1/2, 2/3]) \geq 1/6 \}$$

$$\llbracket \text{inl } \text{true}_{Id} \rrbracket_T^c = \text{inl } X$$

$$\llbracket \text{inr } \beta^{1/6} \langle \text{true}_{Id}, (1/2, 2/3] \rangle \rrbracket_T^c = \text{inr } \{ \mu \in \Delta(X \times [0, 1]) : \mu(X \times (1/2, 2/3]) \geq 1/6 \}$$

$$\llbracket \llbracket \text{next} \rrbracket \text{inl } \text{true}_{Id} \rrbracket_{Id}^c = \{x\}$$

$$\llbracket \llbracket \text{next} \rrbracket \text{inr } \beta^{1/6} \langle \text{true}_{Id}, (1/2, 2/3] \rangle \rrbracket_{Id}^c = \{y, z\}$$

In contrast,

$$\llbracket \llbracket \text{next} \rrbracket \text{inr } \beta^{1/2} \langle \text{true}_{Id}, (1/2, 2/3] \rangle \rrbracket_{Id}^c = \emptyset$$

**Lemma 4.1.** *Coalgebra morphisms reflect the semantics. In other words, if  $f : (X, c) \rightarrow (Y, b)$  is a*

$T$ -coalgebra morphism, then for every  $\varphi \in \mathcal{L}(T)$ , if  $\varphi : S$  then  $(Sf)^{-1}(\llbracket \varphi \rrbracket_S^b) = \llbracket \varphi \rrbracket_S^c$ .

*Proof.* We prove this by induction on the formulas of  $\mathcal{L}(T)$ . If  $\varphi$  is  $\text{true}_S$ , then  $(Sf)^{-1}(\llbracket \varphi \rrbracket_S^b) = (Sf)^{-1}SY = SX = \llbracket \varphi \rrbracket_S^c$ . Similarly, for  $A : M$  for some constant sort  $M$ ,  $(Mf)^{-1}(\llbracket A \rrbracket_S^b) = (1_M)^{-1}A = A = \llbracket A \rrbracket_S^c$ .

Since inverse images preserve intersections, the Lemma is trivial for the case of conjunctions. For products, we have to use the inductive hypothesis:

$$\begin{aligned}
 ((U \times V)f)^{-1}(\llbracket \langle \varphi, \psi \rangle \rrbracket_{U \times V}^b) &= (Uf \times Vf)^{-1}(\llbracket \varphi \rrbracket_U^b \times \llbracket \psi \rrbracket_V^b) \\
 &= (Uf)^{-1}(\llbracket \varphi \rrbracket_U^b) \times (Vf)^{-1}(\llbracket \psi \rrbracket_V^b) \\
 &= \llbracket \varphi \rrbracket_U^c \times \llbracket \psi \rrbracket_V^c \\
 &= \llbracket \langle \varphi, \psi \rangle \rrbracket_{U \times V}^c
 \end{aligned}$$

For sums, we have:

$$\begin{aligned}
 ((U + V)f)^{-1}(\llbracket \text{inl } \varphi \rrbracket_{U+V}^b) &= (Uf + Vf)^{-1}(\text{inl } \llbracket \varphi \rrbracket_U^b) \\
 &= \text{inl } (Uf)^{-1}(\llbracket \varphi \rrbracket_U^b) \\
 &= \text{inl } \llbracket \varphi \rrbracket_U^c \\
 &= \llbracket \text{inl } \varphi \rrbracket_{U+V}^c
 \end{aligned}$$

and similarly for  $V$  and the inclusion to the right  $\text{inr}$ .

For formulas of sort  $\Delta S$  and of the form  $\beta^p \varphi$ , we also use Lemma 3.1

$$\begin{aligned}
 (\Delta S f)^{-1}(\llbracket \beta^p \varphi \rrbracket_{\Delta S}^c) &= (\Delta S f)^{-1}(\beta^p \llbracket \varphi \rrbracket_S^c) \\
 &= \beta^p((S f)^{-1}(\llbracket \varphi \rrbracket_S^c)) \\
 &= \beta^p \llbracket \varphi \rrbracket_S^b \\
 &= \llbracket \beta^p \varphi \rrbracket_{\Delta S}^b
 \end{aligned}$$

Finally, for formulas of the form  $[\text{next}] \varphi$  with  $\varphi : T$ ,

$$\begin{aligned}
 f^{-1}(\llbracket [\text{next}] \varphi \rrbracket_{Id}^b) &= f^{-1}(b^{-1}(\llbracket \varphi \rrbracket_T^c)) \\
 &= c^{-1}((T f)^{-1}(\llbracket \varphi \rrbracket_T^c)) \\
 &= c^{-1}(\llbracket \varphi \rrbracket_T^c) \\
 &= \llbracket [\text{next}] \varphi \rrbracket_{Id}^c
 \end{aligned}$$

Here we used that  $b \circ f = T f \circ c$ , because  $f$  is a  $T$ -coalgebra morphism.  $\square$

## 4.2 The Space of Satisfied Theories

We will use the formulas in  $\mathcal{L}(T)$  to describe the elements of coalgebras and other sets  $SX$ . We do this by defining for each  $T$ -coalgebra  $(X, c)$  and each  $x \in SX$  where  $S \in \text{Ing}(T)$ ,

$$d_S^c(x) = \{\varphi : S \mid x \in \llbracket \varphi \rrbracket_S^c\} \quad (4.1)$$

The set  $d_S^c(x)$  will be called the *theory satisfied* by point  $x \in SX$ . We call the maps  $d_S^c$  the *description maps*. As a consequence of Lemma 4.1, we have:

**Corollary 4.1.** *If  $f : (X, c) \rightarrow (Y, b)$  is a morphism of coalgebras, then for every  $S \in \text{Ing}(T)$ ,*

$d_S^b \circ Sf = d_S^c$ . That is, coalgebra morphisms preserve description maps.

**Definition 4.5.** For each of the ingredients  $S$  of a polynomial functor  $T$ , we define the set  $S^*$  as follows:

$$S^* = \{d_S^c(x) \mid \text{for some coalgebra } (X, c), x \in SX\}. \quad (4.2)$$

Notice that  $S^*$  is a collection of subsets of  $\mathcal{L}(T)$ , and therefore a set. For each formula  $\varphi$  of sort  $S$ , we let

$$|\varphi|_S = \{s \in S^* \mid \varphi \in s\}. \quad (4.3)$$

Now we can endow  $S^*$  with the  $\sigma$ -algebra generated by the sets  $|\varphi|_S$  for all  $\varphi :: S$ . Since  $|\varphi| \cap |\psi| = |\varphi \wedge \psi|$ , we have in the sets  $|\varphi|$  a  $\pi$ -system of generators of the  $\sigma$ -algebra on  $S^*$ .

Now that we defined  $S^*$  as a measurable space, we can ask the question if the function  $d_S^c : SX \rightarrow S^*$  is measurable.

**Lemma 4.2.** For all  $(X, c)$  and all  $S \in \text{Ing}(T)$ :

1. For all  $\varphi : S$ ,  $\llbracket \varphi \rrbracket_S^c = (d_S^c)^{-1}(|\varphi|)$ .
2.  $d_S^c : SX \rightarrow S^*$  is measurable.

*Proof.* The first part follows immediately from the definition, while the second follows from the first. Recall that to prove that a function is measurable, it is enough to prove that for a family of generators, the inverse images under the function are measurable sets. So, it's enough to recall that for all  $\varphi :: S$ ,  $\llbracket \varphi \rrbracket_S^c$  is measurable.  $\square$

Our purpose is to prove that  $Id^*$  is the carrier of the final coalgebra for a measure polynomial functor. The space  $Id^*$  is similar to the *canonical model* often used in Modal Logic (see, for example,

[BdRV01]). The difference is that we don't use a deductive system of inference to determine the sets of formulas that make up the model, but use instead all the sets of formulas that are realised in *some* model.

The idea is that each point  $s$  in  $Id^*$  is a set of formulas and these formulas will contain the information of what  $c^*(s)$  will be, if  $c^*$  is the structure map of the final coalgebra. Each  $s \in Id^*$  is the description of some element  $x$  in a coalgebra  $(X, c)$ . The description of  $c(x)$  will be already contained in  $s$ . The following Lemma makes this formal.

**Lemma 4.3.** *There is a measurable map  $g : Id^* \rightarrow T^*$  such that for every  $T$ -coalgebra  $(X, c)$ , the following diagram commutes,*

$$\begin{array}{ccc} X & \xrightarrow{c} & TX \\ d_{Id}^c \downarrow & & \downarrow d_T^c \\ Id^* & \xrightarrow{g} & T^* \end{array} \quad (4.4)$$

and for all  $\varphi : T$ ,  $g^{-1}|\varphi|_T = |[\mathbf{next}]\varphi|_{Id}$ .

*Proof.* We define  $g$  by

$$g(s) = \{\varphi : T \mid [\mathbf{next}]\varphi \in s\}. \quad (4.5)$$

Assume that  $(X, c)$  is a coalgebra with  $x \in X$  such that  $s = d_{Id}^c(x)$ . Then  $c(x) \in TX$ , and

$$\begin{aligned} g^{-1}(d_{Id}^c(x)) &= g^{-1}(\{\varphi : Id \mid x \in \llbracket \varphi \rrbracket_{Id}\}) \\ &= \{\psi : T \mid [\mathbf{next}]\psi \in \{\varphi : Id \mid x \in \llbracket \varphi \rrbracket_{Id}\}\} \\ &= \{\psi : T \mid x \in \llbracket [\mathbf{next}]\psi \rrbracket_{Id}\} \\ &= \{\psi : T \mid c(x) \in \llbracket \psi \rrbracket_T\} \\ &= d_T^c(c(x)) \end{aligned}$$

Now take  $s \in g^{-1}(|\varphi|_T)$ . Then  $g(s) \in |\varphi|_T$ , which is equivalent to saying that  $\varphi \in g(s)$ , or that



[next] $\varphi \in s$ . This proves the second statement, and when applied to formulas  $\varphi :: T$ , it proves the measurability of  $g$ .  $\square$

### 4.3 From $S^*$ to $S(Id^*)$

Now we have the map  $g : Id^* \rightarrow T^*$  and for every coalgebra, the map  $d^c : X \rightarrow Id^*$  makes the diagram in (4.4) commute. But to get the final coalgebra, we need a map  $c^* : Id^* \rightarrow T(Id^*)$ . We'll achieve this by finding maps  $r_S$  connecting  $S^*$  with  $S(Id^*)$  for each ingredient of  $T$ . As a preliminary step, we need some measurable maps introduced in the following three Lemmas.

**Lemma 4.4.** *Let  $U \times V \in \text{Ing}(T)$ . There is a measurable map  $\langle \pi_1, \pi_2 \rangle : (U \times V)^* \rightarrow U^* \times V^*$  such that for every coalgebra  $(X, c)$ ,  $\langle \pi_1, \pi_2 \rangle \circ d_{U \times V}^c = d_U^c \times d_V^c$  that is, the following diagram commutes:*

$$\begin{array}{ccc}
 (U \times V)X & & \\
 d_{U \times V}^c \downarrow & \searrow^{d_U^c \times d_V^c} & \\
 (U \times V)^* & \xrightarrow{\langle \pi_1, \pi_2 \rangle} & U^* \times V^*
 \end{array} \tag{4.6}$$

Furthermore, if  $\varphi : U$  and  $\psi : V$ ,  $\langle \pi_1, \pi_2 \rangle^{-1}(|\varphi| \times |\psi|) = |\langle \varphi, \psi \rangle|$ .

*Proof.* We define  $\pi_1 : (U \times V)^* \rightarrow U^*$  by

$$\pi_1(s) = \{\varphi : U \mid \langle \varphi, \text{true}_V \rangle \in s\}.$$

$\pi_2 : (U \times V)^* \rightarrow V^*$  is defined in a similar way. First we check that for every  $s \in (U \times V)^*$ ,  $\pi_1(s) \in U^*$ .

We know there must exist some  $(X, c)$  such that for some  $x \in (U \times V)X$ ,  $d_{U \times V}^c(x) = s$ . But then  $x = \langle u, v \rangle$  with  $u \in UX$  and  $v \in VX$ . Now it's easy to check that  $\pi_1(s) = \{d_U^c(u) : \varphi \in \pi_1(s)\}$  iff  $\langle \varphi, \text{true}_V \rangle \in s = d_{U \times V}^c(\langle u, v \rangle)$  iff  $\langle u, v \rangle \in \llbracket \langle \varphi, \text{true}_V \rangle \rrbracket_{U \times V}^c$  iff  $u \in \llbracket \varphi \rrbracket_U^c$  iff  $\varphi \in d_U^c(u)$ . In the same

manner, one can check that  $\pi_2(s) = d_V^c(v)$  and therefore  $\pi_2(s) \in V^*$ .

To verify that  $\langle \pi_1, \pi_2 \rangle^{-1}(|\varphi| \times |\psi|) = |\langle \varphi, \psi \rangle|$ , consider the following equivalent statements:

$$\begin{aligned}
s &\in |\langle \varphi, \psi \rangle|_{U \times V} \\
\langle \varphi, \psi \rangle &\in s \\
\langle \varphi, \text{true}_V \rangle, \langle \text{true}_U, \psi \rangle &\in s \\
\varphi \in \pi_1(s) \text{ and } \psi \in \pi_2(s) \\
\pi_1(s) \in |\varphi|_U \text{ and } \pi_2(s) \in |\psi|_V \\
\langle \pi_1, \pi_2 \rangle(s) &\in |\varphi|_U \times |\psi|_V
\end{aligned}$$

The measurability of  $\langle \pi_1, \pi_2 \rangle$  follows from the equation above and the fact that the  $\sigma$ -algebra on  $U^* \times V^*$  is generated by the sets  $|\varphi|_U \times |\psi|_V$ , with  $\varphi :: U$  and  $\psi :: V$ .  $\square$

**Lemma 4.5.** *Let  $U + V \in \text{Ing}(T)$ . There is a measurable map  $\alpha : (U + V)^* \rightarrow U^* + V^*$  such that for every coalgebra  $(X, c)$ ,  $\alpha \circ d_{U+V}^c = d_U^c + d_V^c$ , and if  $\varphi : U$  and  $\psi : V$ , then  $\alpha^{-1}(\text{inl}_{U+V} |\varphi|_U) = |\text{inl } \varphi|_{U+V}$  and  $\alpha^{-1}(\text{inr}_{U+V} |\psi|_V) = |\text{inr } \psi|_{U+V}$ .*

*Proof.* We define  $\alpha : (U + V)^* \rightarrow U^* + V^*$  by

$$\alpha(s) = \begin{cases} \text{inl}_{U^*+V^*}(\{\varphi : U \mid \text{inl } \varphi \in s\}), & \text{if } \text{inl } \text{true}_U \in s \\ \text{inr}_{U^*+V^*}(\{\varphi : V \mid \text{inr } \varphi \in s\}), & \text{if } \text{inr } \text{true}_V \in s \end{cases} \quad (4.7)$$

We check that for every  $s \in (U + V)^*$ ,  $\alpha(s) \in U^* + V^*$ . We know as before that there must exist some  $(X, c)$  such that for some  $x \in (U + V)X$ ,  $d_{U+V}^c(x) = s$ . But then  $x = \text{inl } u$  with  $u \in UX$  or  $x = \text{inr } v$  with  $v \in VX$ , and only one of these is valid. Let's assume that  $x = \text{inl } u$  (the other case is handled in the same way). In this case we have that  $u \in UX = \llbracket \text{true}_U \rrbracket_U^c$  and therefore  $\text{inl } \text{true}_U \in s$ . So  $\alpha(s) = \text{inl}_{U^*+V^*}(\{\varphi : U \mid \text{inl } \varphi \in s\})$ . We claim that  $\{\varphi : U \mid \text{inl } \varphi \in s\} = d_U^c(u)$ . To verify this,

it is enough to note that  $\text{inl } \varphi \in s = d_{U+V}^c(\text{inl } u)$  iff  $\text{inl } u \in \llbracket \text{inl } \varphi \rrbracket = \text{inl } \llbracket \varphi \rrbracket$  iff  $u \in \llbracket \varphi \rrbracket$ .

Now  $(\alpha \circ d_{U+V}^c \circ \text{inl})(u) = (\alpha \circ d_{U+V}^c)(x) = \text{inl } d_U^c(u)$ , and similarly for  $v$ , which proves that  $\alpha \circ d_{U+V}^c = d_U^c + d_V^c$ ,

Finally, to prove the last assertion, and that  $\alpha$  is measurable, consider the following equivalent statements:

$$\begin{aligned} s &\in |\text{inl } \varphi| \\ \text{inl true}_U &\in s \text{ and } \text{inl } \varphi \in s \\ \text{inl true}_U &\in s \text{ and } \varphi \in \{\psi : U \mid \text{inl } \psi \in s\} \\ \text{inl true}_U &\in s \text{ and } \{\psi : U \mid \text{inl } \psi \in s\} \in |\varphi| \\ \alpha(s) &\in \text{inl}_{U^*+V^*}(|\varphi|) \end{aligned}$$

□

**Lemma 4.6.** *Let  $\Delta S \in \text{Ing}(T)$ . There is a measurable map  $\epsilon : (\Delta S)^* \rightarrow \Delta(S^*)$  such that for any coalgebra  $(X, c)$ ,  $\epsilon \circ d_{\Delta S}^c = \Delta d_{S^*}^c$ , and for all  $\varphi :: S$ ,  $\epsilon^{-1}(\beta^p(|\varphi|)) = |\beta^p \varphi|$ .*

*Proof.* For a given  $s \in (\Delta S)^*$ , fix  $(X, c)$  and  $\mu \in \Delta SX$  such that  $d_{\Delta S}^c(\mu) = s$ . We define

$$\epsilon(s) = (\Delta d_{S^*}^c)\mu. \tag{4.8}$$

This definition depends on  $(X, c)$  and  $\mu$ , but notice that we get a probability measure on  $S^*$ .

Let  $\mathcal{F}$  be the family of the sets of the form  $|\varphi|$  for  $\varphi :: S$ . For each set in  $\mathcal{F}$  we get that:

$$\begin{aligned} ((\Delta d_{S^*}^c)\mu)(|\varphi|) &= \mu((d_S^c)^{-1}(|\varphi|)) && \text{by the definition of } \Delta \text{ as a functor} \\ &= \mu(\llbracket \varphi \rrbracket_S^c) && \text{by Lemma 4.2} \\ &= \max\{p \mid \beta^p \varphi \in s\} \end{aligned}$$

The last equality deserves a little explanation. If we let  $q = \mu \llbracket \varphi \rrbracket_S^c$ , then for all  $p \leq q$ ,  $\mu \in \beta^p \varphi$ , and this proves that  $\max\{p \mid \beta^p \varphi \in s\} \leq q$ . But also  $q \in \{p \mid \beta^p \varphi \in s\}$ , so the equality follows.

Since the expression ‘ $\max\{p \mid \beta^p \varphi \in s\}$ ’ depends only on  $s$ , for the sets in  $\mathcal{F}$  we have proved that the definition is in fact independent of  $(X, c)$  and  $\mu$ . But  $\mathcal{F}$  is a  $\pi$ -system of generators of the  $\sigma$ -algebra on  $S^*$ . By Lemma 3.4, it follows that  $\epsilon(s)$  is also well defined, and it follows directly from (4.8) that  $\epsilon \circ d_{\Delta}^c S = \Delta d_S^c$ .

Finally, we need to prove that  $\epsilon$  is measurable. By Lemma 3.6, the sets  $E$  of the form  $\beta^p |\varphi|$  with  $\varphi :: S$  or finite intersections of them,  $\epsilon^{-1}(E)$  are enough to generate the  $\sigma$ -algebra of measurable subsets of  $\Delta(S^*)$ . These are easier to handle than the sets  $\beta^p(E)$  for just any measurable subset  $E$ .

So for these sets we have:

$$\begin{aligned}
 \epsilon^{-1}(\beta^p |\varphi|) &= \{s \in (\Delta S)^* \mid \epsilon(s) \in \beta^p(|\varphi|)\} \\
 &= \{s \in (\Delta S)^* \mid \epsilon(s)(|\varphi|) \geq p\} \\
 &= \{s \in (\Delta S)^* \mid \beta^p \varphi \in s\} \quad (*) \\
 &= |\beta^p \varphi|
 \end{aligned}$$

The second case (finite intersections of sets of this kind), follows easily from the fact that inverse images preserve intersections.

The line marked above with (\*) also needs to be explained. The argument is similar to one we had before in this proof. Suppose that  $\epsilon(s)(|\varphi|) = q \geq p$ . Then  $\beta^q \varphi \in s$ , but also  $\beta^p \varphi \in s$  for all  $p \leq q$ , since all theories of all points must have this monotonicity property. In the other direction, if  $\beta^p \varphi \in s$ , then the largest  $q$  such that  $\beta^q \varphi \in s$  is at least  $p$ , and  $\epsilon(s)(|\varphi|)$  is the maximum of those, so  $\epsilon(s)(|\varphi|) \geq p$ . □

Before we proceed, we need to define some natural sets  $\hat{\varphi}$  that act as interpretations of the

formulas of  $\mathcal{L}(T)$  in the sets  $S(Id^*)$ , as we'll see shortly.

**Definition 4.6.** For the special case of the functor  $Id$ , which is always an ingredient of  $T$ ,  $Id(Id^*) = Id^*$ , so we can take for all  $\varphi : Id$ ,  $\widehat{\varphi} = |\varphi|$ . For all  $S$ ,  $\widehat{\text{true}_S} = S(Id^*)$ . For  $A : M$ ,  $\widehat{A} = A$ .  $\widehat{\varphi \wedge \psi} = \widehat{\varphi} \cap \widehat{\psi}$ . For  $\langle \varphi, \psi \rangle : U \times V$ ,  $\widehat{\langle \varphi, \psi \rangle} = \widehat{\varphi} \times \widehat{\psi}$ . For  $\varphi : U$ ,  $\widehat{\text{inl } \varphi} = \text{inl } (\widehat{\varphi})$ ; and  $\widehat{\text{inr } \varphi} = \text{inr } (\widehat{\varphi})$ . For  $\varphi :: S$ ,  $\widehat{\beta^p \varphi} = \beta^p \widehat{\varphi}$ .

Recall that our goal was to establish a connection between  $T^*$  and  $T(Id^*)$ . To do this we need a slightly stronger result, that presents the connection between  $S^*$  and  $S(Id^*)$  for all the ingredients  $S$  of  $T$ .

**Lemma 4.7.** *There is a family of measurable maps  $r_S : S^* \rightarrow S(Id^*)$  indexed by the ingredients of  $T$  such that for all coalgebras  $(X, c)$ , the diagram below commutes:*

$$\begin{array}{ccc}
 SX & & \\
 d_S^c \downarrow & \searrow Sd_{Id}^c & \\
 S^* & \xrightarrow{r_S} & S(Id^*)
 \end{array} \tag{4.9}$$

and for all  $\varphi : S$ ,  $r_S^{-1}(\widehat{\varphi}) = |\varphi|$ .

*Proof.* For  $S = Id$ , it is enough to let  $r_{Id}$  be the identity on  $Id^*$  and then the result is immediate.

For a constant functor  $M$ , notice that when we defined  $\mathcal{L}(T)$  we included each singleton  $\{m\}$  as a formula of sort  $M$ . This way, for each  $s \in M^*$ , there is a unique  $m \in M$  such that  $\{m\} \in s$ . So we let  $r_M(s) = m$ . Then for  $m \in M = M(X)$ ,  $r_M(d_M^c(m)) = m = 1_M(m) = M(d_{Id}^c(m))$ , so (4.9) holds. For any  $A : M$ , if  $s = d_M^c(m)$  as before, then  $s \in |A|$  iff  $A \in s$  iff  $r_M(s) = m \in A = \widehat{A}$ , so  $r_M^{-1}(\widehat{A}) = |A|$ . As in all the following cases, it is trivial to check the equality holds also for conjunctions of formulas, so we won't mention it any more.

For a product functor  $U \times V$ , we use define  $r_{U \times V} = (r_U \times r_V) \circ \langle \pi_1, \pi_2 \rangle$ .

$$\begin{array}{ccc}
& UX \times VX & \\
d_{U \times V}^c \swarrow & \downarrow d_U^c \times d_V^c & \searrow (U \times V)d_{Id}^c \\
(U \times V)^* & \xrightarrow{\epsilon} U^* \times V^* & \xrightarrow{r_U \times r_V} U(Id^*) \times V(Id^*)
\end{array} \tag{4.10}$$

The triangle on the left commutes because of Lemma 4.4, and the one on the right by induction hypothesis, and this proves (4.9) for this case.

The second part is easy to check:

$$\begin{aligned}
r_{U \times V}^{-1}(\widehat{\langle \varphi, \psi \rangle}) &= \langle \pi_1, \pi_2 \rangle^{-1}((r_U \times r_V)^{-1}(\widehat{\varphi} \times \widehat{\psi})) \\
&= \langle \pi_1, \pi_2 \rangle^{-1}(|\varphi| \times |\psi|) && \text{by induction hypothesis} \\
&= |\langle \varphi, \psi \rangle| && \text{by Lemma 4.4}
\end{aligned}$$

For coproducts, we take  $r_{U+V} = (r_U + r_V) \circ \alpha$ . Using the induction hypothesis and Lemma 4.5, we get (4.9). The second part is also similar:

$$\begin{aligned}
r_{U+V}^{-1}(\widehat{\text{inl } \varphi}) &= \alpha^{-1}((r_U + r_V)^{-1}(\text{inl } \widehat{\varphi})) \\
&= \alpha^{-1} \text{inl } r_U(\widehat{\varphi}) \\
&= \alpha^{-1} \text{inl } |\varphi| && \text{by induction hypothesis} \\
&= |\text{inl } \varphi| && \text{by Lemma 4.5}
\end{aligned}$$

Similar calculations are valid for the case of  $\Delta S$ : using Lemma 4.6 one gets (4.9) and for the second part:

$$\begin{aligned}
r_{\Delta S}^{-1}(\widehat{\beta^p \varphi}) &= \epsilon^{-1}(\Delta r_S)^{-1} \beta^p(\widehat{\varphi}) \\
&= \epsilon^{-1} \beta^p(r_S)^{-1}(\widehat{\varphi}) && \text{by Lemma 3.1} \\
&= \epsilon^{-1} \beta^p(|\varphi|) && \text{by induction hypothesis} \\
&= |\beta^p \varphi| && \text{by Lemma 4.6}
\end{aligned}$$

□

## 4.4 Final Coalgebras

Now we can define the coalgebra  $(Id^*, c^*)$  by letting  $c^*$  be the composition:

$$r_T \circ g : Id^* \rightarrow T^* \rightarrow T(Id^*) \quad (4.11)$$

where  $g$  is the function from Lemma 4.3.

**Lemma 4.8.** *For each  $T$ -coalgebra  $(X, c)$ ,  $d_{Id}^c$  is a morphism of coalgebras.*

*Proof.* Consider

$$\begin{array}{ccccc}
 X & \xrightarrow{c} & TX & & \\
 d_{Id}^c \downarrow & & \downarrow d_T^c & \searrow Td_{Id}^c & \\
 Id^* & \xrightarrow{g} & T^* & \xrightarrow{r_T} & T(Id^*)
 \end{array}$$

The square commutes as we proved in Lemma 4.3, and the triangle is a special case of equation (4.9) in Lemma 4.7. □

In order to prove that  $(Id^*, c^*)$  is the final coalgebra for  $T$ , it remains to be proved that  $d_{Id}^c$  is the only morphism from  $(X, c)$  to  $(Id^*, c^*)$ . For the proof we need the following two Lemmas.

**Lemma 4.9 (Truth Lemma).** *For all  $\varphi : S$ ,  $\llbracket \varphi \rrbracket_S^{c^*} = \widehat{\varphi}$ .*

*Proof.* We prove this by induction on  $\varphi$ . The first base case is  $\text{true}_S : S$ . Here  $\llbracket \text{true} \rrbracket_S^{c^*} = S(Id^*) = \widehat{\text{true}}$ .

The other base case is  $A : M$ , where  $M$  is a constant functor in  $\text{Ing}(T)$ . Now we have that  $\llbracket A \rrbracket_S^{c^*} = A = \widehat{A}$ .

The cases for the formulas of the form  $\varphi \wedge \psi$ ,  $\langle \varphi, \psi \rangle$ ,  $\text{inl } \varphi$ ,  $\text{inr } \psi$  are trivial from the definition of  $\hat{\varphi}$ .

For the formulas  $[\text{next}]\varphi : Id$  however, a bit more of calculation is required:

$$\begin{aligned}
[[\text{next}]\varphi]_{Id}^{c^*} &= (c^*)^{-1}([\varphi]_T^{c^*}) \\
&= (c^*)^{-1}(\hat{\varphi}) && \text{by induction hypothesis} \\
&= g^{-1}(r_T^{-1}(\hat{\varphi})) && \text{by the definition of } c^* \text{ in (4.11)} \\
&= g^{-1}(|\varphi|_T) && \text{by Lemma 4.7} \\
&= [[\text{next}]\varphi]_{Id} && \text{by Lemma 4.3} \\
&= \widehat{[\text{next}]\varphi}
\end{aligned}$$

□

**Lemma 4.10.** *The description function  $d_{Id}^{c^*} : Id^* \rightarrow Id^*$  is the identity  $1_{Id^*}$*

*Proof.* Recall that for every  $\varphi : Id$ , by the previous Lemma,  $[[\varphi]_{Id}^{c^*} = \hat{\varphi} = |\varphi|$ . Then, for all  $s \in Id^*$ ,

$$d_{Id}^{c^*}(s) = \{\varphi : Id \mid s \in [[\varphi]_{Id}^{c^*}\} = \{\varphi : Id \mid s \in |\varphi|\} = s.$$

□

**Theorem 4.1.**  *$(Id^*, c^*)$  is the final  $T$ -coalgebra.*

*Proof.* Given an arbitrary  $T$ -coalgebra  $(X, c)$ , we have seen in Lemma 4.8 that  $d_{Id}^c$  is a coalgebra morphism. If  $f : (X, c) \rightarrow (Id^*, c^*)$  is another one, then by Lemma 4.1 and Lemma 4.10,  $f = d_{Id}^{c^*} \circ f = d_{Id}^c$ . □



# 5

---

## Final Coalgebras Found in Final Sequences

In this chapter we will give a different construction of the final coalgebras for measure polynomial functors. This time we will use the final sequence of a functor  $T$  instead of the language  $\mathcal{L}(T)$ . The resulting construction is of course isomorphic to the one presented in Chapter 4. To emphasize the parallel between the constructions, we re-use the names of the functions  $g, \langle \pi_1, \pi_2 \rangle, \alpha$  and  $\epsilon$ . They will now have different definitions, but they will play similar roles in the structure of the proof.

### 5.1 The Space of Realised Points in the Final Sequence

For each ingredient  $S$  of  $T$  we will consider the  $\omega^{op}$ -sequence  $(ST^n 1, ST^{n+1} 1)_{n \in \omega}$

$$S1 \xleftarrow{S!} ST1 \xleftarrow{ST!} ST^2 1 \xleftarrow{ST^2!} ST^3 1 \xleftarrow{\dots} \dots$$

Let  $P_S$  be the projective limit of this chain, and let  $\pi_S^n : P_S \rightarrow ST^n 1$  be the corresponding projections. To determine an element  $z$  of  $P_S$  it will be enough to know its components, that is,  $\pi_S^n z$  for all  $n \geq 0$ .

For the particular case of  $S = Id$  we will denote with  $\pi_n$  the projections, as we did in Example 2.6. It is worth noting that unless the functor  $S$  is  $\omega^{op}$ -continuous,  $\pi_S^n$  won't in general be equal, or even isomorphic, to  $S\pi_n$ . For each coalgebra  $(X, c)$ , recall the maps  $h_n^c : X \rightarrow T^n 1$  defined in Section 2.4 by setting  $h_0^c = !_X$  the unique function from  $X$  to  $1$  and  $h_{n+1}^c = (Th_n^c) \circ c$ . The natural source  $Sh_n^c : SX \rightarrow ST^n 1$  induces a mapping  $h_S^n^c : SX \rightarrow P_S$  such that

$$\pi_S^n h_S^n^c = Sh_n^c. \quad (5.1)$$

We will sometimes omit the superscript  $c$  when this is not a cause of confusion.

**Definition 5.1.** Let  $Z_S$  be the set of all  $z \in P_S$  such that for some coalgebra  $(X, c)$ , and some  $x \in SX$ ,  $h_S^n(x) = z$ . It follows that the functions  $h_S^n$  map  $SX$  to  $Z_S$ .

Thus  $Z_S$  is the analog of the space  $S^*$  in the previous chapter. It is the collection of all sequences in the projective limit that are realized in some coalgebra. In the particular case of  $S = Id$ , we will denote  $Z_{Id}$  just by  $Z$ , and  $\pi_{Id}^n$  by  $\pi_n$ .

The sets  $Z_S$  inherit the measurable space structure (i.e., the  $\sigma$ -algebra) from the projective limit  $P_S$ . Furthermore, this  $\sigma$ -algebra is generated by the family  $\mathcal{F}$  of sets of the form  $(\pi_S^n)^{-1}(E_n)$  with  $E_n$  measurable in  $ST^n 1$ .

**Lemma 5.1.** *The family  $\mathcal{F}$  of sets of the form  $(\pi_S^n)^{-1}(E_n)$  with  $E_n$  measurable in  $ST^n 1$  is a  $\pi$ -system.*

*Proof.* Suppose that  $E = (\pi_S^n)^{-1}(E_n)$  and  $F = (\pi_S^k)^{-1}(E_k)$ , that is,  $E$  and  $F$  are in  $\mathcal{F}$ . We can assume also that  $n \leq k$ . Then  $E \cap F = (\pi_S^n)^{-1}(E_n) \cap (\pi_S^k)^{-1}(E_k) = (\pi_S^k)^{-1}(\tau_{kn}^{-1}E_n \cap E_k)$ . Since  $\tau_{kn}$  is measurable,  $\tau_{kn}^{-1}E_n \cap E_k$  is a measurable subset of  $ST^k 1$  and  $E \cap F \in \mathcal{F}$ .  $\square$

The measurable space  $Z$  will be the carrier of the final coalgebra for  $T$ . To define the structure

map on  $Z$ , we will proceed in two stages. First, we define a map  $g : Z \rightarrow Z_T$ , and then we find a map  $r_T$  from  $Z_T$  to  $T(Z)$  that will establish the result. The first part is taken care of in the following Lemma:

**Lemma 5.2.** *There exists a measurable map  $g : Z \rightarrow Z_T$ , such that for all coalgebras  $(X, c)$  and  $x \in X$ ,  $gh^c = h_T^c c$ .*

$$\begin{array}{ccc} X & \xrightarrow{c} & TX \\ h^c \downarrow & & \downarrow h_T^c \\ Z & \xrightarrow{g} & Z_T \end{array}$$

*Proof.* We define  $g : Z \rightarrow P_T$  by  $\pi_T^n g = \pi_{n+1}$  for all  $n \geq 0$ . We need to prove that  $\pi_T^n gh = \pi_T^n h_T c$ .

$$\begin{aligned} \pi_T^n h_T c &= (Th_n)c \\ &= h_{n+1} \\ &= \pi_{n+1}h \\ &= \pi_T^n gh \quad \text{by the definition of } g \end{aligned}$$

This proves that  $gh = h_T c$  and also that  $g : Z \rightarrow Z_T$ .

Next we prove that  $g$  is measurable. Consider a measurable set  $E_n \subseteq TT^n 1$ ; since the sets of the form  $E = (\pi_T^n)^{-1}(E_n)$  generate the  $\sigma$ -algebra on  $Z_T$ , it will be enough to prove that  $g^{-1}(E)$  is measurable for all such sets  $E$ . But  $g^{-1}(E) = g^{-1}(\pi_T^n)^{-1}(E_n) = (\pi_T^n \circ g)^{-1}(E_n) = (\pi_T^{n+1})^{-1}(E_n)$ , which we know to be measurable.  $\square$

## 5.2 From $Z_S$ to $S(Z)$

To find the map  $r_T : Z_T \rightarrow T(Z)$  satisfying the appropriate conditions, we need to work making reference to the structure of  $T$ . First, we introduce some auxiliary results.

**Lemma 5.3.** *If  $(U \times V) \in \text{Ing}(T)$ , there is a measurable map  $\langle \pi_1, \pi_2 \rangle : Z_{U \times V} \rightarrow Z_U \times Z_V$  so that for every  $T$ -coalgebra  $(X, c)$ ,  $\langle \pi_1, \pi_2 \rangle h_{U \times V}^c = h_U^c \times h_V^c$  and for every  $n \geq 0$ ,  $(\pi_U^n \times \pi_V^n) \langle \pi_1, \pi_2 \rangle = \pi_{U \times V}^n$ .*

*Proof.* Let  $p_1^n : (U \times V)T^n \mathbf{1} \rightarrow UT^n \mathbf{1}$ ,  $p_2^n : (U \times V)T^n \mathbf{1} \rightarrow VT^n \mathbf{1}$ ,  $p_U : (U \times V)X \rightarrow UX$ ,  $p_V : (U \times V)X \rightarrow VX$ ,  $p_{Z_U} : Z_U \times Z_V \rightarrow Z_U$ ,  $p_{Z_V} : Z_U \times Z_V \rightarrow Z_V$  be the natural projections. The following diagram will be helpful throughout the proof:

$$\begin{array}{ccccc}
 (U \times V)X & \xrightarrow{p_U} & UX & & \\
 \downarrow h_{U \times V} & \searrow h_U \times h_V & \searrow h_U & & \\
 Z_{U \times V} & \xrightarrow{\langle \pi_1, \pi_2 \rangle} & Z_U \times Z_V & \xrightarrow{p_{Z_U}} & Z_U \\
 & \searrow \pi_{U \times V}^n & \downarrow \pi_U^n \times \pi_V^n & & \downarrow \pi_U^n \\
 & & (U \times V)T^n \mathbf{1} & \xrightarrow{p_1^n} & UT^n \mathbf{1}
 \end{array}$$

We start by defining  $\pi_1 : Z_{U \times V} \rightarrow P_U$  through  $\pi_U^n \pi_1 = p_1 \pi_{U \times V}^n$ . One needs to check that the sequence thus obtained is actually in  $P_U$ , i.e., for every  $n \geq 0$ ,  $UT^{n+1}! p_1^{n+1} \pi_{U \times V}^{n+1} = p_1^n \pi_{U \times V}^n$ . The following diagram commutes:

$$\begin{array}{ccc}
 (U \times V)T^{n+1} \mathbf{1} & \xrightarrow{p_1^{n+1}} & UT^{n+1} \mathbf{1} \\
 (U \times V)T^n \mathbf{1} \downarrow & & \downarrow UT^n ! \\
 (U \times V)T^n \mathbf{1} & \xrightarrow{p_1^n} & UT^n \mathbf{1}
 \end{array}$$

Furthermore, for any  $z \in P_{U \times V}$  (and therefore also for  $z \in Z_{U \times V}$ ) and  $n \geq 0$ , we have that  $(U \times V)T^n ! \pi^{n+1} z = \pi_{U \times V}^n z$ . So we can calculate  $UT^{n+1}! p_1^{n+1} \pi_{U \times V}^{n+1} = p_1^n (U \times V)T^n ! \pi_{U \times V}^{n+1} = p_1^n \pi_{U \times V}^n$ .

From the definition it follows that  $(\pi_U^n \times \pi_V^n)\langle \pi_1, \pi_2 \rangle = \pi_{U \times V}^n$ .

To prove that  $\langle \pi_1, \pi_2 \rangle h_{U \times V}^c = h_U \times h_V = \langle h_U p_U, h_V p_V \rangle$ , we need to show that  $\pi_1 h_{U \times V} = h_U p_U$  (and the corresponding equation for  $V$ ). This will be proved once we prove that for all  $n$ ,  $\pi_U^n \pi_1 h_{U \times V} = \pi_U^n h_U p_U$ . In fact:

$$\pi_U^n \pi_1 h_{U \times V} = p_1^n \pi_{U \times V}^n h_{U \times V} = p_1^n (U \times V) h_n = p_1^n (U h_n \times V h_n) = U h_n p_U = \pi_U^n h_U p_U.$$

The above equation also proves that  $\pi_1$  maps  $Z_{U \times V}$  to  $Z_U$ .

To prove that  $\langle \pi_1, \pi_2 \rangle$  is measurable, it's enough to prove each of the components is measurable. We do it for  $\pi_1$ . Let  $E, E_n$  be as in the proof of Lemma 5.2. Then  $\pi_1^{-1}(E) = \pi_1^{-1}(\pi_U^n)^{-1}(E_n) = (\pi_U^n \pi_1)^{-1}(E_n) = (p_1 \pi_{U \times V}^n)^{-1}(E_n)$ , which is measurable.  $\square$

**Lemma 5.4.** *If  $(U + V) \in \text{Ing}(T)$ , there is a measurable map  $\alpha : Z_{U+V} \rightarrow Z_U + Z_V$  so that for every  $T$ -coalgebra  $(X, c)$ ,  $\alpha h_{U+V}^c = h_U^c + h_V^c$ , and for all  $n \geq 0$ ,  $(\pi_U^n + \pi_V^n)\alpha = \pi_{U+V}^n$ .*

*Proof.*

$$\begin{array}{ccccc}
 (U + V)X & \xleftarrow{\text{inl}_{UX}} & UX & & \\
 \downarrow h_{U+V} & \searrow h_U + h_V & \searrow h_U & & \\
 Z_{U+V} & \xrightarrow{\alpha} & Z_U + Z_V & \xleftarrow{\text{inl}_{Z_U}} & Z_U \\
 & \searrow \pi_{U+V}^n & \downarrow \pi_U^n + \pi_V^n & & \downarrow \pi_U^n \\
 & & (U + V)T^n 1 & \xleftarrow{\text{inl}_U^n} & UT^n 1
 \end{array}$$

Since for each  $z \in Z_{U+V}$ ,  $z = h_{U+V}^c(x)$  for some  $x \in (U + V)X$  for some coalgebra  $(X, c)$ , we define  $\alpha$  by  $\alpha h_{U+V}^c(x) = (h_U^c + h_V^c)(x)$ .

For every  $n \geq 0$  we have:

$$\begin{aligned}
\pi_{U+V}^n h_{U+V} &= (U+V)h_n \\
&= Uh_n + Vh_n \\
&= \pi_U^n h_U + \pi_V^n h_V && \text{by(5.1)} \\
&= (\pi_U^n + \pi_V^n)(h_U + h_V)
\end{aligned}$$

Now we can check  $\alpha$  is well-defined. If  $(Y, d)$  is another coalgebra and  $y \in (U+V)Y$  is such that  $h_{U+V}^c(x) = h_{U+V}^d(y)$ , then for all  $n \geq 0$ ,  $(\pi_U^n + \pi_V^n)(h_U^c + h_V^c)x = (\pi_U^n + \pi_V^n)(h_U^d + h_V^d)y$ . So  $(h_U^c + h_V^c)x = (h_U^d + h_V^d)y$ , i.e.,  $\alpha h_{U+V}^c(x) = \alpha h_{U+V}^d(y)$ .

It also follows from the computation above that  $(\pi_U^n + \pi_V^n)\alpha h_{U+V} = (\pi_U^n + \pi_V^n)(h_U + h_V) = \pi_{U+V}^n h_{U+V}$ .

To prove that  $\alpha$  is measurable, consider  $E = \text{inl}_{Z_U}(\pi_U^n)^{-1}(E_n)$ , with  $E_n$  a measurable subset of  $UT^n\mathbf{1}$ . Then  $\alpha^{-1}(E) = \alpha^{-1}(\text{inl}_{Z_U}(\pi_U^n)^{-1}(E_n)) = \alpha^{-1}(\pi_U^n + \pi_V^n)^{-1}\text{inl}_U^n(E_n) = ((\pi_U^n + \pi_V^n)\alpha)^{-1}\text{inl}_U^n(E_n) = (\pi_{U+V}^n)^{-1}\text{inl}_U^n(E_n)$  is a measurable set. Here we used the fact that  $\text{inl}_{Z_U}(\pi_U^n)^{-1}(E_n) = (\pi_U^n + \pi_V^n)^{-1}\text{inl}_U^n(E_n)$ , which is easy to verify.  $\square$

**Lemma 5.5.** *Let  $\Delta S$  be an ingredient of  $T$ . Then there exists a measurable function  $\epsilon : Z_{\Delta S} \rightarrow \Delta Z_S$  so that for every  $T$ -coalgebra  $(X, c)$   $\epsilon h_{\Delta S}^c = \Delta h_S^c$  and for every  $n \geq 0$ ,  $(\Delta \pi_S^n)\epsilon = \pi_{\Delta S}^n$ .*

*Proof.* To define  $\epsilon(z)$  for a given  $z \in Z_{\Delta S}$ , we start by doing it for the family  $\mathcal{F}$  of sets of the form  $E = (\pi_S^n)^{-1}(E_n)$  with  $E_n$  measurable in  $ST^n\mathbf{1}$ .

Given  $z \in Z_{\Delta S}$ , we define

$$\epsilon(z)(E) = \pi_{\Delta S}^n(z)(E_n)$$

It's worth remarking this definition just depends on  $z$  and not on any  $\mu$  such that  $h(\mu) = z$ . To check that it does not depend on the selection of  $n$ , consider  $(X, c)$  and  $\mu \in \Delta SX$  so that  $h_{\Delta S}\mu = z$ .

$$\begin{aligned}
\epsilon(z)(E) &= \pi_{\Delta S}^n(z)(E_n) \\
&= \pi_{\Delta S}^n(h_{\Delta S}(\mu))(E_n) \\
&= (\Delta S h_n)(\mu)(E_n) && \text{by (5.1)} \\
&= \mu(S h_n)^{-1}(E_n) \\
&= \mu(h_S)^{-1}(\pi_S^n)^{-1}(E_n) && \text{by (5.1)} \\
&= \mu(h_S)^{-1}(E) \\
&= (\Delta h_S)(\mu)(E)
\end{aligned}$$

The above equation not only proves the independence of the definition from the selection of  $n$ , but also that  $\epsilon h_{\Delta S} \mu(E) = \Delta h_S \mu(E)$  for every  $E \in \mathcal{F}$  and  $\mu \in \Delta S X$ . Now we extend the definition of  $\epsilon(z)$  to every measurable subset  $F$  of  $Z_S$  by letting  $\epsilon(z)(F) = \Delta h_S \mu(F)$ . We need to check that the probability measure  $\epsilon(z)$  is well defined. If  $z = h_{\Delta S}^c(\mu) = h_{\Delta S}^d(y)$ , then we know that  $\epsilon h_{\Delta S}^c(\mu)$  and  $\epsilon h_{\Delta S}^d(y)$  agree on all the elements of the family  $\mathcal{F}$ . By Lemma 5.1,  $\mathcal{F}$  is a  $\pi$ -system, then by Lemma 3.4, the measures must agree on all measurable subsets.

To prove that  $\epsilon$  is measurable, first notice that for any measurable subset  $E_n$  of  $ST^{n-1}$ ,  $(\Delta \pi_S^n \epsilon)(z)(E_n) = \epsilon(z)(\pi_S^n)^{-1}(E_n) = \pi_{\Delta S}^n(z)(E_n)$ . By Lemma 3.5 it will be enough to prove that for a measurable subset  $E_n \subseteq ST^{n-1}$ ,  $\epsilon^{-1} \beta^p (\pi_S^n)^{-1}(E_n)$  is measurable.

$$\begin{aligned}
\epsilon(\beta^p (\pi_S^n)^{-1}(E_n)) &= \epsilon^{-1} (\Delta \pi_S^n)^{-1} \beta^p (E_n) \\
&= (\Delta \pi_S^n \epsilon)^{-1} \beta^p (E_n) \\
&= (\pi_{\Delta S}^n)^{-1} \beta^p (E_n)
\end{aligned}$$

We know the set in the last line to be measurable. We used Lemma 3.1 in the first line of the equation. □

**Lemma 5.6.** *There exists a measurable function  $r_T : Z_T \rightarrow T(Z)$  so that for every  $(X, c)$ ,  $r_T \circ h^c =$*

$Th^c$ , and for every  $n \geq 0$ ,  $T\pi_n \circ r_T = \pi_T^n$ .

*Proof.* We will prove this by induction over the ingredients of  $T$ . This is, if  $S \in \text{Ing}(T)$ , then there exists a measurable map  $r_S : Z_S \rightarrow S(Z)$  such that  $r_S h_S = S h$  and for all  $n \geq 0$ ,  $S\pi_n \circ r_S = \pi_S^n$ .

For  $S = Id$ ,  $r_{Id} = 1_Z$  is measurable and trivially satisfies the conditions.

For  $S = M$ , a constant functor, we let  $r_M = \pi_M^0 : Z_M \rightarrow M = M(1)$ . Then  $r_M h_M = \pi_M^0 \langle 1_M \rangle_{n \geq 0} = 1_M = M(h)$ , and  $M\pi_n r_M = 1_M \pi_M^0 = \pi_M^n$  for all  $n \geq 0$ .

**Products** We define  $r_{U \times V}$  as  $(r_U \times r_V) \circ \langle \pi_1, \pi_2 \rangle$ .

$$\begin{array}{ccccc}
 & & (U \times V)X & & \\
 & \swarrow h_{U \times V}^c & \downarrow h_U^c \times h_V^c & \searrow (U \times V)h_{Id}^c & \\
 Z_{U \times V} & \xrightarrow{\langle \pi_1, \pi_2 \rangle} & Z_U \times Z_V & \xrightarrow{r_U \times r_V} & (U \times V)(Z) \\
 & \searrow \pi_{U \times V}^n & \downarrow \pi_U^n \times \pi_V^n & \swarrow (U \times V)\pi_n & \\
 & & \Delta(U \times V)T^n 1 & & 
 \end{array} \tag{5.2}$$

The triangles on the left commute by Lemma 5.3, and the ones on the right by the induction hypothesis. Hence the diagram commutes.

**Coproducts** We take  $r_{U+V}$  to be  $(r_U + r_V) \circ \alpha$ . We use the diagram from (5.2), replacing  $U \times V$  with  $U + V$ , and Lemma 5.3 with Lemma 5.4.



**Probability measures** We define  $r_{\Delta S}$  as  $\Delta r_S \circ \epsilon$ .

$$\begin{array}{ccccc}
 & & \Delta SX & & \\
 & \swarrow h_{\Delta S}^c & \downarrow \Delta h_S^c & \searrow \Delta S h_{Id}^c & \\
 Z_{\Delta S} & \xrightarrow{\epsilon} & \Delta(Z_S) & \xrightarrow{\Delta r_S} & \Delta S(Z) \\
 & \searrow \pi_{\Delta S} & \downarrow \Delta \pi_S^n & \swarrow \Delta S \pi_n & \\
 & & \Delta ST^{n1} & & 
 \end{array} \tag{5.3}$$

The triangles on the left commute by Lemma 5.5, and the ones on the right by the induction hypothesis. Hence the diagram commutes.  $\square$

### 5.3 Final Coalgebras

Now we are ready to define  $\gamma : Z \rightarrow T(Z)$  as

$$r_T \circ g : Z \rightarrow Z_T \rightarrow T(Z) \tag{5.4}$$

We shall show that  $(Z, \gamma)$  is a final  $T$ -coalgebra.

**Lemma 5.7.** *For each coalgebra  $(X, c)$ ,  $h^c$  is a morphism of coalgebras.*

*Proof.* Consider the diagram:

$$\begin{array}{ccccc}
 X & \xrightarrow{c} & TX & & \\
 h^c \downarrow & & \downarrow h_T^c & \searrow Th^c & \\
 Z & \xrightarrow{g} & Z_T & \xrightarrow{r_T} & T(Z)
 \end{array}$$

The square commutes by Lemma 5.2, and the triangle by Lemma 5.6.  $\square$

**Lemma 5.8.**  $h^\gamma = 1_Z$ .

*Proof.* It will be enough to prove that for each  $n \geq 0$ ,  $\pi_n h^\gamma = \pi_n$ . For  $n = 0$ , we have that  $\pi_0 h^\gamma = h_0 = \pi_0 1_Z$ . Now assume that  $h_n^\gamma = \pi_n h^\gamma = \pi_n$ . Then,

$$\begin{aligned}
\pi_{n+1} h^\gamma &= h_{n+1}^\gamma \\
&= Th_n^\gamma \circ \gamma \\
&= T\pi_n \circ \gamma && \text{by inductive hypothesis} \\
&= T\pi_n \circ r_T \circ g && \text{by the definition of } \gamma \\
&= \pi_T^n \circ g && \text{by Lemma 5.6} \\
&= \pi_{n+1} && \text{by the definition of } g
\end{aligned}$$

□

**Theorem 5.1.**  $(Z, \gamma)$  is a final coalgebra of  $T$ .

*Proof.* Let  $(X, c)$  be a  $T$ -coalgebra. By Lemma 5.7,  $h^c$  is a coalgebra morphism. For the uniqueness, suppose that  $f$  is any morphism.

By Lemma 2.2,  $h^\gamma \circ f = h^c$ . But by Lemma 5.8,  $h^\gamma = 1_Z$ , so  $f = h^\gamma \circ f = h^c$ .

□

**Example 5.1.** Returning to the functor  $T$  of example 4.1, now we have that the structure map  $\gamma$  for  $Z$  is

$$\begin{array}{ccccc}
Z & \xrightarrow{g} & Z_T & \xrightarrow{\alpha} & Z + Z_{\Delta(Id \times M)} & \xrightarrow{1_Z + \epsilon} & Z + \Delta Z_{Id \times M} \\
\downarrow \gamma & & & & & & \downarrow 1_Z + \Delta \langle \pi_1, \pi_2 \rangle \\
Z + \Delta(Z \times M) & \xleftarrow{1_Z + \Delta(1_Z \times \pi_M^0)} & & & & & Z + \Delta(Z + Z_M)
\end{array}$$

$$\gamma = (1_Z + \Delta((1_Z \times \pi_M^0)\langle \pi_1, \pi_2 \rangle))\epsilon) \alpha g$$

Here we see how we use the different functions we defined in this Chapter to build the map  $\gamma$  for this specific functor  $T$ . Since we have given the definition of each of the intermediate maps, we see here that this method yields some more information on the structure map of the final coalgebra of a measure polynomial functor.

## 6

---

# Probabilistic Kripke Polynomial Functors in Set

The work from the previous Chapters can also be carried out in **Set**, for probabilistic Kripke polynomial functors as introduced in [MV04]. These are functors built from the identity functor, constant functors for fixed sets, the finite, covariant *finite* power set functor denoted by  $\mathcal{P}$ , functions from a fixed set  $E$ , denoted by  $\cdot^E$  (see Example 2.1), and the discrete measure functor  $\mathcal{D}$ , that assigns to a set  $X$  the set of all functions  $\mu : A \rightarrow [0, 1]$  with finite support and such that  $\sum_{a \in A} \mu(a) = 1$  (Example 2.3).

The set  $\text{Ink}(T)$  is defined similarly as done for measure polynomial functors, omitting the clause for  $\Delta$  and adding  $\text{Ink}(\mathcal{P}S) = \{\mathcal{P}S\} \cup \text{Ink}(S)$ ;  $\text{Ink}(S^E) = \{S^E\} \cup \text{Ink}(S)$ ;  $\text{Ink}(\mathcal{D}S) = \{\mathcal{D}S\} \cup \text{Ink}(S)$ .

In the paper [MV05], a construction of final coalgebras for probabilistic Kripke polynomial functors in **Set** using the method from Chapter 4 is given. In this Chapter we will do the same but using the final sequence as in Chapter 5.

Recall that in **Set**, the projective limit of an  $\omega^{op}$ -sequence  $(D_n, f_n)_{n \in \omega}$  is constructed by taking

the set

$$P = \{(d_n)_n \in \prod_{n \geq 0} D_n : f_n(d_{n+1}) = d_n\}$$

and the projections  $\pi_n : P \rightarrow D_n$ . We will use the following fact about finite subsets of such limit  $P$ :

**Lemma 6.1.** *Suppose  $x \in \mathcal{P}P$  and  $y \in P$  satisfies  $\pi_n(y) \in (\mathcal{P}\pi_n)(x)$  for all  $n \geq 0$ . Then  $y \in x$ .*

*Proof.* Let's write  $x = \{y_1, \dots, y_r\}$ , and assume the claim is not true. Then, for every  $i, 1 \leq i \leq r, y \neq y_i$  so

$$\exists n_i \pi_{n_i}(y) \neq \pi_{n_i}(y_i). \quad (6.1)$$

Let  $K = \max\{n_1, \dots, n_r\}$ . By hypothesis,  $\pi_K(y) \in (\mathcal{P}\pi_K)(x)$ , so for some  $j, 1 \leq j \leq r$ ,

$$\pi_K(y) = \pi_K(y_j).$$

Now let  $\tau_{mn} : D_m \rightarrow D_n$  be the composition  $f_n \circ f_{n+1} \circ \dots \circ f_{m+1}$  and therefore for all  $n \leq m, \tau_{mn}\pi_m = \pi_n$ . We have that

$$\pi_{n_j}(y) = \tau_{Kn_j}\pi_K(y) = \tau_{Kn_j}\pi_K(y_j) = \pi_{n_j}(y_j),$$

a contradiction with (6.1). □

Following the structure of the proof of Theorem 5.1, three new connecting maps are required, and we present them in the following three Lemmas.

**Lemma 6.2.** *Let  $S^E \in \text{Ing}(T)$ . There is a function  $\eta : Z_{S^E} \rightarrow (Z_S)^E$  such that*

1. For all  $T$ -coalgebras  $(X, c)$ ,  $\eta \circ h_{S^E}^c = (h_S^c)^E$ .

2. For all  $n \geq 0, z \in Z_{S^E}$  and  $e \in E$ ,  $\pi_S^n(\eta(z)(e)) = \pi_{S^E}^n(z)(e)$ .

$$\begin{array}{ccc}
 & (SX)^E & \\
 h_{S^E}^c \swarrow & & \searrow (h_S^c)^E \\
 Z_{S^E} & \xrightarrow{\eta} & (Z_S)^E \\
 \pi_{S^E}^n \searrow & & \swarrow (\pi_S^n)^E \\
 & (ST^{n1})^E &
 \end{array}$$

*Proof.* We define  $\eta : Z_{S^E} \rightarrow (P_S)^E$  by

$$\pi_S^n(\eta(z)(e)) = \pi_{S^E}^n(z)(e) \quad (6.2)$$

Since  $z \in Z_{S^E}$ , for every  $n \geq 0$  we have  $(ST^{n!})^E \pi_{S^E}^{n+1}(z) = \pi_{S^E}^n(z)$ , that is to say that  $(ST^{n!}) \circ \pi_{S^E}^{n+1}(z) = \pi_{S^E}^n(z)$ . So the sequence in the right hand of (6.2) is actually in  $(P_S)^E$ .

From the definition of  $\eta$  the second condition of the Lemma is immediately satisfied. We also note that it makes  $\eta$  well-defined.

For all  $n \geq 0$ ,  $(\pi_S^n)^E(\eta \circ h_{S^E}^c) = \pi_S^n \circ \eta \circ h_{S^E}^c = \pi_{S^E}^n \circ h_{S^E}^c = (Sh_n^c)^E = (\pi_S^n \circ h_S^c)^E = (\pi_S^n)^E(h_S^c)^E$ . Since they agree on all the projections, it follows that  $\eta \circ h_{S^E}^c = (h_S^c)^E$ , and also proves that the image of  $\eta$  is in  $(Z_S)^E$ .  $\square$

**Lemma 6.3.** *Let  $\mathcal{P}S \in \text{lng}(T)$ . There is a function  $\zeta : Z_{\mathcal{P}S} \rightarrow \mathcal{P}(Z_S)$  such that*

1. For all  $T$ -coalgebras  $(X, c)$ ,  $\zeta \circ h_{\mathcal{P}S}^c = \mathcal{P}h_S^c$ .

2. For all  $n \geq 0$ ,  $\mathcal{P}\pi_S^n \zeta = \pi_{\mathcal{P}S}^n$ .

*Proof.* Let  $z \in Z_{\mathcal{P}S}$ . We set

$$\zeta(z) = \{u \in Z_S \mid \forall n \geq 0 \pi_S^n(u) \in \pi_{\mathcal{P}S}^n(z)\} \quad (6.3)$$

Given  $x \in \mathcal{P}SX$  such that  $z = h_{\mathcal{P}S}(x)$ ,

$$\begin{aligned} \zeta(z) &= \zeta h_{\mathcal{P}S}(x) \\ &= \{u \in Z_S \mid \forall n \geq 0, \pi_S^n(u) \in \pi_{\mathcal{P}S}^n h_{\mathcal{P}S}(x)\} \\ &= \{u \in Z_S \mid \forall n \geq 0, \pi_S^n(u) \in \mathcal{P}Sh_n(x)\} \\ &= \{u \in Z_S \mid \forall n \geq 0, \pi_S^n(u) \in \mathcal{P}(\pi_S^n h_S)(x)\} \\ &= \{u \in Z_S \mid \forall n \geq 0, \pi_S^n(u) \in \mathcal{P}(\pi_S^n) \mathcal{P}(h_S)(x)\} \end{aligned}$$

Using the Lemma 6.1, we get that the condition defining the set in the last line is equivalent to  $u \in \mathcal{P}h_S(x)$ , so  $\zeta \circ h_{\mathcal{P}S} = \mathcal{P}(h_S)$ , and we also get that  $\zeta(z)$  is a finite set for all  $z \in Z_{\mathcal{P}S}$ .

To prove the second part, we use the fact that every  $z \in Z_{\mathcal{P}S}$  is  $z = h_{\mathcal{P}S}^c(x)$  for some coalgebra  $(X, c)$  and  $x \in \mathcal{P}SX$ . So for all  $n \geq 0$ ,

$$\begin{aligned} (\mathcal{P}\pi_S^n) \circ \zeta(h_{\mathcal{P}S}(x)) &= (\mathcal{P}\pi_S^n)(\mathcal{P}h_S)(x) && \text{by the first part of this Lemma} \\ &= (\mathcal{P}Sh_n)(x) && \text{by (5.1)} \\ &= \pi_{\mathcal{P}S}^n h_{\mathcal{P}S}(x) && \text{again by (5.1)} \\ &= \pi_{\mathcal{P}S}^n(z) \end{aligned}$$

□

**Lemma 6.4.** *Let  $\mathcal{D}S \in \mathbf{Ing}(T)$ . There is a function  $\theta : Z_{\mathcal{D}S} \rightarrow \mathcal{D}(Z_S)$  such that*

1. *For all  $T$ -coalgebras  $(X, c)$ ,  $\theta \circ h_{\mathcal{D}S}^c = \mathcal{D}h_S^c$ .*

2. For  $n \geq 0$ ,  $\mathcal{D}\pi_S^n \theta = \pi_{\mathcal{D}S}^n$

*Proof.* Given  $z \in Z_{\mathcal{D}S}$ , let  $Q_z = \{u \in Z_S \mid \forall n \geq 0 \pi_S^n(u) \in \text{Supp } \pi_{\mathcal{D}S}^n(z)\}$ . (This will be the support of  $\theta(z)$ ).

Claim:  $Q_z$  is a finite set. Every  $z \in Z_{\mathcal{D}S}$  is  $h_{\mathcal{D}S}(\mu)$  for some  $\mu \in \mathcal{D}SX$ . We will use as an auxiliary construction  $\mathcal{P}SX$ , the finite powerset of  $SX$ , and  $Z_{\mathcal{P}S}$ , together with the maps  $h_{\mathcal{P}S}$  and  $\pi_{\mathcal{P}S}^n$ , although  $\mathcal{P}S$  may not be an actual ingredient of  $T$ .

For each set  $X$ ,  $\text{Supp}_X : \mathcal{D}X \rightarrow \mathcal{P}X$  is the application that sends each  $\mu \in \mathcal{D}X$  to the finite set  $\{x \in X : \mu(x) > 0\}$ . It is easy to check that  $\text{Supp}$  is a natural transformation. Now consider the diagram:

$$\begin{array}{ccc}
 \mathcal{D}SX & \xrightarrow{\text{Supp}_{SX}} & \mathcal{P}SX \\
 \mathcal{D}Sh_n \downarrow & \searrow h_{\mathcal{D}S} & \downarrow \mathcal{P}Sh_n \\
 & Z_{\mathcal{D}S} & \\
 \mathcal{D}ST^{n_1} & \xrightarrow{\text{Supp}_{ST^{n_1}}} & \mathcal{P}ST^{n_1} \\
 \downarrow \pi_{\mathcal{D}S}^n & \nearrow \pi_{\mathcal{D}S}^n & \downarrow \pi_{\mathcal{P}S}^n \\
 & Z_{\mathcal{P}S} & 
 \end{array}$$

The square commutes because of the naturality of  $\text{Supp}$ , and the two triangles commute because of (5.1). Now we calculate:

$$\begin{aligned}
 Q_{h_{\mathcal{D}S}(x)} &= \{u \in Z_S \mid \forall n \geq 0 \pi_S^n(u) \in \text{Supp}_{ST^{n_1}} \pi_{\mathcal{D}S}^n h_{\mathcal{D}S} \mu\} \\
 &= \{u \in Z_S \mid \forall n \geq 0 \pi_S^n(u) \in (\mathcal{P}Sh_n) \text{Supp}_X \mu\} \\
 &= \{u \in Z_S \mid \forall n \geq 0 \pi_{\mathcal{P}S}^n(u) \in \pi_{\mathcal{P}S}^n h_{\mathcal{P}S} \text{Supp}_X \mu\} \\
 &= \{u \in Z_S \mid u \in h_{\mathcal{P}S} \text{Supp}_X \mu\} && \text{by Lemma 6.1} \\
 &= h_{\mathcal{P}S} \text{Supp}_{SX} \mu
 \end{aligned}$$

To define  $\theta : Z_{\mathcal{D}S} \rightarrow \mathcal{D}(Z_S)$  one must first consider for each  $z \in Z_{\mathcal{D}S}$  the set  $Q_z$ . Since this is equal to  $h_{\mathcal{P}S} \text{Supp}_{SX} \mu$ , it is finite, so there's a number  $n$  such that if  $u, u' \in Q_z$ , and  $u \neq u'$ , then



$\pi^n(u) \neq \pi^n(u')$ . Let  $N$  be the first such number. Then

$$\theta(z)(u) = \begin{cases} \pi_{\mathcal{D}S}^N(z)(\pi_S^N(u)) & \text{if } u \in Q_z \\ 0 & \text{otherwise.} \end{cases}$$

If  $u \in Q_z$ , then

$$\begin{aligned} \theta(h_{\mathcal{D}S}(x))(u) &= (\pi_{\mathcal{D}S}^N h_{\mathcal{D}S}(x))(\pi_S^N(u)) \\ &= \mathcal{D}Sh_N(x)(\pi_S^N(u)) \\ &= x(Sh_N)^{-1}(\pi_S^N(u)) \end{aligned}$$

On the other hand,  $(\mathcal{D}h_S)(x)(u) = x(h_S)^{-1}(u)$ . So if we prove that  $(Sh_N)^{-1}(\pi_S^N(u)) = h_S^{-1}(u)$  we'll have proved that  $\theta h_{\mathcal{D}S} = \mathcal{D}h_S$ . As a consequence, this will also prove that for all  $z \in Z_{\mathcal{D}S}$ ,  $\theta(z)$  is a discrete measure.

If  $s \in h_S^{-1}(u)$ , then  $h_S(s) = u$  and therefore  $\pi_S^N h_S(s) = Sh_N(s) = \pi_S^N(u)$ , i.e.  $s \in (Sh_N)^{-1}(\pi_S^N(u))$ .

Now we'll assume that  $\pi_S^N(h_S)(s) = Sh_N(s) = \pi_S^N(u)$ , and we'll prove that  $h_S(s) = u$ . This will be accomplished by showing that for all  $n$ ,  $\pi_S^n h_S(s) = \pi_S^n(u)$ . We have two cases:

*Case 1:*  $n < N$ . From  $Sh_N(s) = \pi_S^N(u)$  it follows that

$$\pi_S^{N-1} h_S(s) = Sh_{N-1}(s) = S(T^{N-1}! \circ h_N)(s) = ST^{N-1}! \pi_S^N(u) = \pi_S^{N-1}(u).$$

by induction, one can prove the same for all  $n \leq N$ .

*Case 2:*  $n \geq N$ . We have chosen  $N$  so that if  $u, u' \in Q_z$  are different, then  $\pi_S^N(u) \neq \pi_S^N(u')$ . but it follows that for all  $n \geq N$ ,  $\pi_S^n(u) \neq \pi_S^n(u')$ . Otherwise, we'd have  $\pi_S^N(u) = S\tau_{nN}\pi_S^n(u) = S\tau_{nN}\pi_S^n(u') = \pi_S^N(u')$ , contradiction.

Then, if for some  $n \geq N$  we had  $\pi_S^n h_S(s) \neq \pi_S^n(u)$ , then  $h_S(s) \neq u$ , so  $\pi_S^N(h_S(s)) \neq \pi_S^N(u)$ , contradicting the hypothesis. This concludes the proof of part 1.

Finally, to prove the second part of the Lemma, let  $v \in SF^n 1$ .

$$\begin{aligned} \mathcal{D}\pi_S^n \theta(z)(v) &= \theta(z)(\pi_S^n)^{-1}(v) \\ &= \pi_{\mathcal{D}S}^n(z)(\pi_S^n[(\pi_S^n)^{-1}(v)]) \\ &= \pi_{\mathcal{D}S}^n(z)(v) \end{aligned}$$

□

**Lemma 6.5.** *There is a family of maps  $r_S : Z_S \rightarrow S(Z)$  indexed by the ingredients of  $T$  such that the following hold:*

1. *For all  $T$ -coalgebras  $(X, c)$  and all  $S \in \text{Ing}(T)$ ,  $r_S \circ h_S^c = Sh^c$ .*
2. *For all  $n \geq 0$ ,  $S\pi_n r_S = \pi_S^n$ .*

*Proof.* By induction on the ingredients  $S$  of  $T$ . The base cases and the induction steps for products, and coproducts are the same as in Lemma 5.6.

We treat in detail the induction step for functors  $\mathcal{P}S$ . Let  $r_{\mathcal{P}S}$  be  $\mathcal{P}r_S \circ \zeta$ . Then for part 1 we have that

$$r_{\mathcal{P}S} \circ h_{\mathcal{P}S}^c = \mathcal{P}r_S \circ \zeta \circ h_{\mathcal{P}S}^c = \mathcal{P}r_S \circ \mathcal{P}h_S^c = \mathcal{P}(r_S \circ h_S^c) = \mathcal{P}Sh_{Id}^c.$$

For part 2 we have:

$$\begin{aligned}
\mathcal{P}S\pi_n r_{\mathcal{P}S} &= \mathcal{P}(S\pi_n) \circ (\mathcal{P}r_S) \circ \zeta && \text{by the definition of } r_{\mathcal{P}S} \\
&= \mathcal{P}(S\pi_n \circ r_S) \circ \zeta \\
&= \mathcal{P}(\pi_S^n) \circ \zeta && \text{by inductive hypothesis} \\
&= \pi_{\mathcal{P}S}^n && \text{by Lemma 6.3}
\end{aligned}$$

In a similar manner, we define  $r_{SE}$  by  $r_{SE} = r_S^E \circ \eta$  and  $r_{DS}$  by  $r_{DS} = (\mathcal{D}r_S) \circ \theta$ . □

The rest of the results of Chapter 5 are valid in Set, yielding the following Theorem:

**Theorem 6.1.** *All the probabilistic Kripke polynomial functors have a final coalgebra.*

## Type Spaces

### 7.1 Motivation

The theory of type spaces has its origins in game theory. The intuitive idea is that a type describes a player. A player in a game can be optimistic, pessimistic, cautious, daring, suspicious, paranoid, etc. To get a mathematical definition, we need to be clear on which kind of games we are talking about, and then we can proceed to see how we can describe the ‘type’ of a player. We will not go into too much detail about game theory, but just enough to understand the setting in which type spaces originated.

**Definition 7.1.** (following [OR94]) An *extensive game with perfect information*  $G = (N, A, H, P, (U_n)_{n \in N})$  consists of:

- A set  $N$ , the set of players.
- A set  $A$ , the set of actions.
- A set  $H$  of sequences (finite or infinite) of elements in  $A$  that satisfies the following three properties:


- The empty sequence  $\emptyset$  is in  $H$ .
- If  $(a_k)_{k=1,\dots,K} \in H$  (where  $K$  may be infinite) and  $L < K$  then  $(a_k)_{k=1,\dots,L} \in H$ .
- If an infinite sequence  $(a^k)_{k=1}^\infty$  is such that  $(a^k)_{k=1,\dots,L} \in H$  for every positive integer  $L$ , then  $(a^k)_{k=1}^\infty \in H$ .

The members of  $H$  are called *histories*. A history  $(a^k)_{k=1,\dots,K} \in H$  is *terminal* if it is infinite or there is no  $a^{K+1}$  such that  $(a^k)_{k=1,\dots,K+1} \in H$ . The set of terminal histories is denoted with  $Z$ . The set of actions available after the nonterminal history  $h$  is denoted  $A(h) = \{a \in A \mid (h, a) \in H\}$ .

- A function  $P : H \setminus Z \rightarrow N$ , that indicates for each non-terminal history in  $H$  which one of the players takes an action after the history.
- Functions  $U_n : Z \rightarrow \mathbb{R}$  for  $n \in N$  that give for each terminal history and each player, the *payoff* of that player after that history.

The set  $H$  can be seen as a tree with root  $\emptyset$ , and with its nodes labeled by the function  $P$ , and the leaves labeled by the functions  $U_n$ . We indicate the elements  $a^k$  on the edges of the tree so following a particular branch from the root will give the history that names each node.

**Example 7.1.**



In the diagram above we have a game where  $N = \{1, 2, 3\}$ ;  $P(\emptyset) = 1$  meaning that player 1 gets to decide the first move in the game, and has three options available:  $l, c, r$  (the letters stand for left, center or right, respectively). If player 1 chooses  $l$  or  $c$ , then player 2 decides what's the next action, and she has options  $l$  and  $r$  available. If player 1 chooses  $r$  instead, it is player 3 who decides what's the final move. Under each terminal node in the tree, a triple indicates the values of the utility functions  $U_1, U_2$  and  $U_3$ . So, for example if the history of the game is  $(c, l)$ , then player 1 gets a payoff of 3, while players 2 and 3 get a payoff of 1 each.

Alternatively, extensive games with complete information can be given by indicating a family of preorders  $(\prec_n)_{n \in N}$  that indicate the *preferences* of the players. For our purposes, it will be enough to assume that all players prefer to maximize their payoffs and are indifferent to what other players' payoffs are.

*Games with incomplete information* are games in which the incompleteness of the information arises in three main ways.

1. The players may not know the *physical outcome function* of the game which specifies the physical outcome produced by each strategy available to the players.
2. The players may not know their own or some other players' *utility functions*, which specify the utility payoff that a given player  $i$  derives from every physical outcome.
3. The players may not know their own or some other players' *strategy space*, i.e. the set of all strategies available to various players.


'All other causes of incomplete information can be reduced to these three basic cases— indeed sometimes this can be done in two or more different (but essentially equivalent) ways' [Har67]. The challenge is to be able to take the best possible decisions when these uncertainties are present.

A breakthrough in this field was made in 1967, when a series of papers by John C. Harsanyi, [Har67, Har68a, Har68b] saw print. The idea was to tame the uncertainty by transforming the games with incomplete information into games with complete but *imperfect* information.

**Definition 7.2.** An extensive game with *imperfect information* is a game  $G = (N, A, H, P, U_n, \mathcal{J}_n)$  where  $N, A, H, P$  and  $U_n$  are as in Definition 7.1, and for each player  $n \in N$ ,  $\mathcal{J}_n$  is a partition on the set  $H_n = \{h \in H \setminus Z : P(h) = n\}$  such that for two elements  $h, h'$  in the same component of the partition,  $A(h) = A(h')$ . The equivalence classes in this partitions are called *information sets*.

The idea here is that player  $n$  knows in which information set the game currently is, but doesn't know exactly the whole history that has lead the game into that set. Note that the players still have perfect information. They know the payoffs in all the possible outcomes.

**Example 7.2.**



Now the dotted line indicates that the set  $\{l, c\}$  is an information set for player 2. She does not have information about whether player 1 moved to the right or to the center, but she does know what the payoffs will be in each case, and also knows that, since it's her turn, player 1 did not choose  $r$ .

If all the information sets contain exactly one node of the tree, we have a game with perfect information. The information sets allow us to represent games in which the players make their moves simultaneously (and thus don't know when making their decision what are the other players'

---

moves), and also to represent situations in which “nature” or “chance” makes a move we cannot predict. This feature will be exploited later.

Some further assumptions are made about the games under study. In the first place, it is assumed that the beliefs the players of the game have can be represented through probability measures (this is called the *Bayesian approach*). It is also assumed that the players are aware of the extent of the knowledge or ignorance of the other players, and that they will always act “rationally”, that is, they will take the action that gives them the highest possible expected payoff, based on the information available to them. The notion of rationality is quite hard to formulate and still topic of debate among game theorists (see, e.g., [Bra04]).

In Harsanyi’s words, [Har67]:

It seems to me that the basic reason why the theory of games with incomplete information has made so little progress so far lies in the fact that these games give rise, or at least appear to give rise, to an infinite regress in reciprocal expectations on the part of the players.

The argument is the following: suppose the game has incomplete information and just two players. Player 1 has some beliefs about what are the actual values of the missing information. This is represented as a probability measure over the space of all possible values the unknown could take. Player 1 also knows that player 2 cannot know the actual value and hence resorts to using a probability distribution representing her beliefs as well. In order to take a decision, player 1 then must form some mental model of what player 2’s beliefs are. Player 2’s beliefs include those that, in turn, player 2 has about player 1’s beliefs. This kind of reasoning promptly leads to an infinite regression of unfolding beliefs. Harsanyi calls any model of this kind a *sequential-expectations* model for games with incomplete information.



---

Harsanyi was concerned with finding ways of analyzing these games with incomplete information. The solution he offered involved the construction of a game with complete but imperfect information based on the given one with incomplete information. In the new game, there are new chance moves that are assumed to occur before the two players choose their strategies. In these random moves, the actual payoff of the two players are determined, but being a game with imperfect information, the players only know they are in some information set, and a probability distribution for the random moves (this probability distribution is assumed to be common knowledge to all the players). Using conditional probabilities, they can then derive the different expected values they need to assess the strategies to be taken in the game.

There is an alternative interpretation of the random moves added to the game, which originates the intuition in which we base our model. Instead of assuming that they determine important characteristics of the players (in particular, their payoffs), it could be assumed that the players themselves are being chosen at random from ‘certain hypothetical populations containing individuals of different “types”, each possible “type” of a player  $i$  being characterized by a different attribute vector  $c_i$ , i.e., by a different combination of production costs, financial resources, and states of information’ [Har67].

It is these populations that we’ll call *type spaces*, and their elements will be of course, *types*. While Harsanyi assumes the type space was given, he already suggested they could be constructed from the considerations about beliefs explained above:

As we have seen, if we use the Bayesian approach, then the sequential-expectations model for any given [incomplete information] game  $G$  will have to be analyzed in terms of infinite sequences of higher and higher-order subjective probability distributions, i.e. subjective probability distributions over subjective probability distributions [Har67].

Harsanyi was discouraged from this approach by the technical difficulties it presented:

Probability distributions over some space of payoff functions or of probability distributions, and more generally probability distributions over function spaces, involve certain mathematical difficulties [...]. However, as Aumann has shown [Aum61] and [Aum64], these mathematical difficulties can be overcome. But even if we succeed in defining the relevant higher order probability distributions in a mathematically admissible way, the fact remains that the resulting model –like *all* models based on the sequential-expectations approach–will be extremely complicated and cumbersome.

The difficulty pointed out by Aumann in [Aum61] is that if  $X$  and  $Y$  are measurable spaces and we denote by  $Y^X$  the set of all measurable functions from  $X$  to  $Y$ , then there is no natural way of endowing  $Y^X$  with a  $\sigma$ -algebra that makes the evaluation function  $ev : Y^X \times X \rightarrow Y$  given by  $ev(f, x) = f(x)$  measurable. Aumann proposes in [Aum64] to choose a single real number that represents a probability distribution. In our approach, the problem is overcome by considering the spaces  $\Delta X$  instead of looking at all the measurable functions in  $[0, 1]^X$  that have integral 1 over  $X$ .

So, to formalize the notion of types that Harsanyi had in mind, we want a mathematical object, the *type space*, such that each element or *type* will have associated to it, in a natural way, beliefs (represented by probability distributions) over the states of nature and the types of the other players in the game. In a game with  $N$  players, each player will assume one of the types  $t \in T$ , as if they were roles in a play.

A first approach would be to solve the equation  $T \cong \Delta(S \times T)$ , where the set  $T$  would be the type space and  $S$  the *states of nature*. The states of nature are the possible values the unknown variables in the game can take. We want both  $S$  and  $T$  to be measurable spaces so we can define probability measures on them. Let  $m : T \rightarrow \Delta(S \times T)$  be the desired isomorphism. Then for each

$t \in T$ ,  $m(t)$  represents the beliefs of a player of type  $t$ .

There are some problems with this approach. If the game has  $N$  players, then each player  $i$  with type  $t_i$  should have beliefs about the types of all the other players, so the equation to solve could be

$$T \cong \Delta(S \times T^N). \quad (7.1)$$

Furthermore, we want each type to know what his own type is, so we don't want  $T$  to be isomorphic to  $\Delta(S \times T^N)$ , but to the subset of  $\Delta(S \times T^N)$  of probability distributions in which the marginal of each  $m(t_i)$  on the  $i$ -th copy of  $T$  is the distribution  $\delta_{t_i}$  which has support on the point  $t_i$ . Adding this extra condition to the definition would steer us away from the definition of coalgebras on  $\mathbf{Meas}$ , but we can overcome this difficulty by changing the functor in an appropriate way. The key observation here is that for any product of measurable spaces  $A \times B$  and  $b_0 \in B$  such that the singleton  $\{b_0\}$  is measurable, there is an isomorphism between the spaces  $\{\mu \in \Delta(A \times B) : \text{mar}_B \mu = \delta_{b_0}\}$  and  $\Delta A$ .

Recall that a probability measure  $\mu$  on  $A \times B$  induces, via the projections, a measure on each of the factor spaces. These measures are called *marginals*, and denoted by  $\text{mar}_A \mu = (\Delta \pi_A) \mu = \mu \circ \pi_A^{-1}$ ;  $\text{mar}_B \mu = (\Delta \pi_B) \mu = \mu \circ \pi_B^{-1}$ .

The following Lemma proves that in the case above, it is enough to know the marginals to determine the measure.

**Lemma 7.1.** *Let  $\mu$  be a probability measure on a product measurable space  $A \times B$ . If  $\text{mar}_B \mu = \delta_{b_0}$  for some  $b_0 \in B$ , then  $\mu = \text{mar}_A \mu \times \delta_{b_0}$ .*

*Proof.* We only need to prove it for rectangles  $G \times F$ , where  $G$  is a measurable subset of  $A$  and  $F$  is a measurable subset of  $B$ .

We want to prove that  $\mu(G \times F) = (\text{mar}_A \mu)(G) \times \delta_{b_0}(F)$ . We have two cases: if  $b_0 \notin F$ , this

reduces to proving that  $\mu(G \times F) = 0$ , and if  $b_0 \in F$ , then we want to show that  $\mu(G \times F) = \text{mar}_A \mu(G) = \mu(\pi_A^{-1}(G)) = \mu(G \times B)$ .

Notice first that for  $\mu(G \times B) = \mu(\pi_A^{-1}(G)) = \text{mar}_A \mu(G) = \text{mar}_A \mu(G) \times \delta_{b_0}(B)$ . Also  $\mu(A \times F) = \text{mar}_B \mu(F) = \delta_{b_0}(F) = (\text{mar}_A \mu)(A) \times \delta_{b_0}(F)$ .

Now we can prove that if  $b_0 \notin F$ , then  $\mu(G \times F) \leq \mu(A \times F) = 0$ , and if  $b_0 \in F$ , then  $\mu(G \times F) = \mu(G \times \{b_0\}) + \mu(G \times (F \setminus \{b_0\})) \leq \mu(G \times \{b_0\}) + \mu(A \times (F \setminus \{b_0\})) = \mu(G \times \{b_0\})$ . On the other hand,  $\mu(G \times B)$  is also equal to  $\mu(G \times \{b_0\}) + 0$ .  $\square$

Note that even though the Lemma requires the singletons in  $B$  to be measurable, once we decide to model types using the isomorphism, we can drop the condition. Now we can model the introspection condition by considering equations like

$$T \cong \Delta(S \times T^{N-1}). \quad (7.2)$$

The problem of finding a *universal type space*, that is, a type space containing all the possible types a player could adopt, could be solved by finding the final coalgebra for the functor  $F(X) = \Delta(S \times X^{N-1})$ . This can be done using Theorem 4.1 or Theorem 5.1. Lambek's Lemma 2.3 provides the isomorphism we are looking for.

But when we look at a single coalgebra for this functor, that is, a measurable map  $m : T \rightarrow \Delta(S \times T^{N-1})$  we get a somewhat unsatisfactory model. Why should all the players come from the same type space? It would be better to be more general and to assume that there are type spaces  $T_1, T_2, \dots, T_N$  and the type of player  $i$  is selected from the corresponding  $T_i$ .

**Definition 7.3.** Let  $\text{Meas}^N$  be the  $N$ -fold product of the category  $\text{Meas}$ . Each object  $M$  in  $\text{Meas}^N$  is a  $N$ -tuple of measurable spaces  $(M_1, \dots, M_N)$ , and the morphisms are  $N$ -tuples of measurable

functions  $f_i : M_i \rightarrow M'_i$ . Let  $Proj_i^N : \text{Meas}^N \rightarrow \text{Meas}$  be the  $i$ -th projection functor.

**Definition 7.4.** We define then a *type space* for a game with  $N$  players over the measurable space  $S$  of states of nature, as a coalgebra for the endofunctor in  $\text{Meas}^N$  given by  $T = (T_1, T_2, \dots, T_N)$  where for  $1 \leq i \leq N$ ,

$$T_i = \Delta(S \times \prod_{j \neq i} Proj_j^N). \quad (7.3)$$

The diagram for a coalgebra  $(X, m)$  of this functor is:

$$\begin{array}{ccccccc} (X_1 & , & X_2 & , & \dots & , & X_N \\ m_1 \downarrow & & m_2 \downarrow & & & & m_N \downarrow \\ (\Delta(S \times \prod_{j \neq 1} X_j)) & , & \Delta(S \times \prod_{j \neq 2} X_j) & , & \dots & , & \Delta(S \times \prod_{j \neq N} X_j) \end{array}$$

The definition above is a particular case of the more general one that follows.

## 7.2 Measure polynomial functors in many variables

**Definition 7.5.** A *measure polynomial functor in many variables*  $T : \text{Meas}^N \rightarrow \text{Meas}$  is a functor built from the functors  $Proj_1^N, \dots, Proj_N^N$  and constant functors for measurable spaces, using either products, coproducts and  $\Delta$ . For any natural number  $N'$ , we can extend the notion of a measure polynomial functor to functors  $T = (T_1, \dots, T_{N'}) : \text{Meas}^N \rightarrow \text{Meas}^{N'}$  such that each  $T_i, 1 \leq i \leq N'$ , is a measure polynomial functor in many variables from  $\text{Meas}^N$  to  $\text{Meas}$  as defined above.

**Example 7.3.** For a fixed measurable space  $M$ , consider the polynomial functor in three variables  $F : \text{Meas}^3 \rightarrow \text{Meas}^2$  given by:

$$F = ( \Delta(Proj_1^3 + Proj_2^3) , ((\Delta Proj_3^3) \times Proj_2^3) + M )$$

**Definition 7.6.** The *ingredients* of a measure polynomial functor in many variables  $T = (T_1, T_2, \dots, T_{N'}) : \text{Meas}^N \rightarrow \text{Meas}^{N'}$  are defined by:

- $\text{Ing}(T) = \cup_{i=1}^{N'} \text{Ing}(T_i)$
- $\text{Ing}(Id_i) = \{Proj_j^N : 1 \leq j \leq N\}$
- $\text{Ing}(M) = \{M, Proj_j^N : 1 \leq j \leq N\}$
- $\text{Ing}(U \times V) = \{U \times V\} \cup \text{Ing}(U) \cup \text{Ing}(V)$
- $\text{Ing}(U + V) = \{U + V\} \cup \text{Ing}(U) \cup \text{Ing}(V)$
- $\text{Ing}(\Delta U) = \{\Delta U\} \cup \text{Ing}(U)$

$\text{Ing}(T)$  is a finite set of functors from  $\text{Meas}^N$  to  $\text{Meas}$ .

We are going to center our attention on measure polynomial functors in many variables that are endofunctors of the category  $\text{Meas}^N$ , and the coalgebras for those functors.

**Theorem 7.1.** *If  $T : \text{Meas}^N \rightarrow \text{Meas}^N$  is a measure polynomial functor in many variables, then it has a final coalgebra.*

*Proof.* We could prove this theorem using the methods from Chapter 4 or Chapter 5. Using the method of modal languages, we have to start by defining for each  $i, 1 \leq i \leq N$  new sorts  $Proj_i^N$ , the formulas  $\text{true}_i : Proj_i^N$  and modal operators  $[\text{next}_i]$  so that if  $\varphi : T_i$ , then  $[\text{next}_i]\varphi : Proj_i^N$ , with semantics given by

$$\llbracket \text{true}_i \rrbracket_{Proj_i^N}^m = X_i,$$

$$\llbracket [\text{next}_i]\varphi \rrbracket_{Proj_i^N}^m = m_i^{-1} \llbracket \varphi \rrbracket_{T_i}^m$$

for every  $T$ -coalgebra  $(X, m)$ .

For each ingredient  $S$  of  $T$ , the measurable spaces  $S^*$  are defined as in Definition 4.5. Let  $d_i^m : X_i \rightarrow (Proj_i^N)^*$  be the description map:

$$d_i^m(x_i) = \{\varphi : Proj_i^N \mid x_i \in \llbracket \varphi \rrbracket_{Proj_i^N}^m\}$$

These descriptions maps establish the function  $d_{Id}^m : X \rightarrow Id^*$ , where  $Id^* = ((Proj_1^N)^*, \dots, (Proj_N^N)^*)$ .

In a similar way, we have the maps  $d_S^m : SX \rightarrow S^*$  for each ingredient  $S$ .

For each  $i$ , we let  $g_i : (Proj_i^N)^* \rightarrow T_i^*$  be defined by

$$g_i(s) = \{\varphi : T_i \mid \llbracket \text{next}_i \rrbracket \varphi \in s\} \quad (7.4)$$

for every  $s \in (Proj_i^N)^*$ . Using the same arguments as in Lemma 4.3, we get that for each  $i$  the following diagram commutes.

$$\begin{array}{ccc} X_i & \xrightarrow{m_i} & T_i X \\ d_{Proj_i^N}^m \downarrow & & \downarrow d_{T_i}^m \\ Proj_i^* & \xrightarrow{g_i} & T_i^* \end{array}$$

Therefore, the diagram

$$\begin{array}{ccc} X & \xrightarrow{m} & TX \\ d_{Id}^m \downarrow & & \downarrow d_T^m \\ Id^* & \xrightarrow{g} & T^* \end{array}$$

also commutes. Here  $T^* = (T_1^*, \dots, T_N^*)$ ,  $g = (g_1, \dots, g_N)$  and  $d_T^m = (d_1^m, \dots, d_N^m)$ .

Defining  $r_T$  as  $N$ -tuple of the maps  $r_{T_i} : T_i^* \rightarrow T_i(Id^*)$ , and letting  $m^* = r_T \circ g$ , we get that  $d_{Id}^m : (X, m) \rightarrow (Id^*, m^*)$  is a  $T$ -coalgebra morphism. Furthermore, using Lemma 4.10 for each component of  $d_{Id}^m$ , this function is the identity on  $Id^*$ , from which it follows that  $(Id^*, m^*)$  is a final coalgebra for  $T$ .  $\square$

### 7.3 Universal type spaces

Going back to the type spaces for a game with  $N$  players, application of the Theorem above yields a final type space, also known in the literature as *universal type space*. We also get the following Lemma:

**Lemma 7.2.** *If  $T : \text{Meas}^N \rightarrow \text{Meas}^N$  is the functor given by  $(T_i = \Delta(S \times \prod_{j \neq i} \text{Proj}_j^N))_{1 \leq i \leq N}$ , then for each  $i$ ,  $(\text{Proj}_i^N)^*$  is isomorphic to  $\Delta(S \times \prod_{j \neq i} (\text{Proj}_j^N)^*)$  and all the spaces  $(\text{Proj}_i^N)^*$ ,  $1 \leq i \leq N$  are isomorphic.*

*Proof.* By Lambek's Lemma, 2.3,  $Id^*$  is isomorphic to  $T(Id^*)$ , from which follows that for each  $i$ ,  $(\text{Proj}_i^N)^* \cong T_i(Id^*) = \Delta(S \times \prod_{j \neq i} (\text{Proj}_j^N)^*)$ .

For the second assertion, it is enough to notice the symmetry in the definition of the functor  $T$ . The construction of each space  $(\text{Proj}_i^N)^*$  is the same, up to a reassignment of the indices.  $\square$

The fact that all the type spaces in the universal type space for a game with  $N$  players are isomorphic justifies naming it *the* universal type space for the game: each one of the players are of one of the types in this universal space.

**Example 7.4.** Assume that  $N = 4$ , se take a look at some of the formulas for the functor  $T$  and its ingredients.

$$\text{true}_j : \text{Proj}_j^4, 1 \leq j \leq 4$$

For  $A$  measurable in  $S$ ,  $A : S$

$$\langle A, \text{true}_1, \text{true}_3, \text{true}_4 \rangle : S \times \prod_{j \neq 2} \text{Proj}_j^4$$

$$\beta^P \langle A, \text{true}_1, \text{true}_3, \text{true}_4 \rangle : T_2$$



$$[\text{next}]_2^{\beta^p} \langle A, \text{true}_1, \text{true}_3, \text{true}_4 \rangle : \text{Proj}_2^4$$

This last formula expresses that player 2 in the game believes that the event  $A$  has probability bigger or equal than  $p$ . If we name this formula  $\varphi_2$ , then we can build more complex formulas, like

$$[\text{next}]_1^{\beta^q} \langle A, \varphi_2, \text{true}_3, \text{true}_4 \rangle : \text{Proj}_1^4.$$

In this way, we can express beliefs about beliefs, and by further nesting formulas, beliefs about beliefs about beliefs, etc.

It is in this kind of formulas that we see the relevance of using coalgebras in dealing with this problem. The repeated application of the structure map allows us to unfold the different levels of beliefs of the players.

## 7.4 A brief review of the literature on type spaces

There have been several constructions of type spaces and universal type spaces in the literature, each one trying to capture the intuitive idea behind the definition in a slightly different way. Here we review them, as we compare them with the framework we just developed.

### Armbruster, Böge and Eisele

In *Bayesian Game Theory* [AB79], W. Armbruster and W. Böge present their approach to the study of games with unknown utility functions, in which the players “will have at least a subjective probability distribution on [the] alternatives”. This is called the *Bayesian assumption*. In order to construct “canonical representations for the players’ subjective probability measures”, the following notion is introduced, and attributed to Böge, in a lecture on game theory given in 1970.

**Definition 7.7.** Let  $S_1^0, \dots, S_N^0$  be compact Hausdorff spaces. An  $N$ -tuple of compact sets and continuous maps  $(S_1, \dots, S_N, \rho_1, \dots, \rho_N)$  is called an *oracle system* for  $S_1^0, \dots, S_N^0$  if for all  $i$ ,  $\rho_i : S_i \rightarrow S_i^0 \times \prod_{j \neq i} \Delta_r(S_j)$ .

This is the same as saying that  $(S, \rho)$  is a coalgebra for the functor  $T = (S_i^0 \times \prod_{j \neq i} \Delta_r(\text{Proj}_j^N))_{1 \leq i \leq N}$  in the category **CHaus** of compact Hausdorff spaces and continuous functions. The underlying assumption here is that each player has a different space of states of nature  $S_i^0$  in which their unknowns lie.

The final coalgebra is constructed by taking the projective limit of the corresponding final sequence. This final coalgebra is called the *canonical oracle system*. Note that not all the components of the functor are the same, so in general the spaces  $Id_i^*$  will not be isomorphic to each other as in Lemma 7.2. This is a reasonable assumption, and using Theorem 7.1, one can extend the definition and existence of canonical oracle systems to the general case of measurable spaces.

It is important to note that here appears for the first time a coalgebra (not necessarily the final one) as a model of the beliefs of a player. This transcends the idea of just looking for the space of all possible types, to give more restricted models that can be useful to describe situations in more manageable terms.

W. Böge and Th. Eisele present a slightly different approach in the paper *On Solutions of Bayesian Games*, [BE79]. Here again the topological setting is the category **CHaus**. The space over which the behavior of the players is selected is similar to the one we proposed in (7.1), but with certain restrictions.

Given a compact space of states of nature  $R^0$ , a nonempty subspace  $R^1 \subseteq R^0 \times (\Delta_r R^0)^N$  of common a-priori information is selected.

**Definition 7.8.** A system  $(R, \rho)$  with

$$\rho : R \rightarrow R^0 \times (\Delta_r R)^N$$

is called a *system of complete reflections over the information set  $R^1$*  if

$$(1_{R^0} \times (\Delta_r(\pi_{R^0} \circ \rho))) \circ \rho \subseteq R^1 \subseteq R^0 \times (\Delta_r R^0)^N. \quad (7.5)$$

$$\begin{array}{ccc} R & & R \\ \downarrow \rho & & \downarrow \rho \\ R^0 \times (\Delta_r R)^N & & R^0 \times (\Delta_r R)^N \\ \downarrow \pi_{R^0} & & \downarrow 1_{R^0} \times (\Delta_r(\pi_{R^0} \circ \rho))^N \\ R^0 & & R^0 \times (\Delta_r R^0)^N \end{array}$$

The space  $R^1$  has to satisfy a couple of conditions, the first one specifying that each player knows what their beliefs are, and the second one saying that each player will try to maximize their utility function. These requirements preclude the systems of complete reflections from being coalgebras. We have seen before how the first condition, of each player knowing their beliefs, can be dealt with by taking a different functor.

The construction of the final object in the category of systems of complete reflections is done by taking the projective limit, and restricting the spaces so that the image of the map  $\rho$  for the final object is contained in  $R^1$ . It would be interesting to adapt our methods from Chapters 4 and 5 to include this kind of restrictions.

## Mertens and Zamir

The paper *Formulation of Bayesian Analysis for Games with Incomplete Information* by Jean-François Mertens and Shmuel Zamir, [MZ84], is the most often cited one in the literature about type spaces.

Starting from a compact space  $S$  called *parameter-space* or set of states of nature, they seek to define a set  $Y$  of the “states of the world” in which every point contains all characteristics, beliefs and mutual beliefs of all players. The equations that summarize their goals are:

$$Y = S \times T^N \tag{7.6}$$

$$T = \text{the set of all probability distributions on } (S \times T^{N-1}) \tag{7.7}$$

These equations are, of course, intended to be solved up to isomorphism. Equation (7.7) is essentially our (7.2). Some of the definitions in this work are interesting and we will analyze them here, trying to understand their motivation and how they are accounted for in our model.

**Definition 7.9.** [MZ84] Let  $S$  be a compact space. An  $S$ -based abstract beliefs space (*BL-space*) is an  $(N + 3)$  tuple  $(C, S, f, (t^i)_{i=1}^N)$  where  $C$  is a compact set,  $f$  is a continuous mapping  $f : C \rightarrow S$  and  $t^i, i = 1, \dots, N$ , are continuous mappings  $t^i : C \rightarrow \Delta(C)$  (with respect to the weak-\* topology) satisfying:

$$\tilde{c} \in C \text{ and } \tilde{c} \in \text{Supp}(t^i(c)) \Rightarrow t^i(\tilde{c}) = t^i(c). \tag{7.8}$$

The condition (7.8) specifies that “a player assigns positive probability (in the discrete case) only to those points in  $C$  in which he has the same beliefs. In other words, he is certain of his own

beliefs.” It can be rewritten as:

$$\tilde{c} \in C \text{ and } \tilde{c} \in \text{Supp}(t^i(c)) \Rightarrow \tilde{c} \in (t^i)^{-1}[t^i(c)].$$

Or the following equivalent equations:

$$\text{Supp}(t^i(c)) \subseteq (t^i)^{-1}[t^i(c)]$$

$$t^i(c)[(t^i)^{-1}[t^i(c)]] = 1$$

$$(\Delta t^i)t^i(c) = \delta_{t^i(c)}.$$

Thus, even though the first impression could be that Belief spaces are coalgebras for the functor  $FX = S \times \Delta X$ , we see immediately that we need the function  $f$  to have the specific codomain  $S$ , and we need many different functions  $t^i$  with codomain  $\Delta C$ .

However, we can see that an adaptation from our definitions yields spaces with the same properties. Furthermore, we can drop the requirements about compactness for the space  $S$  and continuity for the functions. If  $(X, m)$  is a type space for a game over  $S$  with  $N$  players, as in Definition 7.4, then let

$$C = S \times \prod_{i=1}^N X_i$$

$$C_{-i} = S \times \prod_{j \neq i} X_j$$

Let  $\pi_i$  and  $\pi_{-i}$  be the projections from  $C$  to  $X_i$  and  $C_{-i}$ , respectively. Now for all  $c \in C$ , let  $t^i : C \rightarrow \Delta C$  be defined by

$$t^i(c) = m_i \pi_i(c) \times \delta_{\pi_{-i}(c)}.$$

Thus  $t^i(c) \in \Delta C$ . Letting  $\pi_S : C \rightarrow S$  be the projection, we have that

**Proposition 7.1.**  $(C, S, \pi_S, (t^i)_{i=1}^N)$  is a BL-space.

*Proof.* We only need to check that condition (7.8) is satisfied. Notice that the type spaces of Definition 7.4 are defined for any measurable space  $S$ , and the functions  $m_i$  need not be continuous, just measurable. Condition (7.8) is stated in terms of the support of the probability measure  $t^i(c)$ , which does not necessarily exist in the more general case. We will prove the condition  $t^i(c)[(t^i)^{-1}[t^i(c)]] = 1$  which is equivalent to (7.8) when the support is defined.

$$\begin{aligned} t^i(c)[(t^i)^{-1}(t^i(c))] &= t^i(c)[(t^i)^{-1}(m_i \pi_i(c) \times \delta_{\pi_i(c)})] \\ &= t^i(c)[((m_i \times \delta) \circ \pi_i)^{-1}(m_i \pi_i(c) \times \delta_{\pi_i(c)})] \\ &= t^i(c)[(\pi_i)^{-1}(m_i \times \delta)^{-1}(m_i \pi_i(c) \times \delta_{\pi_i(c)})] \end{aligned}$$

The set  $(m_i \times \delta)^{-1}(m_i \pi_i(c) \times \delta_{\pi_i(c)})$  is not empty, since at least  $\pi_i(c)$  is in it. It is also equal to the set  $m_i^{-1} m_i \pi_i(c) \cap \delta^{-1}(\delta_{\pi_i(c)}) = m_i^{-1} m_i \pi_i(c) \cap \{\pi_i(c)\}$  so its inverse image under  $\pi_i$  is  $C_{-i} \times \{\pi_i(c)\}$ .

Therefore

$$\begin{aligned} t^i(c)[(t^i)^{-1}(t^i(c))] &= t^i(c)[C_{-i} \times \{\pi_i(c)\}] \\ &= m_i \pi_i(c)(C_{-i} \times \delta_{\pi_i(c)}(\pi_i(c))) \\ &= 1 \end{aligned}$$

□

Note that in Mertens and Zamir's approach, the universal type spaces are constructed by constructing first the universal BL-space  $Y$  and then taking taking  $T = t^i(Y)$ , while here we have shown how to construct belief spaces from the type spaces.

**Definition 7.10.** [MZ84] A *coherent beliefs hierarchy [over  $S$ ] of level  $K$*  ( $K = 1, 2, \dots$ ) is a sequence  $(C_0, C_1, \dots, C_K)$  where:

1.  $C_0$  is a compact subset of  $S$  and for  $k = 1, \dots, K$ ,  $C_k$  is a compact subset of  $C_{k-1} \times [\Delta(C_{k-1})]^N$  (as topological spaces). We denote by  $\rho_{k-1}$  and  $t^i$  the projections of  $C_k$  onto  $C_{k-1}$  and the  $i$ -th copy of  $\Delta(C_{k-1})$  respectively.

$$C_0 \xleftarrow{\rho_0} C_1 \xleftarrow{\rho_1} \dots \xleftarrow{\rho_{K-1}} C_K$$

2.

$$\rho_{k-1}(C_k) = C_{k-1}; k = 1, \dots, K$$

3. For all  $c_k \in C_k$ , let  $c_{k-1} = \rho_{k-1}(c_k)$ . Then for all  $i$ , and  $k = 2, \dots, K$ ,

**H1)** the marginal distribution of  $t^i(c_k)$  on  $C_{k-2}$  is  $t^i(c_{k-1})$ ;

**H2)** the marginal distribution of  $t^i(c_k)$  in the  $i$ -th copy of  $\Delta(C_{k-2})$  is the unit mass at

$$t^i(c_{k-1}) = t^i(\rho_{k-1}(c_k)).$$

The coherent hierarchies are used to build the universal beliefs space  $Y$ . They can be seen as the first  $K$  steps in the iteration that leads to the final sequence. The additional conditions we see come from different complications introduced in the construction. Part 2 of the definition states that the projections should be surjective. This condition is necessary here because the spaces  $C_k$  are compact subspaces of  $C_{k-1} \times (\Delta C_{k-1})^N$  and not that whole space.

Conditions **H1)** and **H2)** of part 3 have the following intuitive meaning:

**H1)** says that player  $i$ 's  $k$ -level beliefs coincide with his  $(k-1)$  level beliefs in whatever concerns hierarchies up to level  $(k-2)$ . Condition **H2)** says that player  $i$  knows his own previous order beliefs. [MZ84]

Under a more technical light, **H1**) can be written as

$$(\Delta\rho_{k-2})t^i(c_k) = t^i(c_{k-1}) = t^i(\rho_{k-1}(c_k)) \quad (7.9)$$

for every  $c_k \in C_k$ . This condition is saying that  $c_k$  is an element of the projective limit of the spaces  $C_k$ . The condition **H2**) can be written as: for every  $c_k \in C_k$ ,

$$(\Delta\rho_{k-2})t^i(c_k) = t^i(c_{k-1}) = t^i\rho_{k-1}(c_k). \quad (7.10)$$

There is some abuse of notation here: for each number  $k \geq 1$ , functions  $t^i : C_k \rightarrow \Delta C_{k-1}$  are defined, so there is a different function  $t^i$  that is applied to  $c_k$  and another one that's applied to  $c_{k-1}$ , and it should be clear which one is needed in each occurrence of  $t^i$ . Having (7.10) is needed in order to obtain (7.8) in the projective limit.

Morphisms between BL-spaces are defined as follows:

**Definition 7.11.** [MZ84] A *beliefs morphism* (BL-morphism) from a BL-space  $(C, S, f, (t^i)_{i=1}^N)$  to a BL-space  $(\tilde{C}, \tilde{S}, \tilde{f}, (\tilde{t}^i)_{i=1}^N)$  is a pair  $(\varphi, \varphi')$  where  $\varphi'$  is a continuous mapping from  $C$  to  $\tilde{C}$  and  $\varphi$  is a continuous mapping of  $S$  to  $\tilde{S}$  such that for each  $i; i = 1, 2, \dots, n$ , the following diagram commutes:

$$\begin{array}{ccc} S & \xrightarrow{\varphi} & \tilde{S} \\ f \uparrow & & \uparrow \tilde{f} \\ C & \xrightarrow{\varphi'} & \tilde{C} \\ t_i \downarrow & & \downarrow \tilde{t}^i \\ \Delta C & \xrightarrow{\Delta\varphi'} & \Delta\tilde{C} \end{array}$$

Given a fixed space  $S$  of states of nature, the *universal BL-space* is the final object in the category of BL-spaces over  $S$  and BL-morphisms. The universal BL-space over a fixed space  $S$  is built by



taking the projective limit  $Y$  of a sequence of coherent beliefs hierarchies:

Let  $Y_0 = S$ ;  $Y_1 = S \times \Delta S \times \dots \times \Delta S$  and for  $k \geq 2$ , let  $Y_k = \{y_k \in Y_{k-1} \times [\Delta(Y_{k-1})]^N : \mathbf{H1}\}$

For all  $i$  the marginal distribution of  $t^i(y_k)$  on  $Y_{k-2}$  is  $t^i(y_{k-1})$  and **H2** the marginal distribution of  $t^i(y_k)$  on  $\Delta^i(Y_{k-2})$  is the unit mass at  $t^i(y_{k-1})$ .

With this definition, for each value of  $k$ , the sequence  $(Y_0, \dots, Y_k)$  is a coherent beliefs hierarchy over  $S$  of level  $k$ , and it also is the biggest one that can be constructed. All the coherent hierarchies of beliefs can be mapped to the ones constructed above, and all the BL-spaces can be mapped in a unique way to their limit  $Y$ .

It is clear from the proof given that the spaces under consideration are assumed to be compact Hausdorff topological spaces. Mertens and Zamir use Riesz's Representation theorem to prove what essentially amounts to Theorem 3.4, but one needs to also assume that the probability measures involved are all regular, as [AB79] and [Hei93] point out.

## Heifetz and Samet

Aviad Heifetz and Dov Samet, in their paper *Topology-Free Typology of Beliefs*, [HS98], are the first to solve the problem of finding the universal type space in the general case of measurable spaces. They present two constructions of the space, much in the spirit of the two constructions of final coalgebras for measure polynomial functors presented in Chapters 4 and 5, respectively.

Their methods have provided us with guiding insight for the extensions we presented here. In the language-based construction, we have presented new languages  $\mathcal{L}(T)$  based on each measure polynomial functor  $T$ . We have also introduced two important refinements:

Their operator  $B_i^p(e)$  is used to express that a player  $i$  believes that an event represented by  $e$  has probability bigger than  $p$ . In our formulation, this would be expressed as  $[\mathbf{next}]_i \beta^p(e)$ , a formula of

sort  $Proj_i^N$  for the functor  $T$  from definition 7.3 that is, the syntactic operator  $B_i^p$  has been factored in two parts. This allows us to have more expressive power and describe points of coalgebras that are not of the form  $\Delta S(X)$  for some measure polynomial functor  $S$ .

Our refinement of their Lemma 3.5 to get Lemma 3.6 allows us to work with languages without negation, by proving that a  $\pi$ -system of generators is enough to generate the  $\sigma$ -algebra on  $\Delta X$ , and a boolean algebra is not necessary.

On section 5 of the paper, *The Universal Type Space in Terms of Hierarchies*, they describe types by their *histories*, and therefore the notation  $h$  we borrowed for the map  $h^c : X \rightarrow Z$ . Here the coalgebraic and categorial machinery we have used allowed us to simplify the construction, providing both more clarity and generality.

## Other related work

Among other work related to type spaces, we'd like to mention some in particular.

Spyros Vassilakis, in [Vas91], identifies the final sequence method as the right one to obtain a solution for  $X = \Delta(S \times X)$  in the category of Compact Hausdorff spaces. He also suggests further applications in [Vas90].

Brandenburger and Dekel in [BD93] propose a similar construction to that of [MZ84], and explore the relation of the concept of types with the one of common knowledge.

Luc Lismont in [Lis92] and Aviad Heifetz in [Hei96] present models of type spaces in the framework of non-wellfounded set theory, which has a close relation to the theory of coalgebras since its origin (see [Acz88] and [BM96]).

Probabilistic logic applied to type spaces has been studied by Heifetz and Mongin in [HM01], and Meier in [Mei01]. Meier also explored the simpler case of type spaces when the probabilities are

given by *finitely additive* measures in [Mei02].

This being just a cursory overview of the literature on this topic, it shows the interest in the problem, and also suggest directions for further development of both the applications and the general theory presented in previous chapters.

## Conclusions

In this dissertation we have tried to expand the usefulness of the theory of coalgebras, by studying them in a previously unexplored category, that of measurable spaces and functions. The main purpose on doing so, is to be able to express and model problems involving probability measures.

Most of the applications of coalgebras have been done in the category of sets, with a particular emphasis on theoretical computer science. Some examples of this include data structures and automata theory. But these notions have been generalized to include probabilities: in some systems the transitions are not deterministic, but they may follow some probability distribution.

Coalgebras provide a formal framework to think about processes, transitions between states are the essence of the structure maps  $c : X \rightarrow T(X)$ . The particular functor  $T$  used in the model determines what may be referred to as “observable behavior” of the system: some new information about an element  $x \in X$  is obtained after we apply the transition  $c$  to it. But this element  $c(x)$  will also make reference to new elements in  $X$ , the new state (or set of states, or probabilities about the states) in the system. To this new state we can apply  $c$  again to look further into the behavior of the element  $x$ . This idea leads naturally to the quest of somehow collecting all possible behaviors under a certain functor. Finding the final coalgebra for the functor achieves even more than this:

two elements in different coalgebras with the same history of behavior are mapped to the very same *element* in the final coalgebra.

We have presented two constructions of the final coalgebra for measure polynomial functors. These functors comprehend a wide class of useful functors for probabilistic systems. The purpose of giving two constructions is that they illuminate different aspects of the mathematical object under study (i.e. the final coalgebras for the measure polynomial functors).

Using the languages  $\mathcal{L}(T)$  from Chapter 4 provides a way of reasoning about the systems. Although an axiomatization of a logic has not been given yet, some of the literature points in that direction (for example, [Mos99], [HM01], [Mei01], [CP04]). One would like to have a deductive method for telling when a set of formulas is actually a realized theory of  $Id^*$  without appealing to a model. In other words, the goal would be to produce a complete logic for these languages.

Reasoning about probabilistic systems is specially appealing when we regard the probabilities as subjective beliefs as we've done in Chapter 7. We have used the category  $\mathbf{Meas}^N$  to model several agents having beliefs about the beliefs of each other, and we were able to model introspection (knowledge of one's own beliefs) in this setting. It will be interesting to take these insights to the category of sets and use them to create modal logics in which to study not only introspection but also the problem of common knowledge.

The second construction offered is simpler, possibly facilitating the use of these coalgebras for a wider audience not interested in logic. In this approach, the idea of describing behaviors is accomplished not by formulas but by trajectories a space that is easy to build just by iterating applications of the functor. Furthermore, the category-theoretic presentation helps clarifying the different assumptions made in the existing solutions for the problem of universal type spaces.

In both constructions the *ingredients* of the functor play an important role. For each one of

---

the ingredients  $S$  a space is constructed,  $S^*$  in the language driven construction and  $Z_S$  in the final sequence approach. We have kept in both approaches the same names for the measurable maps connecting the different spaces so to make the parallel between the constructions clear. The structure map of the final coalgebra is presented factored as a series of maps that have easy and natural definitions. That is not to say that proving they have the required properties is easy. Some technical difficulties need to be overcome, in particular for the map  $\epsilon$  going from  $(\Delta S)^*$  in Lemma 4.6 and from  $Z_{\Delta S}$  to  $\Delta Z_S$  in Lemma 5.5. In particular, it is in the proof of Lemma 4.6 that our Lemma 3.6 lets us do the construction without incorporating negations (they are not needed) to the languages  $\mathcal{L}(T)$ .

The general intuition on how the two given proofs work is that a given element in the final coalgebra must be the description of a particular element  $x$  in some coalgebra  $(X, c)$ . This element already has a prescribed target  $c(x)$ , and we can look for the description of  $c(x)$  in  $T^*$  or  $Z_T$ . Having one of the constructions does not automatically yield the other. By the nature of their underlying sets, the proofs of finality require different techniques.

We hope that the tools presented in this dissertation will find their applications both in computer science, economics and any other fields where spaces of probabilities may appear, and need to be reasoned upon.

# Bibliography

- [AB79] W. Armbruster and W. Böge. Bayesian game theory. In *Game theory and related topics (Proc. Sem., Bonn and Hagen, 1978)*, pages 17–28. North-Holland, Amsterdam, 1979.
- [Acz88] Peter Aczel. *Non-Well-Founded Sets*, volume 14 of *CSLI Lecture Notes*. CSLI Publications, Stanford, 1988.
- [AHS90] Jiří Adámek, Horst Herrlich, and George E. Strecker. *Abstract and concrete categories*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1990.
- The joy of cats, A Wiley-Interscience Publication.
- [AM89] Peter Aczel and Nax Mendler. A final coalgebra theorem. In *Category theory and computer science*, volume 389 of *Springer Lecture Notes in Computer Science*, pages 357–365. Springer, Berlin, 1989.
- [Aum61] Robert J. Aumann. Borel structures for function spaces. *Illinois Journal of Mathematics*, 5:614–630, 1961.
- [Aum64] Robert J. Aumann. Mixed and behavior strategies in infinite games. In *Advances in Game Theory*, volume 52 of *Annals of Mathematical Studies*. Princeton University Press, 1964.

- [BD93] Adam Brandenburger and Eddie Dekel. Hierarchies of beliefs and common knowledge. *J. Econom. Theory*, 59(1):189–198, 1993.
- [BDEP97] Richard Blute, Josée Desharnais, Abbas Edalat, and Prakash Panangaden. Bisimulation for labelled markov processes. In *Logic in Computer Science*, pages 149–158, 1997.
- [BdRV01] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal logic*, volume 53 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, Cambridge, 2001.
- [BE79] W. Böge and Th. Eisele. On solutions of Bayesian games. *Internat. J. Game Theory*, 8(4):193–215, 1979.
- [Bla51] J. H. Blau. The space of measures on a given set. *Fund. Math.*, 38:23–34, 1951.
- [BM96] Jon Barwise and Lawrence Moss. *Vicious Circles*, volume 60 of *CSLI Lecture Notes*. CSLI Publications, Stanford, 1996.
- [Bra04] Adam Brandenburger. The power of paradox: Some recent developments in interactive epistemology. Technical report, Stern School of Business, New York University, 2004.
- [BSdV03] Falk Bartels, Ana Sokolova, and Erik de Vink. A hierarchy of probabilistic system types. In H. Peter Gumm, editor, *Electronic Notes in Theoretical Computer Science*, volume 82. Elsevier, 2003.
- [CP04] Corina Cirstea and Dirk Pattinson. Modular construction of modal logics. In *Proceedings of Fifteenth International Conference on Concurrency Theory*, volume 3170 of *Lecture Notes in Computer Science*, pages 258–275. Springer, 2004.



- [DEP02] Josée Desharnais, Abbas Edalat, and Prakash Panangaden. Bisimulation for labelled Markov processes. *Inform. and Comput.*, 179(2):163–193, 2002. LICS’97 (Warsaw).
- [DGJP03] Josée Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. Approximating labelled Markov processes. *Inform. and Comput.*, 184(1):160–200, 2003.
- [Dud89] Richard M. Dudley. *Real analysis and probability*. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Pacific Grove, CA, 1989.
- [Dyn61] E. B. Dynkin. *Theory of Markov processes*. Translated from the Russian by D. E. Brown; edited by T. Köváry. Prentice-Hall Inc., Englewood Cliffs, N.J., 1961.
- [Gir82] Michèle Giry. A categorical approach to probability theory. In *Categorical aspects of topology and analysis*, volume 915 of *Springer Lecture Notes in Mathematics*, pages 68–85. Springer, Berlin-New York, 1982.
- [Gum99] H. Peter Gumm. Elements of the general theory of coalgebras. Technical report, LUATCS’99, Rand Africaans University, Johannesburg, South Africa, 1999.
- [Gum01] H. Peter Gumm. Functors for coalgebras. *Algebra Universalis*, 45(2-3):135–147, 2001. Conference on Lattices and Universal Algebra (Szeged, 1998).
- [Hal50] Paul R. Halmos. *Measure Theory*. D, Van Nostrand Company, Inc., New York, N. Y., 1950. xi + 304 pp.
- [Har67] John C. Harsanyi. Games with incomplete information played by “Bayesian” players. I. The basic model. *Management Sci.*, 14:159–182, 1967.
- [Har68a] John C. Harsanyi. Games with incomplete information played by “Bayesian” players. II. Bayesian equilibrium points. *Management Sci.*, 14:320–334, 1968.

- 
- [Har68b] John C. Harsanyi. Games with incomplete information played by “Bayesian” players. III. The basic probability distribution of the game. *Management Sci.*, 14:486–502, 1968.
- [Hei93] Aviad Heifetz. The Bayesian formulation of incomplete information- the non-compact case. *International Journal of Game Theory*, 21:329–338, 1993.
- [Hei96] Aviad Heifetz. Non-well-founded type spaces. *Games and Economic Behavior*, 16(2):202–217, 1996.
- [HM01] Aviad Heifetz and Phillippe Mongin. Probability logic for type spaces. *Games and Economic Behavior*, 35:31–53, 2001.
- [HS98] Aviad Heifetz and Dov Samet. Topology-free typology of beliefs. *Journal of Economic Theory*, 82(2):324–341, 1998.
- [Jac01] Bart Jacobs. Many-sorted coalgebraic modal logic: a model-theoretic study. *Theor. Inform. Appl.*, 35(1):31–59, 2001. Coalgebraic methods in computer science (Berlin, 2000).
- [Kec95] A. S. Kechris. *Classical Descriptive Set Theory*. Springer-Verlag, New York, 1995.
- [KP02] Alexander Kurz and Dirk Pattinson. Definability, canonical models, compactness for finitary coalgebraic modal logic. In *CMCS 2002: coalgebraic methods in computer science (Berlin)*, volume 65 of *Electron. Notes Theor. Comput. Sci.*, page 21 pp. (electronic). Elsevier, Amsterdam, 2002.
- [Kur01] Alexander Kurz. Specifying coalgebras with modal logic. *Theoret. Comput. Sci.*, 260(1-2):119–138, 2001. Coalgebraic methods in computer science (Lisbon, 1998).

- [Lam68] Joachim Lambek. A fixpoint theorem for complete categories. *Math. Z.*, 103:151–161, 1968.
- [Lis92] Luc Lismont. Une classe universelle anti-fondée pour les croyances. *C. R. Acad. Sci. Paris Sr. I Math.*, 315(11):1117–1120, 1992.
- [LS91] Kim G. Larsen and Arne Skou. Bisimulation through probabilistic testing. *Inform. and Comput.*, 94(1):1–28, 1991.
- [Mei01] Martin Meier. An infinitary probability logic for type spaces. Technical report, CORE, Université Catholique de Louvain, Belgium, 2001.
- [Mei02] Martin Meier. Finitely additive beliefs and universal type spaces. Technical report, Center for operations research and econometrics, Université catholique de Louvain, 2002.
- [Mèt63] Michel Métivier. Limites projectives de mesures. martingales. applications. *Ann. Mat. Pura Appl. (4)*, 63:225–352, 1963.
- [Mos99] Lawrence S. Moss. Colagebraic logic. *Annals of Pure and Applied Logic*, 96(1-3):277–317, 1999.
- [MV04] Lawrence S. Moss and Ignacio D. Viglizzo. Harsanyi type spaces and final coalgebras constructed from satisfied theories. *Electronic Notes in Theoretical Computer Science*, 106:279–295, 2004.
- [MV05] Lawrence S. Moss and Ignacio D. Viglizzo. Final coalgebras for functors on measurable spaces. *Information and Computation*, 2005. Accepted, under revision.
- [MZ84] J. F. Mertens and S. Zamir. Formulation of Bayesian analysis for games of incomplete information. *International Journal of Game Theory*, 14:1–29, 1984.

- [OR94] Martin J. Osborne and Ariel Rubinstein. *A course in game theory*. MIT Press, Cambridge, MA, 1994.
- [Par67] K. R. Parthasarathy. *Probability measures on metric spaces*, volume 3 of *Probability and Mathematical Statistics*. Academic Press, Inc., New York-London, 1967.
- [RdV99] J. J. M. M. Rutten and E. P. de Vink. Bisimulation for probabilistic transition systems: a coalgebraic approach. *Theoretical Computer Science*, 211:271–293, 1999.
- [Röß99] Martin Rößiger. Languages for coalgebras on datafunctors. In *CMCS'99 Coalgebraic Methods in Computer Science (Amsterdam, 1999)*, volume 19 of *Electron. Notes Theor. Comput. Sci.*, page 22 pp. (electronic). Elsevier, Amsterdam, 1999.
- [Röß00] Martin Rößiger. *Coalgebras, Clone Theory, and Modal Logic*. PhD thesis, Technischen Universität Dresden, 2000.
- [Röß01] Martin Rößiger. From modal logic to terminal coalgebras. *Theoret. Comput. Sci.*, 260(1-2):209–228, 2001. Coalgebraic methods in computer science (Lisbon, 1998).
- [Rut00] J. J. M. M. Rutten. Universal coalgebra: a theory of systems. *Theoretical Computer Science*, 249(1):3–80, 2000.
- [SP82] M. B. Smyth and G. D. Plotkin. The category-theoretic solution of recursive domain equations. *SIAM J. Comput.*, 11(4):761–783, 1982.
- [TR98] Daniele Turi and Jan Rutten. On the foundations of final coalgebra semantics: non-well-founded sets, partial orders, metric spaces. *Math. Structures Comput. Sci.*, 8(5):481–540, 1998.
- [Vas90] Spyros Vassilakis. Rules for changing the rules. Technical report, Stanford - Institute for Thoretical Economics, 1990.

- 
- [Vas91] S. Vassilakis. Functorial fixed points: a non-technical introduction. Working Paper 266, Department of Economics, University of Pittsburg, 1991.
- [vBMOW03] Franck van Breugel, Michael Mislove, Joël Ouaknine, and James Worrell. An intrinsic characterization of approximate probabilistic bisimilarity. In *Foundations of software science and computation structures*, volume 2620 of *Lecture Notes in Comput. Sci.*, pages 200–215. Springer, Berlin, 2003.
- [vBMOW05] Franck van Breugel, Michael Mislove, Joël Ouaknine, and James Worrell. Domain theory, testing and simulation for labelled Markov processes. *Theoret. Comput. Sci.*, 333(1-2):171–197, 2005.
- [vBSW02] Franck van Breugel, Steven Shalit, and James Worrell. Testing labelled Markov processes. In *Automata, languages and programming*, volume 2380 of *Lecture Notes in Comput. Sci.*, pages 537–548. Springer, Berlin, 2002.
- [Vig05] Ignacio D. Viglizzo. Final sequences and final colagebras for measurable spaces. Accepted in CALCO 05, 2005.
- [Wor99] James Worrell. Terminal sequences for accessible endofunctors. In *CMCS'99 Coalgebraic Methods in Computer Science (Amsterdam, 1999)*, volume 19 of *Electron. Notes Theor. Comput. Sci.*, page 15 pp. (electronic). Elsevier, Amsterdam, 1999.

# Curriculum Vitae

Ignacio Darío Viglizzo

Department of Mathematics, Indiana University, Bloomington

Contact Information: (812)857-6787 / igvigliz@indiana.edu

## Education

- 2005. PhD (Mathematics) Indiana University, Bloomington, USA. Advisor: Lawrence S. Moss.
- 2005. Logic Certificate. Logic Program, Indiana University, Bloomington, USA.
- 1999. Magister en Matemática, Universidad Nacional del Sur, Bahía Blanca, Argentina. Advisor: Luiz F. Monteiro.
- 1994. Licenciado en Matemática, Universidad Nacional del Sur, Bahía Blanca, Argentina.

## Teaching Experience

- August 1999-July 2005. Associate Instructor. Department of Mathematics, Indiana University, Bloomington.
- July 1998-August 1999. Teaching Assistant. Department of Mathematics, Universidad Nacional del Sur, Argentina.
- March 1993-March 1998. Junior Teaching Assistant. Department of Mathematics, Universidad Nacional del Sur, Argentina.

## Professional Service

- Local Student co-chair for NASSLLI 2003, Indiana University, Bloomington.

## Honors

- Rothrock Teaching Award for Associate Instructors, Department of Mathematics, Indiana University, April 2004.
- James P. Williams Memorial Award, Department of Mathematics, Indiana University, April 2000.
- Fulbright Fellowship. 1999-2001.
- Fellowship from CONICET (National Research Council, Argentina). 1995-1999.
- “25 de Mayo” Award, (Best GPA among graduates in Exact, Physical and Natural Sciences), Universidad Nacional del Sur, June 1995.

### Articles in Journals and Conference Proceedings

- I. Viglizzo. *Final sequences and final coalgebras for measurable spaces*. To appear in the proceedings of CALCO (Conference on Algebra and Coalgebra in Computer Science) 2005, LNCS.
- L. Moss and I. Viglizzo. *Final Coalgebras for functors on measurable spaces*. To appear in *Information and Computation*.
- L. Moss and I. Viglizzo. *Harsanyi Type Spaces and Final Coalgebras Constructed from Satisfied Theories*. Electronic Notes in Theoretical Computer Science, Volume 106, Proceedings of the Workshop on Coalgebraic Methods in Computer Science (CMCS) Pages 279-295.
- I. Viglizzo. *Free Monadic Three-valued Lukasiewicz Algebras*. Revista de la Unión Matemática Argentina, Volumen 41,2,1998, pp. 109-117.