



Italics indicate geographic areas in Indiana where unit names are used.
*Simplified to illustrate common lithologies.
[†]These units normally occur only in the subsurface. However, some units of Ordovician and Lower Silurian age are exposed at the Kentland and Hammer structures in northwestern Indiana.

Note: Mixtures and interbedding of lithologies can occur.

REFERENCES

¹ Gray, H. H., Ault, C. H., and Keller, S. J., 1987. Bedrock geologic map of Indiana: Indiana Geological Survey Miscellaneous Map 48, scale 1:500,000.
² Gradstein, F. M., Ogg, J. G., Smith, A. G., Agterberg, F. P., and others, 2004. A geological time scale 2004: Cambridge, U.K., Cambridge University Press, 384 p.
³ Rupp, J. A., 1991. Structure and isopach maps of Paleozoic rocks of Indiana: Indiana Geological Survey Special Report 48, 106 p.
⁴ Shaver, R. H., Ault, C. H., Burger, A. M., Carr, D. D., and others, 1986. Compendium of Paleozoic rock-unit stratigraphy in Indiana—a revision: Indiana Geological Survey Bulletin 59, 203 p.
⁵ Hasenmueller, W. A., Hasenmueller, N. R., and Daniels, M. S., 2013. Indiana Geologic Names Information System: Indiana Geological Survey website, <http://igs.indiana.edu/IGNIS/Search.cfm>, date accessed, July 1, 2013.

- sandstone, conglomerate
- shale, mudstone, siltstone
- limestone
- dolomite
- coal
- volcanic rocks
- igneous rocks
- rocks absent
- unconformity
- bioturbation
- chert