

Vitamin B₁₂ regulates photosystem gene expression via the CrtJ antirepressor AerR in *Rhodobacter capsulatus*

Zhuo Cheng,¹ Keran Li,¹ Loubna A. Hammad,² Jonathan A. Karty² and Carl E. Bauer^{1*}

Departments of ¹Molecular and Cellular Biochemistry and ²Chemistry, Indiana University, Bloomington, IN 47405, USA.

Summary

The tetrapyrroles haem, bacteriochlorophyll and cobalamin (B₁₂) exhibit a complex interrelationship regarding their synthesis. In this study, we demonstrate that AerR functions as an antirepressor of the tetrapyrrole regulator CrtJ. We show that purified AerR contains B₁₂ that is bound to a conserved histidine (His145) in AerR. The interaction of AerR to CrtJ was further demonstrated *in vitro* by pull down experiments using AerR as bait and quantified using micro-scale thermophoresis. DNase I DNA footprint assays show that AerR containing B₁₂ inhibits CrtJ binding to the *bchC* promoter. We further show that *bchC* expression is greatly repressed in a B₁₂ auxotroph of *Rhodobacter capsulatus* and that B₁₂ regulation of gene expression is mediated by AerR's ability to function as an antirepressor of CrtJ. This study thus provides a mechanism for how the essential tetrapyrrole, cobalamin controls the synthesis of bacteriochlorophyll, an essential component of the photosystem.

Introduction

Cobalamin (vitamin B₁₂) is one of the most complex non-polymeric biomolecules synthesized by cells. B₁₂ is important cofactor of many enzymes, such as methyltransferases, reductases and isomerases, which catalyse a range of biological processes (Gruber *et al.*, 2011). There are two forms of B₁₂ that are commonly used *in vivo*, adenosylcobalamin (AdoB₁₂) and methylcobalamin (MeB₁₂) that typically have 5'-deoxyadenosyl or a methyl group coordinated as an upper axial ligand to a corrin bound cobalt respectively. The upper carbon-cobalt bond is relatively weak, generating a reactive leaving group with a homolytic bond disassociating energy of ~30 kcal mol⁻¹

(Halpern, 1985; Waddington and Finke, 1993). Both AdoB₁₂ and MeB₁₂ undergo spontaneous homolytic cleavage of the carbon-cobalt bond resulting in the generation of hydroxocobalamin (OHCB₁₂) when photoexcited at room temperature in aqueous solutions (Weissbach *et al.*, 1960; Hogenkamp, 1966).

Only certain members of bacteria and archaea synthesize B₁₂ *de novo* (Perlman, 1959). The metabolically diverse photosynthetic α -proteobacterium *Rhodobacter capsulatus* is capable of synthesizing B₁₂ under both aerobic and anaerobic conditions (McGoldrick *et al.*, 2002). In this species haem, bacteriochlorophyll (Bchl) and B₁₂ biosynthesis share common early intermediates from δ -aminolevulinic acid (ALA) to uroporphyrinogen III at which point the B₁₂ pathway branches off from the haem/Bchl pathways. The haem and Bchl pathways further diverge at protoporphyrin IX, where iron is chelated to form haem or magnesium is chelated to form the first dedicated intermediate in the Bchl pathway.

Among photosynthetic organisms such as *R. capsulatus*, the levels of haem and B₁₂ stay relatively stable while the amount of Bchl is altered dramatically in response to different environmental conditions. Under aerobic conditions only trace amounts of Bchl are synthesized as expression of genes coding for enzymes for the Bchl branch are repressed by the aerobic repressor CrtJ (Ponnampalam *et al.*, 1995; Ponnampalam and Bauer, 1997). Under anaerobic conditions the amount of Bchl is significantly higher as *bch* genes are no longer repressed by CrtJ, and are additionally anaerobically activated by the RegB/RegA two-component system (Du *et al.*, 1998; Bird *et al.*, 1999; Willett *et al.*, 2007).

CrtJ is a major regulator of genes coding for enzymes involved in the biosynthesis of haem, Bchl and carotenoids, as well as structural proteins of light harvesting-II complex (Penfold and Pemberton, 1991; 1994; Ponnampalam *et al.*, 1995). Previous studies have demonstrated that CrtJ binds promoters containing two conserved palindromes (TGTN₁₂ACA) (Ponnampalam *et al.*, 1995; Ponnampalam and Bauer, 1997), and that redox regulated binding is dependent on oxidation of a conserved cysteine (C420) located in the helix–turn–helix DNA-binding motif (Masuda *et al.*, 2002; Cheng *et al.*, 2012). In many proteobacteria, *crtJ* homologues are immediately preceded by a gene (termed *aerR* in *R. capsulatus*) that encodes a regulator of photosynthesis gene expression

Accepted 8 December, 2013. *For correspondence. E-mail bauer@indiana.edu; Tel. (+1) 812 855 6595; Fax (+1) 812 856 5710.




Fig. 1. Diagram showing the conserved chromosomal positioning of *aerR* homologues upstream of *crtJ* homologues in various species. Blue arrows represent homologues of *aerR*; orange arrows represent homologues of *crtJ*.

(Fig. 1). *AerR* does not have any identifiable DNA-binding domains so it has been assumed that it may control gene expression by affecting the DNA binding activity of another protein with *CrtJ* being the most likely candidate (Gomelsky *et al.*, 2003; Masuda *et al.*, 2008). Interestingly, *AerR* has significant sequence similarity to B_{12} -binding domains suggesting that it may be capable of interacting with B_{12} (Ludwig and Matthews, 1997; Gomelsky *et al.*, 2003). However, the tetrapyrrole specificity of *AerR* is somewhat controversial as there is a report that a maltose-binding domain fused to a truncated variant of a homologue from *Rhodobacter sphaeroides* preferentially binds haem over B_{12} (Moskvin *et al.*, 2010).

In addition to the role of B_{12} as an enzyme cofactor, this complex tetrapyrrole is also sometimes involved in regulating gene expression. For example, B_{12} is a ligand for RNA-based riboswitches that typically regulate synthesis of enzymes involved in B_{12} biosynthesis (Nou and Kadner, 2000; Mandal and Breaker, 2004). In *Myxococcus xanthus* CarH functions as a B_{12} -dependent repressor of carotenogenic genes (Ortiz-Guerrero *et al.*, 2011). Ado B_{12} containing CarH binds an operator as a tetramer to repress *crt* gene expression in the dark. Photolysis of Ado B_{12} to OH B_{12} disassembles the tetramer and causes dissociation of CarH from the operator (Ortiz-Guerrero *et al.*, 2011). The involvement of cobalamin in regulating photosystem genes from *R. capsulatus* was initially reported by Pollich *et al.* (Pollich *et al.*, 1993; Pollich and Klug, 1995). These studies demonstrated that light harvesting and reaction centre photosystem gene expression were significantly impaired in a B_{12} auxotroph. Normal photosystem gene expression could be recovered by exogenous addition of B_{12} ; however, the mechanism whereby B_{12} regulates expression of photosystem genes was not established.

In this study, we demonstrate that the *aerR* open reading frame located immediately upstream of *crtJ* codes

for a B_{12} binding antirepressor of *CrtJ*. This study provides the first example where B_{12} regulates gene expression by controlling the interaction of an anti-repressor with a transcription repressor. It also provides a molecular mechanism for the control of photosystem gene expression based on the availability of B_{12} .

Results

AerR is required for photosystem synthesis

A chromosomal deletion of the *aerR* gene was generated by removing its entire coding sequence with the exception of retention of the start codon (the correct start codon was determined as described in *Supporting information*) (Fig. S4) that remained in frame with the stop codon. The Δ *aerR* strain exhibits a much lighter pigmentation in comparison to wild-type *R. capsulatus* and also grows significantly slower under photosynthetic conditions. Spectral analyses of the Δ *aerR* cell lysate revealed that photosystem synthesis is reduced ~2.5-fold in this strain when grown under anaerobic photosynthetic conditions (Fig. 2A). QRT-PCR analysis of *bchC* expression, which is a *CrtJ* regulated gene that codes for an enzyme in the bacteriochlorophyll biosynthetic pathway, is also consistent with spectral analyses (Fig. 2B). Specifically, we observed that the Δ *aerR* strain exhibits a ~18-fold reduction in anaerobic *bchC* expression over that observed with wild-type cells (Fig. 2B).

aerR is located immediately upstream of *crtJ* in *R. capsulatus* as well as in nearly all sequenced species of purple photosynthetic bacteria that contain *CrtJ* (Fig. 1). To address whether removal of *aerR* could cause a polar effect on downstream *crtJ* expression, a FLAG-epitope-tagged chromosomal version of *crtJ* was introduced in a Δ *aerR* strain as described previously (Dong *et al.*, 2002). Δ *aerR*/Flag-*crtJ* and SB1003/Flag-*crtJ* were grown to 100 Klett units under anaerobic photosynthetic conditions and cell extracts were subjected to Western blot analysis with monoclonal antibody to the FLAG epitope tag in *CrtJ*. Levels of *CrtJ* protein in the Δ *aerR* and wild-type parental strains are comparable indicating that deletion of *aerR* did not affect the cellular level of *CrtJ* (Fig. S1). Finally, we also complemented the Δ *aerR* strain by introducing a plasmid-born *aerR* gene *in trans*. As shown in Fig. 2A, the complemented strain exhibits nearly normal amount of photosystem synthesis. QRT-PCR results also indicate that the addition of *AerR in trans* also restored expression of *bchC* to normal levels (Fig. 2B). We conclude, therefore, that the reduction of photopigment synthesis observed by the Δ *aerR* strain is caused by the absence of *AerR* and not due to a polar effect on the downstream expression of the *crtJ* gene.




Fig. 2. AerR activates photosystem synthesis.

A. Spectral scans of crude cell extracts from anaerobically grown wild-type parent strain (SB1003) (black), the *aerR* deletion strain ($\Delta aerR$) (green) and the *aerR* deletion strain complemented with a plasmid encoded copy of *aerR* (red).
B. Quantitative real-time QRT-PCR showing the relative expression level of *bchC* in the *aerR* deletion strain ($\Delta aerR$), and $\Delta aerR$ complementation strain ($\Delta aerR + aerR$) as compared to the wild-type SB1003 strain (defined as a transcript level of 1) under photosynthetic growth condition. Error bars represent standard error of the mean ($n = 3$).

AerR is a B₁₂-binding protein

Analysis of the primary amino acid sequence of AerR shows the presence of a putative cobalamin binding site Asp/Glu-X-His-X-X-Gly-(41)-Ser(Thr)-X-Leu-(26–28)-Gly-Gly (Fig. S2A) (Ludwig and Matthews, 1997). To test whether AerR is capable of binding B₁₂, various forms of B₁₂ were added to purified AerR and after 5 min incubation, unbound B₁₂ was then removed by passing the mixture through a desalting column. Spectral analysis of the eluted protein (Fig. 3A) shows that purified AerR exhibits no observable interaction with AdoB₁₂ or with MeB₁₂. Interestingly, exposure of the AerR and AdoB₁₂ mixture to white light, which promotes upper ligand photolysis to generate OHB₁₂, leads to binding of B₁₂ (Fig. 3A). This is confirmed by the direct addition of OHB₁₂ to purified apo-AerR which subsequently binds stoichiometric amounts of OHB₁₂ even under dark condi-




Fig. 3. AerR binding to OHB₁₂.

A. Light converts AdoB₁₂ to OHB₁₂ and enables AerR association. Dash line is a spectral scan of AerR incubated with AdoB₁₂ in dark for 5 min followed by removal of unbound AdoB₁₂; dotted line is a spectral scan of AerR incubated with MeB₁₂ in dark for 5 min followed by removal of unbound MeB₁₂; Solid line is a spectral scan of AerR incubated with AdoB₁₂ under illumination ($\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of white light) for 5 min followed by removal of unbound AdoB₁₂.
B. Spectral scan of AerR incubated with OHB₁₂ in the dark for 5 min followed by removal of unbound OHB₁₂.
C. Spectral scan of WT AerR and AerR H145A mutant protein purified from *E. coli* with AdoB₁₂ added to the growth medium and lysis buffer with the lysate exposed to $\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of white light for 5 min prior to protein purification. Solid line is WT AerR; dotted line is AerR H145A.
Proteins used for spectral analysis were SUMO tagged.

tions (Fig. 3B). Finally, tight binding of OHB₁₂ to AerR can also be observed by the addition of 10 μ M AdoB₁₂ to the cell lysate of an *Escherichia coli* strain that overexpressed AerR followed by exposure to white light. Spectral analysis of AerR isolated from the light excited lysate exhibits an absorption spectrum that reflects excellent binding of light-generated OHB₁₂ (Fig. 3C, solid line).

In well-studied B₁₂-dependent enzymes such as methionine synthase and methylmalonyl-coenzyme A mutase, a conserved histidine in the B₁₂-binding domain displaces the lower dimethylbenzimidazole ligand to form a histidine lower axial ligand with the cobalt atom of B₁₂ (Drennan *et al.*, 1994). An alignment of the B₁₂-binding domain from methionine synthase to four AerR homologues shows the presence of a conserved histidine (His145) in AerR that aligns with the lower axial cobalt linkage histidine of methionine synthase (Fig. S2). To test if His145 is involved in binding of OHB₁₂, we constructed an alanine substitution at this position and assayed whether this mutant protein could bind B₁₂ as discussed above. Spectral analysis shows that purified AerR H145A protein indeed does not bind any detectable amounts of B₁₂ (Fig. 3C, dotted line).

We further characterized B₁₂ binding to AerR by performing liquid chromatography electrospray ionization mass spectrometry (LC-ESI-MS) on the AerR:B₁₂ complex. ESI-MS has been widely used to study protein-ligand complexes (Loo, 1997) with an example of methionine synthase-B₁₂ complex successfully detected by ESI-MS (Drummond *et al.*, 1993). The deconvoluted mass spectrum generated by MassLynx software of purified AerR:B₁₂ complex treated with 6 M urea gives two distinct species (Fig. 4A) with masses of 26892 ± 2 Da and 28221 ± 2 Da. The mass of the smaller species matches very well with the expected 26893.2 Da mass of apo-AerR as calculated by its protein sequence plus a sodium atom. The mass difference between the two species is 1329 ± 1 Da, which is consistent with the mass of the B₁₂ prosthetic group (1329.4 Da). When 0.04 M HCl was added to the purified AerR:B₁₂ complex, the LC-ESI-MS spectrum still showed the same two species (Fig. 4B). The AerR:B₁₂ bound form at mass 28221 Da is still present, though less intense following the HCl treatment than in the urea-treated sample. This indicates that B₁₂ remains attached to a portion of the protein even under strong acidic conditions and suggests that the interaction between B₁₂ and AerR is stronger than the expected non-covalent axial interaction between cobalt and H145.

To address which amino acid residue in AerR is forming an upper ligand with B₁₂, we utilized liquid chromatography electrospray tandem ionization mass spectrometric analysis (LC-ESI-MS/MS). We first digested purified AerR:B₁₂ complex with pepsin in the presence of 0.04 M HCl overnight at 37°C and then analysed the resulting peptides with LC-ESI-MS/MS. We did not find any peptides containing

H145 that also had an attached B₁₂ but did observe a peptide corresponding to residues 6–16 (VELEHGST-GCL) of AerR that exhibited a UV-Vis spectrum consistent with an attached B₁₂ (Fig. 4E inset). The CID spectrum of this peptide yielded fragment ions (Fig. 4G) that are consistent with a B₁₂ containing peptide and with the fragment ion assignments in Fig. 4H. These results suggest that an amino acid residue in this segment of AerR (residues 6–16) forms the tight upper axial cobalt ligand with B₁₂ (Fig. 4E–G). Since there are two amino acid residues in this peptide that could potentially form an upper axial ligand to cobalt (His10 and Cys15), we mutated each of these two residues to alanine either individually or in combination and then analysed these mutants for peptides containing B₁₂ using LC-ESI-MS/MS. The C15A mutant protein still yielded a B₁₂-containing peptide as shown in Fig. S3 while the H10A mutant and the H10A-C15A double mutant proteins did not yield any B₁₂ containing peptides. These results suggest that His10 is the upper axial cobalt-coordinating residue present above the corrin plane and that it forms a stronger bond to cobalt than does the proposed lower axial His145 bond.

We also used LC-ESI-MS analysis to confirm that the proposed lower linkage between His145 and cobalt is weaker than the proposed upper linkage to His10. For this analysis we observed that LC-ESI-MS analysis of full-length AerR H10A:B₁₂ gave rise to two species after treatment with 6 M urea: a higher mass (28154 ± 2 Da) that is consistent with that of AerR H10A containing an attached B₁₂ plus a sodium atom (28156.6 Da) and a lower mass (26826 ± 2 Da) that is of AerR H10A without bound B₁₂ plus a sodium atom (26827.2 Da) (Fig. 4C). However treatment of AerR H10A:B₁₂ protein with 0.04 M HCl, which would protonate the ϵ -nitrogen of His145 and result in it losing its coordination to cobalt, gave rise only to apo-H10A without bound B₁₂ (26826 ± 2 Da) (Fig. 4D). This is distinctly different from what was observed with wild-type AerR:B₁₂ that has both proposed upper and lower cobalt coordinating His145 and His10 and is still capable of retaining bound B₁₂ even under acidic conditions (Fig. 4B). These results confirm that both His10 and His145 are coordinating to cobalt and that the proposed lower axial bond between cobalt and His145 is weaker than that of the proposed upper axial bond between cobalt and His10.

AerR is an antirepressor of the tetrapyrrole regulator CrtJ

We undertook several independent studies to ascertain whether AerR can interact with CrtJ. In one assay, we performed *in vitro* pull down experiments using resin bound SUMO-AerR containing B₁₂ as 'bait' for the binding of tag-less purified CrtJ. SUMO-AerR:B₁₂ was loaded on the Ni²⁺ column followed by extensive washing to remove




Fig. 4. Mass spectrometric analyses of AerR and AerR mutant proteins.

A–D. The deconvoluted spectrum of full-length protein LC-ESI-MS spectrum of (A) 6 M urea treated WT AerR protein; (B) 0.04 M HCl-treated WT AerR protein; (C) 6 M urea treated AerR H10A protein; (D) 0.04 M HCl-treated AerR H10A protein.

E–G. LC-ESI-MS/MS analyses of a B₁₂-containing peptide (residues 6–16: VELEHGSTGCL) obtained by pepsin digestion. (E) Extracted ion chromatogram (EIC) of the triply charged peptide containing B₁₂ at m/z 824.7 [the asterisk (*) indicates the peak of interest]; the corresponding UV-Vis spectrum of this peptide is shown in the inset. (F) High-resolution mass spectrum of this B₁₂ containing peptide ion at m/z 824.7 (2 ppm mass error). (G) Collision induced dissociation (CID) spectrum of this peptide ion at m/z 824.7. (E) and (G) were obtained using the Bruker HCT ion trap MS instrument whereas (F) was obtained using the Thermo LTQ-Orbitrap MS instrument.

H. Structure and CID fragmentation of the pepsin-generated WT AerR peptide containing B₁₂ attached to His10. The experimental data for this peptide are shown in Fig. 4E–G.




Fig. 5. CrtJ and AerR:B₁₂ form a complex *in vitro*.

A. CrtJ-AerR pull down experiments. Top panel is SUMO-AerR:B₁₂ used as 'bait' to pull down tag-less CrtJ. Bottom panel is tag-less CrtJ used as a control without SUMO-AerR:B₁₂. The imidazole concentration of buffer used in each step was indicated above the picture.

B. Absorbance traces of CrtJ (blue line), AerR:B₁₂ (grey line) and mixture of CrtJ and AerR:B₁₂ (red line), respectively, that were subjected to Superdex 200 chromatography. SDS-PAGE analysis of the middle red peak at volume 15 is shown above the Superdex 200 chromatography absorbance trace.

C. Representative CrtJ-AerR:B₁₂ binding isotherm generated by microscale thermophoresis assay. Continuous line through binding isotherm represents a non-linear regression analysis fitted to a Hill equation (Eq. 1).

unbound SUMO-AerR:B₁₂. Purified tag-less CrtJ was then loaded onto the column followed by extensive washing and then SUMO-AerR:B₁₂ was eluted with a gradient of imidazole. Figure 5A showed that CrtJ co-eluted with SUMO-AerR at an imidazole concentration of 150 mM. As a control, we observed that addition of just tag-less CrtJ to

the Ni²⁺ column resulted in CrtJ eluting at a lower imidazole concentration of 60 mM.

We also performed Superdex 200 size exclusion chromatography to determine whether AerR and CrtJ form a stable complex that co-elutes. The elution profile of CrtJ alone showed that it chromatographs as a single peak with an apparent molecular weight (MW) of 127.3 ± 4.0 kDa which is close to the size of a CrtJ dimer that has a calculated MW of 101.0 kDa (Fig. 5B blue line). The elution profile of AerR:B₁₂ alone also shows a single major peak with a calculated MW of 27.3 ± 4.0 kDa which is close to the size of an AerR monomer containing B₁₂ that has a calculated MW of 28.2 kDa (Fig. 5B grey line). After incubating CrtJ with AerR:B₁₂ (~20 μ M respectively) at 22°C for 1 h, the protein mixture was subjected to chromatography with a resulting elution profile containing three peaks (Fig. 5B red line). One peak corresponds to the same size of a CrtJ dimer, a second peak corresponds to the AerR monomer with a third unique peak present with a calculated MW of 73.0 ± 4.3 kDa. This size corresponds well to the calculated MW of 78.7 kDa for a 1:1 CrtJ-AerR:B₁₂ complex. SDS-PAGE analysis of the peak corresponding to the AerR-CrtJ complex fraction is shown above the absorbance trace. The CrtJ and AerR protein bands have similar intensity further indicating that CrtJ binds to AerR:B₁₂ with a 1:1 stoichiometry.

We also measured the binding affinity for the interaction of CrtJ and AerR:B₁₂, by performing microscale thermophoresis analysis (Monolith NT.115, Nanotemper Technologies GmbH). Microscale thermophoresis measures a change in a protein hydration shell caused by a change in protein conformation or by the formation of a stable interaction between two proteins (Wienken *et al.*, 2010). The binding affinity as measured using replicates of this procedure was calculated to be $EC_{50} = 3.3 \pm 0.7$ μ M (Fig. 5C).

The effect of AerR:B₁₂ binding to CrtJ on the ability of CrtJ to bind operator DNA was addressed by undertaking DNase I DNA footprint assay of CrtJ binding to a pair of CrtJ recognition sequences present in the *bchC* promoter (Fig. 6). Footprint analysis of CrtJ alone shows excellent protection of the tandem CrtJ binding sites (TG₁₂ACA) that overlaps the -35 and -10 promoter recognition sequences. However, when CrtJ was pre-incubated with AerR:B₁₂ at 22°C for 1 h and then used for footprint analysis, these two protection sites disappeared. As a control, AerR:B₁₂ alone does not show any protection on the *bchC* promoter. These results indicate that AerR containing bound B₁₂ effectively inhibits the ability of CrtJ to bind to its target sequence.

AerR senses B₁₂ in vivo

We addressed whether the presence of cobalamin has an effect on *in vivo* photosystem gene expression by con-




Fig. 6. AerR:B₁₂ inhibits CrtJ binding to the *bchC* promoter. DNase I footprint analysis of CrtJ binding to the *bchC* promoter region in the absence of AerR:B₁₂ (third panel), in the presence of AerR:B₁₂ (fourth panel). The first panel is the DNase I digestion pattern of the *bchC* promoter without adding any protein and the second panel is AerR:B₁₂ alone with the *bchC* promoter, which used as a control. CrtJ binding sites are indicated with dashed boxes.

structuring a $\Delta cobN$ B₁₂ auxotroph strain that contains a kanamycin resistance gene cassette inserted into the *cobN* gene that encodes the N subunit of cobalt chelatase. Cells were grown under photosynthetic conditions in the presence of 10 μ M exogenous B₁₂ or under B₁₂ limiting condition. Expression of the bacteriochlorophyll biosynthesis *bchC* gene was then assayed using QRT-PCR as *bchC* is known to be repressed by CrtJ (Ponnampalam *et al.*, 1995; 1998). The QRT-PCR results shown in Fig. 7A demonstrate that *bchC* gene expression is lowered (approximately threefold) when B₁₂ is limiting

and that this effect is reversed by the addition of 10 μ M exogenous B₁₂. Similar studies were carried out in a $\Delta cobN \Delta aerR$ double-deletion strain. In this strain *bchC* expression remains low compared to WT strain (~13-fold lower) with or without exogenous B₁₂ in the growth medium. This indicates that the $\Delta cobN \Delta aerR$ strain can no longer control *bchC* gene expression *in vivo* in response to B₁₂ availability.

To test the effect of disrupting the putative upper and lower cobalt binding ligands on AerR's activity *in vivo* we recombined *aerR*-H10A, *aerR*-H145A and *aerR* H10A




Fig. 7. AerR senses B_{12} in vivo.

A. QRT-PCR showing the relative expression level of *bchC* in the *cobN* disruption strain ($\Delta cobN$) and the *cobN aerR* double-deletion strain ($\Delta cobN \Delta aerR$) under photosynthetic condition in the presence of exogenous 10 μ M B_{12} or under B_{12} limited conditions, as compared with wild-type SB1003 strain (defined as a transcript level of 1).

B. Spectral scans of crude cell extracts from anaerobically grown wild-type parent strain (SB1003) (black), the *aerR* deletion strain ($\Delta aerR$) (green), the *aerR* H10A mutant strain (blue), the *aerR* H145A mutant strain (purple) and the *aerR* H10A H145A double mutant strain (pink).

H145A mutation into the cognate position in the chromosome and assayed for photosystem synthesis. As shown by the spectral analysis in Fig. 7B, the *aerR* H10A (blue spectrum) and H145A strain (purple spectrum) exhibits photosystem production that is intermediate from that of the *AerR* deletion strain (green spectrum) and the wild-type parent strain (black spectrum). Interestingly, photosystem production of a strain that contains a H10A H145A double mutation (pink spectrum) is quite similar to the *aerR* deletion strain (green spectrum), indicating that both of the two His residues play a role in the control of *AerR*'s function.

Finally, we tested if there is an effect of disrupting the *AerR* lower ligand to OHB_{12} on the light sensing

capabilities of these cells. As shown in Fig. 8A, growth of wild-type *R. capsulatus* under high light intensity ($\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) leads to an $\sim 50\%$ reduction in pigment synthesis relative to the amount of pigments synthesized when growth under low light ($\sim 5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) conditions. In contrast, growth of H145A mutant cells under similar high and low light conditions (Fig. 8B) results in cells that contain the same intermediate amounts of pigment production at levels that are similar to that observed in wild-type cells under high light growth conditions. The implications of this phenotype are discussed below.

Discussion

The involvement of B_{12} in the control of photosystem synthesis has been a long observed feature of photosynthetic microorganisms. Nearly five decades ago, Pfennig and Lippert reported the isolation of several strains of purple bacteria that required B_{12} for growth (Pfennig and Lippert, 1966). These strains exhibited severely reduced photosystem synthesis under B_{12} limiting conditions. Decades later this observation was followed by studies by




Fig. 8. Spectral scans of crude cell extracts from photosynthetically grown cells under different light intensities. A. Wild-type strain SB1003 grown under high light (red line) with $\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of white light and low light (blue line) with $\sim 5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ of white light. B. *aerR* H145A mutant strain under high (red line) and low (blue line) light intensity.




Fig. 9. Models for the light-responsive B₁₂-based regulatory switches in *R. capsulatus* and *M. xanthus*.

A. Light converts AdoB₁₂ to OHB₁₂ which enables OHB₁₂ binding to AerR and thus binding to CrtJ and derepresses photosynthesis gene expression in *R. capsulatus*.

B. Light mediated photolysis of the AdoB₁₂::CarH tetramer to the OHB₁₂::CarH monomer allows *M. xanthus* to synthesize carotenoids in the light (Ortiz-Guerrero *et al.*, 2011).

Pollich and Klug which demonstrated that photosystem gene expression is six- to sevenfold lower in a *R. capsulatus* B₁₂ auxotroph than in the wild-type strain (Pollich *et al.*, 1993; Pollich and Klug, 1995). A molecular mechanism for B₁₂ dependence of photosystem gene expression has, however, remained unclear.

A major finding of our study is that AerR functions as a B₁₂-sensing antirepressor of CrtJ. The ability of AerR to bind B₁₂, and subsequently interact with CrtJ, is clearly shown by pull down experiments using AerR:B₁₂ as 'bait', by co-elution of AerR:B₁₂ with CrtJ during size exclusion chromatography, and finally by the determination of the AerR binding affinity to CrtJ (Fig. 5). DNase I DNA footprint assays also demonstrate that AerR:B₁₂ inhibits the binding of CrtJ to the *bchC* promoter which clearly shows that AerR is functioning as an antirepressor of CrtJ (Fig. 6). The *in vitro* B₁₂ binding and antirepressor activity of AerR is also supported by *in vivo* assays which show that a $\Delta cobN$ B₁₂ auxotroph has threefold reduction in *bchC* expression under B₁₂ limiting conditions with resumption of normal *bchC* expression occurring after addition of 10 μ M B₁₂ only when there is functional AerR (Fig. 7A). These results suggest that AerR indeed controls the DNA binding activity of CrtJ in response to B₁₂ *in vivo*, which provides the first molecular understanding as to why B₁₂ is required for

optimal photosystem gene expression in this species. Furthermore, all sequenced strains of purple photosynthetic bacteria that contain a *crtJ* homologue also contain a homologue of *aerR* (Fig. 1). This indicates that B₁₂ control of photosystem gene expression via AerR is most likely a conserved feature of this group of organisms.

Until recently, B₁₂ was considered to be a simple metabolite used as a cofactor for enzyme reactions. However, B₁₂ is also known to be involved in controlling expression of genes coding for enzymes involved in B₁₂ biosynthesis by functioning as a metabolite that is sensed by RNA-based riboswitches that are present upstream of a number of *cob* genes (Nahvi *et al.*, 2002; Rodionov *et al.*, 2003). Most recently it has been observed that *M. xanthus* also contains a DNA binding transcription factor called CarH that utilizes AdoB₁₂ as a cofactor to regulate carotenoid gene expression. CarH containing bound AdoB₁₂ forms a stable tetramer that represses carotenoid gene expression in the dark. Exposure of *M. xanthus* to light promotes photolysis of AdoB₁₂ to OHB₁₂ which causes disassembly of the CarH tetramer and subsequent dissociation of CarH from the carotenoid promoters (Ortiz-Guerrero *et al.*, 2011). This light mediated photolysis of the AdoB₁₂::CarH tetramer to the OHB₁₂::CarH monomer subsequently allows *M. xanthus* to synthesize carotenoids in the light (Fig. 9B).

In the case of *R. capsulatus*, AerR also appears to utilize B₁₂ as a light sensor at some level as it only binds OHB₁₂ that would presumably be elevated under light excitation growth conditions (Fig. 9A). Interestingly the H145A mutant does seem to lack the ability to regulate pigment production in response to alterations in light intensity (Fig. 8). However, this mutant only synthesizes intermediate amount of pigments which is not dim light pigment levels that one would expect if it were only defective in sensing OHB₁₂ production (Fig. 8). Clearly additional studies will have to be undertaken to determine what role, if any, that wild-type AerR has on adjusting the amount of pigment production in response to alterations in light intensity and whether such a response is due to light driven changes in OHB₁₂ production.

CrtJ is also known to contain a redox regulated cysteine in the helix–turn–helix DNA-binding domain that when oxidized promotes an increase in DNA binding activity. Under anaerobic growth conditions this Cys is present in a reduced thiol form that reduces CrtJ DNA binding affinity by ~20-fold (Cheng *et al.*, 2012). Mutagenesis of the redox active Cys to Ala leads to constitutive derepression while mutating the redox active Cys to Ser leads to constitutive repression (Cheng *et al.*, 2012). The phenotype of the *aerR* deletion strain is somewhat surprising in that it has severely reduced *bchC* expression even in the absence of oxygen. This suggests that CrtJ is capable of repressing target gene expression under reducing conditions in strains where *aerR* is deleted. It has also been shown that the CrtJ homologue from *R. sphaeroides*, PpsR, binds haem and regulates expression of haem biosynthesis genes in response to haem availability (Yin *et al.*, 2012). Preliminary studies with *R. capsulatus* CrtJ also indicates that it is also capable of binding haem in a manner similar to that described for PpsR indicating that haem is a likely input that controls CrtJ's activity as is described for PpsR (Z. Cheng and C.E. Bauer, unpubl. obs.). Thus, the regulation of pigment production by CrtJ is highly complex involving an interplay of numerous inputs; redox, potentially haem, and now B₁₂ via AerR. Furthermore, two of these inputs, haem and B₁₂ availability involve end-products of a biosynthetic pathway that is regulated by CrtJ. Determining the mechanism(s) whereby CrtJ responds to these very different inputs remains a distinct challenge going forward.

The similar phenotypes of the *aerR* H145A and H10A mutant strains is also surprising as gene expression and photosystem synthesis of these two mutant are intermediate between that of wild-type cells and that of the *aerR* deletion strain (Fig. 7B). The phenotype of the H145A point mutation is also not as severe as observed with the B₁₂ auxotroph suggesting that even though this point mutation is not capable of loading B₁₂ *in vitro*, it is not a null mutation *in vivo*. Either this mutation is still capable of

binding some form of B₁₂ *in vivo* or it suggests that apo-AerR may also have a role in controlling gene expression. Unfortunately, apo-AerR is very unstable *in vitro*, which has hampered our ability to purify enough protein without B₁₂ to undertake biochemical analysis of its function, if any, in affecting the DNA binding activity of CrtJ.

Sequence alignment of AerR homologues from numerous purple bacteria shows that there is significant homology with the B₁₂-binding domain from *E. coli* methionine synthase as well as to the presumptive B₁₂ binding region of CarH from *M. xanthus* (Fig. S2A). In free B₁₂, the 5,6-dimethylbenzimidazole base forms an axial ligand to cobalt at the lower side of the corrin macrocycle. However when bound to methionine synthase, the 5,6-dimethylbenzimidazole unbinds and swings away to allow histidine to form a lower axial ligand to cobalt. This swapping of the lower ligand is called a base-off His-on conformation (Drennan *et al.*, 1994). All characterized B₁₂-dependent methyltransferases have the conserved B₁₂ binding sequence Asp/Glu-X-His-X-X-Gly-(41)-Ser(Thr)-X-Leu-(26–28)-Gly-Gly which is also present in AerR (Fig. S2A) (Ludwig and Matthews, 1997). This suggests that B₁₂ bound to AerR also has the dimethylbenzimidazole base-off His-on conformation and that His145 may be forming a lower axial ligand with bound cobalt. This lower ligand is also supported by modelling an AerR structure based on the methyltransferases structure that shows the proper positioning of His145 (Fig. S2B), and by our mutational studies which show that a His145 to Ala mutation is incapable of binding B₁₂ *in vitro*.

In methyltransferases the upper cobalt ligand is a methyl group that functions as a leaving group for methylation of substrates (Gruber *et al.*, 2011). AerR does not bind MeB₁₂ and instead binds OHB₁₂ that has a bound water as an upper ligand. Mass spectrometric (MS) analysis shows that AerR tightly binds OHB₁₂ presumably by displacement of the upper OH-cobalt linkage present in OHB₁₂ with His10. This upper axial ligand is extremely tight and indeed remains bound during MS analyses even under conditions where the protein is denatured and digested by proteases. Mammalian B₁₂ transporter transcobalamin is also an example of a B₁₂-binding protein that uses a histidine to form an upper ligand to corrin bound cobalt (Wuerges *et al.*, 2006). The binding of the OHB₁₂ to transporter transcobalamin also includes the displacement of water with a histidine residue (Fedosov *et al.*, 2000). However, this B₁₂ transporter differs in that it is highly selective for the base-on form of B₁₂ which presumably facilitates its release upon transport. The overall structure of transporter transcobalamin, and the orientation of the bound B₁₂ is also quite different from other B₁₂ binding enzymes that have the classic B₁₂ binding sequence as observed with AerR, CarH and methyltransferases (Gruber *et al.*, 2011). AerR from *R. capsulatus* is neither an enzyme nor a B₁₂ trans-

porter so it is interesting that it combines the ligand binding features of these two different B₁₂-binding proteins.

AerR and its homologues from other species also have considerable homology to the SCHIC domain of a well-characterized antirepressor from *Rhodobacter sphaeroides* called AppA (Gomelsky *et al.*, 2003; Moskvina *et al.*, 2007). Along the same lines as AerR, AppA is known to inhibit the DNA binding activity of PpsR, the CrtJ homologue in *R. sphaeroides* (Masuda and Bauer, 2002; Winkler *et al.*, 2013). Interestingly, the AppA SCHIC domain preferentially binds haem instead of B₁₂ as occurs with AerR (Han *et al.*, 2007; Moskvina *et al.*, 2007; Yin *et al.*, 2013). The recent crystal structure of the AppA SCHIC domain, however, highlights several important differences between the putative haem binding pocket in the SCHIC domain and the B₁₂ binding pocket present in the methionine synthase structure (Drennan *et al.*, 1994; Winkler *et al.*, 2013; Yin *et al.*, 2013). Specifically, the His that forms a lower axial ligand to haem in AppA is offset two residues in AppA relative to the location of the lower ligand His present in methyltransferase and for AerR (Yin *et al.*, 2013). In addition, methionine synthase has a space in the core of the MetH:B₁₂ Rossmann fold that anchors the bulky lower nucleotide tail that attached to B₁₂ (Drennan *et al.*, 1994). In the SCHIC domain, the access to this docking space is blocked by Glu289 (Yin *et al.*, 2013). In contrast to the haem binding SCHIC domain, AerR and its homologues have retained extensive homology to region of MetH that allows insertion of the lower nucleotide tail of B₁₂ into the Rossmann fold. Presumably, this difference is promoting tetrapyrrole specificity differences between AerR and AppA.

Given that AerR is a B₁₂-binding protein, it is also surprising that a previous study on an AerR homologue from *R. sphaeroides* called PpaA was shown to bind haem *in vitro* (Moskvina *et al.*, 2010). We have repeatedly attempted to observe haem binding by apo-AerR without success, which is contrasted by the ease at which AerR is capable of binding OHB₁₂. A major difference between our studies and that of Moskvina *et al.* is that we have analysed the tetrapyrrole binding activity of full-length tag-less AerR while the Gomelsky study utilized a truncated variant of PpaA that only contained homology to the SCHIC domain of AppA. The analysed truncated variant of PpaA was further fused to a maltose-binding domain. Ongoing analysis of tetrapyrrole binding by PpaA and other AerR homologues by our laboratory indicates that tetrapyrrole specificity of this class of B₁₂-binding proteins involves residues outside of the SCHIC region of homology. This region was missing in the prior study with PpaA (A.J. Vermeulen and C.E. Bauer, unpublished).

In conclusion, this study, in conjunction with the recent CarH study from *M. xanthus*, indicates that B₁₂ has an important role in controlling gene expression in several

species by modulating the DNA binding activity of transcription factors. This occurs either directly as is the case of CarH or indirectly with an antirepressor as is the case of AerR and CrtJ (Fig. 9). Proteins that contain B₁₂-binding domain similar to AerR or CarH are present in a number of sequenced bacterial genomes (Ortiz-Guerrero *et al.*, 2011), so it is quite possible that this newly discovered form of regulating gene expression may not be an uncommon mechanism.

Experimental procedures

Strains, media and growth conditions

In vivo studies were undertaken with *R. capsulatus* SB1003. *E. coli* strains DH5 α and DH10 β were used for cloning. *E. coli* TEC5 (Taylor *et al.*, 1983) was used in conjugation to introduce various plasmids to *R. capsulatus*. *E. coli* S17-1 λ pir (Miller and Mekalanos, 1988) was used to facilitate the transfer of suicide plasmids into *R. capsulatus*. AerR overexpression was carried out in *E. coli* BL21 (DE3). Various *R. capsulatus* strains were grown in PY salts liquid medium as described previously (Cheng *et al.*, 2012).

Construction of aerR mutations

An *aerR* in-frame deletion was constructed by amplifying ~500 base pairs upstream of *aerR* and ~500 downstream of *aerR* with PCR, using the primers shown in Table S1 (aerRupSacl-F and aerRupEcoRV-R; aerRdownEcoRV-F and aerRdownXbaI-R). The upstream ~500 bp fragment was cloned into SacI and EcoRV sites while the downstream ~500 bp fragment was cloned into EcoRV and XbaI sites of pBluescript SK+ (Stratagene). This generated a ~1 kb fragment that contains the upstream and downstream sequences of *aerR* with the only remaining *aerR* coding sequence ATG-GATATCTGA (underlined sequence introduced an EcoRV restriction site). The SacI-XbaI fragment was subsequently subcloned into corresponding sites in suicide plasmid pZJD29a (Masuda and Bauer, 2004), generating pZJD29a- Δ aerR, and then conjugated into *R. capsulatus* SB1003. pZJD29a- Δ aerR genome recombinants were selected for the presence of gentamicin resistance and sucrose sensitivity. Cells carrying the in-frame deletion of *aerR* gene were further verified by colony-PCR. The strain carrying *aerR* in-frame deletion was designated as Δ aerR.

A chromosomally located *aerR* H145A point mutation was generated by constructing a ~1.6 kb fragment covering the *aerR* gene and both the upstream and downstream ~500 bp region by PCR amplification with primers aerRupSacl-F and aerRdownXbaI-R (Table S1). The PCR product was digested with SacI and XbaI and inserted into pBluescript SK+ (Stratagene). This plasmid was then used as a template for generating a point mutation via PCR mutagenesis as using a Stratagene QuikChange kit with Turbo Pfu DNA polymerase. A pair of primers aerRH145A-f and aerRH145A-r (Table S1) was subsequently used to introduce the point mutation. The mutation was confirmed by sequencing before subcloning into pZJD29a. pZJD29a-aerRH145A was transformed into

S17-1 λ pir and subsequently conjugated into the $\Delta aerR$ strain. Procedures to select for first and second recombination events were the same as used in generating the $\Delta aerR$ strain. The presence of the mutated version of *aerR* gene in the genome was confirmed by PCR amplification of the *aerR* region followed by sequence analysis of the PCR segment. *aerR* H10A point mutant and *aerR* H10A H145A double mutant strain were generated using the same procedure as used for *aerR* H145A mutant strain (primers used are shown in Table S1).

AerR overexpression and purification

A 750 bp fragment of *aerR* gene was PCR amplified using primers *aerRSUMOBbsl-f* and *aerRSUMOBamHI-r* (Table S1). The PCR product was cloned into pCR4Blunt-TOPO plasmid as described using an Invitrogen Zero Blunt TOPO PCR Cloning Kit. After the cloned fragment was checked by sequencing analysis it was then digested with restriction enzymes *Bbsl* and *BamHI* followed by subcloning into a SUMOpro vector plasmid (Lifesensors). The resulting ORF encodes AerR with an N-terminal His₆-tag and a SUMO-tag, which enhances solubility of AerR. The resulting plasmid pSUMO-AerR was then transformed into BL21 (DE3) for protein overexpression.

To overexpress SUMO tagged AerR, *E. coli* strain BL21 (DE3)/pSUMO-AerR was grown in LB medium at 37°C in the presence of 25 $\mu\text{g ml}^{-1}$ kanamycin to an OD₆₀₀ of 0.7–0.8. Cultures were cooled to 16°C and then 50 μM of isopropyl- β -D-thiogalactopyranoside (IPTG) and 1 μM AdoB₁₂ were added with AerR expression continued at 16°C for 16–20 h. Cells were harvested by centrifugation and resuspended in a buffer containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 5 mM imidazole, 10% glycerol and 10 μM AdoB₁₂. Cells were then lysed by passing the cell suspension through a chilled French Press cell three times at 18 000 psi. The lysate was then clarified by centrifugation at 30 000 *g* for 30 min at 4°C with the resulting supernatant illuminated with white light at intensity of $\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ for 5 min. The supernatant was then passed through a 45 μm membrane filter (Millipore) and loaded onto a 1 ml HisTrap column (GE) using an ÄKTA chromatography system (GE), washed with 20-column volume of wash buffer containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 20 mM imidazole, and 10% glycerol. SUMO-AerR was then eluted with a gradient of 20 mM imidazole to 500 mM imidazole in the loading buffer over 15 column volumes total. Eluted protein was then digested with SUMO protease Ulp1 in presence of 1 mM DTT at 22°C for 2 h, followed by a desalting column (Bio-Red, Econo-Pac® 10DG Column) against wash buffer. SUMO tag was then removed from the protein solution by passing through 1 ml Ni Sepharose high-performance resin (GE Healthcare) in an Econo-Pac chromatography column (Bio-Rad) and wash buffer was used to wash the residual protein on the resin. Tag-less AerR was further purified by Superose 12 (S-12) a 16 \times 700 mm size exclusion chromatography in 20 mM Tris-HCl (pH 8.0) and 200 mM NaCl at a flow rate of 1 ml min⁻¹. Peak fractions were analysed by SDS-PAGE with AerR containing fractions collected and pooled. AerR stayed as a monomer in solution, as calculated from its size exclusion chromatography profile.

pSUMO-AerR plasmid was used as a template to generate *aerR* point mutations (primers used are shown in Table S1) and mutant proteins were purified using the same procedure as that for WT AerR.

Complementation analysis

A ~ 1.2 kb fragment containing the *aerR* gene as well as 600 bp upstream was PCR amplified with primers *aerRupSacl-F* and *aerRSUMOBamHI-r* (Table S1). This fragment was cloned into pCR4Blunt-TOPO plasmid (Invitrogen) and then cleaved with *HindIII* and *EcoRV* restriction sites located on pCR4Blunt-TOPO plasmid. The *HindIII*-*EcoRV* fragment was then inserted into pBBR-MCS2 that was pre-digested with the same enzymes. Plasmid pBBR-MCS2-*aerR* was then transformed into S17-1 λ pir and then conjugated into the $\Delta aerR$ strain.

Spectral analysis

Rhodobacter capsulatus parent strain SB1003 and the mutant strains $\Delta aerR$ and *aerRH145A* were grown photosynthetically either under low light intensity ($\sim 5 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), normal light intensity ($\sim 30 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$), or under high light intensity ($\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) to a density of 75 klett units in PY salts medium. Ten millilitres of cultures were chilled on ice and then harvested by centrifugation at 4400 *g* for 10 min at 4°C. Cell pellets were resuspended in 1 ml of buffer containing 20 mM Tris-HCl pH 8.0, NaCl 150 mM and lysed by sonication $10 \text{ s} \times 3$. The cell lysate was clarified by centrifugation at 15 600 *g* for 10 min. The protein concentration of each supernatant was determined using the Bradford assay (Advanced Protein Assay Reagent, Cytoskeleton) with BSA as a standard. Each sample was measured three times and the average concentration was obtained. Absorption spectrum was scanned from 400 nm to 900 nm on a DU640 Beckman spectrophotometer with each supernatant adjusted by its individual protein concentration.

For analysis of B₁₂ binding, purified WT and H145 AerR protein were scanned on a DU640 Beckman spectrophotometer from 300 nm to 700 nm. The AdoB₁₂, MeB₁₂ and OHB₁₂ binding experiments were performed with SUMO tagged AerR as tag-less AerR is unstable without bound B₁₂. Apo SUMO tagged protein was purified using the same procedure as used for SUMO-AerR:B₁₂, except that no AdoB₁₂ was added to the LB medium and lysis buffer and the supernatant was not illuminated. The absence of B₁₂ in the purified SUMO-AerR was confirmed by spectral analysis. SUMO-AerR was incubated with different B₁₂ in the buffer containing 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 20 mM imidazole, and 10% glycerol in the dark or with white light for 5 min (light intensity $\sim 1000 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$) before passing through a 13 \times 26 mm PD MidiTrap G-25 desalting chromatography column (GE). Eluted samples were scanned for spectral absorption.

Mass spectrometry

Purified wild-type and mutant AerR proteins containing bound B₁₂ ($\sim 20 \mu\text{M}$) were treated with either 6 M urea or 0.04 M HCl for 15 min and then used for liquid chromatography electro-

spray ionization mass spectrometry (LC-ESI-MS) analysis, using a Waters/Micromass LCT Classic time of flight (Waters Corp., Milford, MA) mass spectrometer with a Waters CapLC inlet. Five microlitres of sample was loaded onto a 0.5 × 50 mm Thermo BioBasic C₈ reverse-phase column (300 Å pore size, 5 µm particles size, Thermo Scientific, West Palm Beach, FL) in Solvent A (5% acetonitrile, 95% water, 0.1% formic acid) and analysed in positive ion mode. The elution was carried out in a 20 min linear gradient from 10% solvent B to 90% Solvent B (95% acetonitrile, 5% water, 0.1% formic acid) at 15 µl min⁻¹. UV absorption at 216 nm and mass spectra from *m/z* 600 to 2300 were collected and analysed using MassLynx software.

Purified wild-type and mutant AerR proteins (~20 µM) were digested with either 20-fold (weight ratio) pepsin (Promega) or chymotrypsin (Roche Applied Science) overnight at 37°C for pepsin or 22°C for chymotrypsin. The resulting samples were then analysed by LC-ESI-MS/MS analysis using the LTQ Orbitrap instrument. LC-ESI-MS/MS analysis was performed using an Eksigent nano LC system (AB SCIEX, Farmingham, MA) interfaced to an externally calibrated LTQ Orbitrap hybrid mass spectrometer equipped with a nanospray ion source (Thermo Scientific, San Jose, CA). A 4 µl aliquot of the peptide digest dissolved in 20 µl of mobile phase A was washed with mobile phase A for 5 min at a flow rate of 5 µl min⁻¹ using an in-house packed reversed-phase C18 (Michrom Magic C18, 200 Å pore size, 5 µm particle size, Bruker-Michrom, Fremont, CA) trap column made from a 15 mm × 100 µm Integragrit column (New Objective, Woburn, MA). The desalted peptides were then separated using an in-house packed reversed-phase C18 (Magic C18, 100 Å, 5 µm particle size, Bruker-Michrom) column (150 mm × 75 µm, 15 µm tip) (New Objective, Woburn, MA) and analysed in positive ion mode. Mobile phase A consisted of 0.1% formic acid in acetonitrile:water (3%:97%, volume:volume) and mobile phase B consisted of 0.1% formic acid in acetonitrile:water (97%:3%, volume:volume). The flow rate was 300 nl min⁻¹ with the following gradient: 3% B from 0 to 1 min, ramp to 10% B from 1 to 9 min, ramp to 30% B from 9 to 28 min, ramp to 40% B from 28 to 31 min, ramp to 80% B from 31 to 35 min, hold at 90% B from 36 to 43 min, ramp to 3% B from 43 to 44 min, then re-equilibrate the column at 3% B for 14 min (58 min total run time). The mass spectrometer was operated in an automated data-dependent mode alternating between an MS scan in the Orbitrap and five collision-induced dissociation (CID) scans in the linear ion trap. The Orbitrap monitored precursor ions from *m/z* 300 to *m/z* 2000 at 15000 resolving power. The five most intense precursor ions in each scan were sequentially isolated in the linear trap with a 2 *m/z* isolation width and were collisionally activated at 35% normalized collision energy. The cycle was continuously repeated throughout the entire separation with the dynamic exclusion set to 45 s with a repeat count of 2. The Orbitrap provided high mass accuracy (< 5 ppm) for the precursor ions while the CID scan in the linear ion trap provided nominal mass information (± 0.25 *m/z*) for the tandem mass spectra. B₁₂ containing peptides were found by searching the tandem mass spectra for B₁₂ diagnostic fragment ions as shown in Fig. 4E–G and Fig. S3. The presence of B₁₂ was further confirmed by coincidence of the B₁₂ UV absorption peak at 360 nm with the parent *m/z* values and diagnostic fragment ions of peptides containing B₁₂ using liquid

chromatography-UV/Vis spectroscopy-tandem mass spectrometry (LC-UV/Vis-MS/MS). The chromatographic separation was performed with an Agilent 1200 capillary liquid pump (Agilent Technologies, Santa Clara, CA) equipped with a micro well-plate autosampler and a diode array UV-VIS absorbance detector that had a 500 nl flow cell. The column was 0.3 mm i.d., 150 mm long and packed with 5 µm Zorbax SBC18-300 particles (Agilent). Eight microlitres of sample was injected with the peptides separated at 4 µl min⁻¹ using the same gradient conditions as detailed above. The UV-VIS detector recorded absorption spectra from 250 to 750 nm at 2.5 Hz with individual chromatograms at 216 and 360 nm recorded. Tandem mass spectra of the peptides were recorded in manual MS² mode with the instrument set to obtain CID spectra for specific *m/z* values corresponding to the B₁₂ containing peptides previously observed using the Orbitrap. A Bruker HCT-Ultra PTM ion trap mass spectrometer (Bruker Daltonics, Billerica, MA) after the diode array detector used 1.0 V activation energy on the endcaps for fragmentation and recorded data from *m/z* 400 to 1800 in UltraScan mode (50 s delay measured between UV and MS).

RNA isolation, quantitative real-time PCR

Rhodobacter capsulatus wild-type and mutant strains were grown as described above for spectral analysis. One millilitre of cell cultures were harvested with total RNA of each sample extracted using the ISOLATE RNA Kit (Bioline). RNA samples were then treated with TURBO DNase (Ambion) with final OD₂₆₀ to OD₂₈₀ ratio of RNA samples (approximately 2.0) measured using NanoDrop spectrophotometer (Thermo Scientific). Quantitative real-time (QRT-PCR) analysis of gene expression utilized SensiFAST™ SYBR Hi-ROX One-Step Kit (Bioline) with 8 ng of RNA sample used in a 20 µl reaction. cDNA synthesis was at 45°C for 10 min followed by polymerase activation at 95°C for 2 min. A two-step PCR reaction was used, consisting of product melting at 95°C for 5 s, followed by primer annealing and DNA polymerase elongation at 60°C for 20 s. As an internal control, the house-keeping gene *rpoZ* that encodes DNA-directed RNA polymerase omega subunit was used. All primers used in QRT-PCR were as previously described (Cheng *et al.*, 2012).

CrtJ–AerR interaction studies

In vitro pull down experiments were performed with SUMO-AerR that was purified as describe above using tag-less CrtJ as purified previously (Cheng *et al.*, 2012). One millilitre Ni Sepharose high-performance resin (GE Healthcare) was loaded onto an Econo-Pac chromatography column (Bio-Rad) and equilibrated with wash buffer [20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 20 mM imidazole and 10% glycerol]. SUMO-AerR protein was loaded onto the Ni Sepharose column as 'bait' then the column was washed with 20 ml of wash buffer at 22°C. The purified tag-less CrtJ was then loaded onto the column and subjected to a gradient imidazole step elution using a buffer comprised of 20 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% glycerol with increasing imidazole concentrations. Tag-less CrtJ was used as a control

without the loading of SUMO-AerR. Protein composition in each fraction was visualized by SDS-PAGE.

An AerR–CrtJ interaction was also probed using co-elution size exclusion chromatography on a column that was calibrated using commercial gel filtration standards (Bio-Rad). Purified CrtJ (~20 µM) and/or AerR (~20 µM) were incubated (molar ratio 1:1) at 22°C for 1 h before loading on a pre-packed Superdex 200 column (10 × 300 mm) that was equilibrated. The column was equilibrated and eluted with 20 mM Tris-HCl (pH 8.0), 150 mM NaCl at 0.5 ml min⁻¹.

Microscale thermophoresis assays

Purified CrtJ (20 µM) was labelled using Monolith NT™ Protein Labelling Kit RED-NHS (Nano Temper Technologies). 500 nM labelled CrtJ was incubated with different concentration of AerR in a buffer containing 20 Tris-HCl (pH 8.0), 200 mM NaCl for 1 h at 22°C. The reactions were loaded on Monolith NT.115™ Standard Treated Capillaries (Nano Temper Technologies) and analysed with a Monolith NT.115 Blue Red microscale thermophoresis (MST) instrument. Thermophoresis with no jump was used to plot the binding isotherm. The resulting data were fitted to a Hill equation (Eq. 1) with OriginPro 8.6.

$$\log \frac{F - F_C}{F_{CB} - F} = n \cdot \log[A] - \log K_{app} \quad (1)$$

Parameters of the fit included normalized fluorescence intensities of free and bound CrtJ protein (F_C and F_{CB}). Hill coefficient (n) and the apparent dissociation constants (K_{app}) were both obtained. $K_{app} = (EC_{50})^n$; A : AerR concentration. EC_{50} corresponds to the AerR concentration when 50% of CrtJ are bound to AerR.

DNase I DNA footprint assay

The *bchC* promoter region (325 bp) was amplified using a 5-HEX labelled forward primer and a 6-FAM-labelled reverse primer (Table S1). After DNA agarose gel electrophoresis the PCR product was purified using the QIAquick Gel extraction kit (Qiagen) and used as DNA probe. The concentration of the DNA probe was determined spectrophotometrically, using the appropriate extinction coefficient. 10 nM DNA probe was incubated with different concentrations of AerR or CrtJ for 30 min at 22°C in a buffer containing 25 mM Tris-HCl (pH 8.0), 200 mM NaCl, 2 mM MgCl₂ and 500 µg ml⁻¹ heparin (250-fold weight excess over probe). For reactions incubated with CrtJ and AerR co-complex, CrtJ and AerR were mixed and incubated for 1 h at 22°C before adding to the probe. The final concentrations for CrtJ and AerR were at 100 nM and 1 µM respectively. The reaction was set up in 20 µl volume to which 3 µl of 125 U ml⁻¹ DNase I (New England BioLabs) was added and incubated for 15 min at 22°C to digest DNA. Twenty-three microlitres of 0.5 M EDTA (pH 8.0) was added to stop the reaction. DNA digestion product was recovered using Min Elute PCR purification kit (Qiagen), with sample eluted with 15 µl EB buffer from the kit plus 1 µl of 500 LIZ™ size standards (Applied Biosystems). The eluted samples were separated and detected with a 3730 DNA analyser

(Applied Biosystems) and the results analysed using Peak Scanner v1.0 (Applied Biosystems).

Acknowledgements

This study was supported by an NIH Grant R37 GM040941 awarded to C.E.B. The METACyt Biochemical Analysis Center was supported by a grant from the Eli Lilly Endowment.

References

- Bird, T.H., Du, S., and Bauer, C.E. (1999) Autophosphorylation, phosphotransfer, and DNA-binding properties of the RegB/RegA two-component regulatory system in *Rhodobacter capsulatus*. *J Biol Chem* **274**: 16343–16348.
- Cheng, Z., Wu, J., Setterdahl, A., Reddie, K., Carroll, K., Hammad, L.A., et al. (2012) Activity of the tetrapyrrole regulator CrtJ is controlled by oxidation of a redox active cysteine located in the DNA binding domain. *Mol Microbiol* **85**: 734–746.
- Dong, C., Elsen, S., Swem, L.R., and Bauer, C.E. (2002) AerR, a second aerobic repressor of photosynthesis gene expression in *Rhodobacter capsulatus*. *J Bacteriol* **184**: 2805–2814.
- Drennan, C.L., Huang, S., Drummond, J.T., Matthews, R.G., and Lidwig, M.L. (1994) How a protein binds B₁₂: a 3.0 Å X-ray structure of B₁₂-binding domains of methionine synthase. *Science* **266**: 1669–1674.
- Drummond, J.T., Loo, R.R., and Matthews, R.G. (1993) Electrospray mass spectrometric analysis of the domains of a large enzyme: observation of the occupied cobalamin-binding domain and redefinition of the carboxyl terminus of methionine synthase. *Biochemistry* **32**: 9282–9289.
- Du, S., Bird, T.H., and Bauer, C.E. (1998) DNA binding characteristics of RegA. A constitutively active anaerobic activator of photosynthesis gene expression in *Rhodobacter capsulatus*. *J Biol Chem* **273**: 18509–18513.
- Fedosov, S.N., Fedosova, N.U., Nexø, E., and Petersen, T.E. (2000) Conformational changes of transcobalamin induced by aquocobalamin binding. Mechanism of substitution of the cobalt-coordinated group in the bound ligand. *J Biol Chem* **275**: 11791–11798.
- Gomelsky, L., Sram, J., Moskvina, O.V., Horne, I.M., Dodd, H.N., Pemberton, J.M., et al. (2003) Identification and *in vivo* characterization of PpaA, a regulator of photosystem formation in *Rhodobacter sphaeroides*. *Microbiology* **149**: 377–388.
- Gruber, K., Puffer, B., and Krautler, B. (2011) Vitamin B₁₂-derivatives-enzyme cofactors and ligands of proteins and nucleic acids. *Chem Soc Rev* **40**: 4346–4363.
- Halpern, J. (1985) Mechanisms of coenzyme B₁₂-dependent rearrangements. *Science* **227**: 869–875.
- Han, Y.C., Meyer, M.H.F., Keusgen, M., and Klug, G. (2007) A haem cofactor is required for redox and light signalling by the AppA protein of *Rhodobacter sphaeroides*. *Mol Microbiol* **64**: 1090–1104.
- Hogenkamp, H.P. (1966) The photolysis of methylcobalamin. *Biochemistry* **5**: 417–422.
- Loo, J.A. (1997) Studying noncovalent protein complexes by

- electrospray ionization mass spectrometry. *Mass Spec-trom Rev* **16**: 1–23.
- Ludwig, M.L., and Matthews, R.G. (1997) Structure-based perspectives on B₁₂-dependent enzymes. *Annu Rev Biochem* **66**: 269–313.
- McGoldrick, H., Deery, E., Warren, M., and Heathcote, P. (2002) Cobalamin (vitamin B₁₂) biosynthesis in *Rhodobacter capsulatus*. *Biochem Soc Trans* **30**: 646–648.
- Mandal, M., and Breaker, R.R. (2004) Gene regulation by riboswitches. *Nat Rev Mol Cell Biol* **5**: 451–463.
- Masuda, S., and Bauer, C.E. (2002) AppA is a blue light photoreceptor that antirepresses photosynthesis gene expression in *Rhodobacter sphaeroides*. *Cell* **110**: 613–623.
- Masuda, S., and Bauer, C.E. (2004) Null mutation of HvrA compensates for loss of an essential *relA/spoT*-like gene in *Rhodobacter capsulatus*. *J Bacteriol* **186**: 235–239.
- Masuda, S., Dong, C., Swem, D., Setterdahl, A.T., Knaff, D.B., and Bauer, C.E. (2002) Repression of photosynthesis gene expression by formation of a disulfide bond in CrtJ. *Proc Natl Acad Sci USA* **99**: 7078–7083.
- Masuda, S., Berleman, J., Hasselbring, B.M., and Bauer, C.E. (2008) Regulation of aerobic photosystem synthesis in the purple bacterium *Rhodospirillum rubrum* by CrtJ and AerR. *Photochem Photobiol Sci* **7**: 1267–1272.
- Miller, V.L., and Mekalanos, J.J. (1988) A novel suicide vector and its use in construction of insertion mutations: osmoregulation of outer-membrane proteins and virulence determinants in *Vibrio cholerae* requires *toxR*. *J Bacteriol* **170**: 2575–2583.
- Moskvin, O.V., Kaplan, S., Gilles-Gonzalez, M.A., and Gomelsky, M. (2007) Novel heme-based oxygen sensor with a revealing evolutionary history. *J Biol Chem* **282**: 28740–28748.
- Moskvin, O.V., Gilles-Gonzalez, M.A., and Gomelsky, M. (2010) The PpaA/AerR regulators of photosynthesis gene expression from anoxygenic phototrophic proteobacteria contain heme-binding SCHIC domains. *J Bacteriol* **192**: 5253–5256.
- Nahvi, A., Sudarsan, N., Ebert, M.S., Zou, X., Brown, K.L., and Breaker, R.R. (2002) Genetic control by a metabolite binding mRNA. *Chem Biol* **9**: 1043.
- Nou, X., and Kadner, R.J. (2000) Adenosylcobalamin inhibits ribosome binding to *btuB* RNA. *Proc Natl Acad Sci USA* **97**: 7190–7195.
- Ortiz-Guerrero, J.M., Polanco, M.C., Murillo, F.J., Padmanabhan, S., and Elias-Arnanz, M. (2011) Light-dependent gene regulation by a coenzyme B₁₂-based photoreceptor. *Proc Natl Acad Sci USA* **108**: 7565–7570.
- Penfold, R.J., and Pemberton, J.M. (1991) A gene from the photosynthetic gene cluster of *Rhodobacter sphaeroides* induces *trans* suppression of bacteriochlorophyll and carotenoid levels in *R. sphaeroides* and *R. capsulatus*. *Curr Microbiol* **23**: 259–263.
- Penfold, R.J., and Pemberton, J.M. (1994) Sequencing, chromosomal inactivation, and functional expression in *Escherichia coli* of *ppsR*, a gene which represses carotenoid and bacteriochlorophyll synthesis in *Rhodobacter sphaeroides*. *J Bacteriol* **176**: 2869–2876.
- Perlman, D. (1959) Microbial synthesis of cobamides. *Adv Appl Microbiol* **1**: 87–122.
- Pfennig, N., and Lippert, K.D. (1966) Über das Vitamin B₁₂-Bedürfnis phototropher Schwefelbakterien. *Arch Mikrobiol* **55**: 245–256.
- Pollich, M., and Klug, G. (1995) Identification and sequence analysis of genes involved in late steps in cobalamin (vitamin B₁₂) synthesis in *Rhodobacter capsulatus*. *J Bacteriol* **177**: 4481–4487.
- Pollich, M., Jock, S., and Klug, G. (1993) Identification of a gene required for the oxygen-regulated formation of the photosynthetic apparatus of *Rhodobacter capsulatus*. *Mol Microbiol* **10**: 749–757.
- Ponnampalam, S.N., and Bauer, C.E. (1997) DNA binding characteristics of CrtJ. A redox-responding repressor of bacteriochlorophyll, carotenoid, and light harvesting-II gene expression in *Rhodobacter capsulatus*. *J Biol Chem* **272**: 18391–18396.
- Ponnampalam, S.N., Buggy, J.J., and Bauer, C.E. (1995) Characterization of an aerobic repressor that coordinately regulates bacteriochlorophyll, carotenoid, and light harvesting-II expression in *Rhodobacter capsulatus*. *J Bacteriol* **177**: 2990–2997.
- Ponnampalam, S.N., Elsen, S., and Bauer, C.E. (1998) Aerobic repression of the *Rhodobacter capsulatus bchC* promoter involves cooperative interactions between CrtJ bound to neighboring palindromes. *J Biol Chem* **273**: 30757–30761.
- Rodionov, D.A., Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. (2003) Comparative genomics of the vitamin B₁₂ metabolism and regulation in prokaryotes. *J Biol Chem* **278**: 41148–41159.
- Taylor, D.P., Cohen, S.N., Clark, W.G., and Marrs, B.L. (1983) Alignment of genetic and restriction maps of the photosynthesis region of the *Rhodospseudomonas capsulata* chromosome by a conjugation-mediated marker rescue technique. *J Bacteriol* **154**: 580–590.
- Waddington, M.D., and Finke, R.G. (1993) Neopentylcobalamin (NeopentylB₁₂) cobalt-carbon bond thermolysis products, kinetics, activation parameters, and bond dissociation energy: a chemical-model exhibiting 10⁶ of the 10¹² enzymatic activation of coenzyme B₁₂'s cobalt-carbon bond. *J Am Chem Soc* **115**: 4629–4640.
- Weissbach, H., Ladd, J.N., Volcani, B.E., Smyth, R.D., and Barker, H.A. (1960) Structure of the adenylobamide coenzyme: degradation by cyanide, acid, and light. *J Biol Chem* **235**: 1462–1473.
- Wienken, C.J., Baaske, P., Rothbauer, U., Braun, D., and Duhr, S. (2010) Protein-binding assays in biological liquids using microscale thermophoresis. *Nat Commun* **1**: 100. doi:10.1038/ncomms1093
- Willett, J., Smart, J.L., and Bauer, C.E. (2007) RegA control of bacteriochlorophyll and carotenoid synthesis in *Rhodobacter capsulatus*. *J Bacteriol* **189**: 7765–7773.
- Winkler, A., Heintz, U., Lindner, R., Reinstein, J., Shoeman, R.L., and Schlichting, I. (2013) A ternary AppA–PpsR–DNA complex mediates light regulation of photosynthesis-related gene expression. *Nat Struct Mol Biol* **20**: 859–867.
- Wuerges, J., Garau, G., Geremia, S., Fedosov, S.N., Petersen, T.E., and Randaccio, L. (2006) Structural basis

- for mammalian vitamin B₁₂ transport by transcobalamin. *Proc Natl Acad Sci USA* **103**: 4386–4391.
- Yin, L., Dragnea, V., and Bauer, C.E. (2012) PpsR, a regulator of heme and bacteriochlorophyll biosynthesis, is a heme-sensing protein. *J Biol Chem* **287**: 13850–13858.
- Yin, L., Dragnea, V., Feldman, G., Hammad, L.A., Karty, J.A., Dann, C.E., 3rd, and Bauer, C.E. (2013) Redox and

light control the heme-sensing activity of AppA. *MBio* **4**: e00563-13.

Supporting information

Additional supporting information may be found in the online version of this article at the publisher's web-site.